

KILGORE'S *Bred-Rite* SEEDS

*1957 Annual Catalog
for Florida*



SANFORD—300-2 W. First St.



WEST PALM BEACH
1310 Belvedere Rd.



PAHOKEE—313-19 Belle Glade Rd.




MIAMI—600 N.W. Fifth St.



PLANT CITY—101 N. Evers St.



GAINESVILLE—203 S.E. First Ave.



FORT MYERS—2216 Anderson Ave.



POMPANO BEACH
228 N. Flagler St.




HOMESTEAD—370 S. Flagler Ave.



GENERAL OFFICES, WAREHOUSE
AND MAIL ORDER DEPARTMENT
214 S. Collins St.



OCALA—909 N. Magnolia St.



BELLE GLADE—304-12 S. Main St.



WAUCHULA—702-6 W. Main St.



PALMETTO—710 - 13th St.



VERO BEACH—1109 20th St.

14 Kilgore Stores Serving Florida



THE KILGORE SEED COMPANY
General Offices and Mail Order Department, Plant City, Florida



TOMATO
JEFFERSON (Wilt Resistant)



PEPPER
CALIFORNIA WONDER



COLLARDS
GEORGIA 912



CABBAGE
MARION MARKET
(Yellows Resistant)



CELERY
FLORIDA GOLDEN
No. 15

PLANT CITY, FLORIDA

Exp. or Frt. Office _____ Date _____ 19____

AMOUNT ENCLOSED		
Check		
Money Order..		
Cash.....		
Total.....		

Notice of any defect or variation resulting from the failure of any seed to be as represented must be given within a reasonable time and not later than 15 days from the time such defect or variation becomes apparent and is observed by the purchaser. THE KILGORE SEED COMPANY. Section 578.26, Florida Statutes.

[illegible]

[SEE OTHER SIDE]

Tear Off Along This Line

PLEASE WRITE BELOW The names of neighbors or friends who might be interested in receiving a copy of our annual catalog.

NAMES	POST OFFICE	P.O. Box, Street, or R.F.D.	STATE
(SEE OTHER SIDE)			

Important Directions to Customers

By carefully observing the following directions, mistakes and misunderstandings will be very largely avoided.

HOW to order—Please be careful to sign your name, box number, street or rural route, post office and state on every order. Be sure to tell us your nearest express or freight office should you desire the seed or merchandise forwarded by express, freight or truck line. **Please write plainly.**

Cash with order or C. O. D.—Ours is a cash business and our prices are made on that basis. Customers are requested to remit when ordering, and by so doing save the c.o.d. fee charged by express companies, banks, and postal authorities for collecting and returning money.

Perishable items not sent C. O. D.—Perishable items, such as strawberry plants, rose bushes, seed potatoes, onion sets, etc., are not sent c.o.d. under any circumstances. Remittances must be sent with order.

How to send remittance—Remittances should be made by postal money orders, drafts, checks or cash. **Please do not send stamps.**

We pay postage to points in Florida—The postpaid prices indicated in this catalog include delivery to any point in Florida.

Prices subject to change—This catalog is printed during the early fall months. The market on many items of garden and field seeds, as well as supplies, has not yet been fully established. Consequently, prices on these items indicate value at time these pages are printed. In practically all cases orders will be filled at prices indicated in this catalog, **but all prices are subject to change without notice.** Write for special prices on larger quantities.

Export Shipments to Cuba and the West Indies—For the convenience of our customers in the West Indies, we have established the following agencies: E. G. Del Campo, Nueva Gerona, Isle of Pines, Cuba; F. C. Albury, Nassau, Bahamas.

Each order of pepper and tomato seed sent to Cuba and Dominican Republic requires an export certificate costing \$5.40.

For direct deliveries outside the continental United States, all prices quoted as "Postpaid" in this catalog are on an f.o.b. Plant City, Florida, basis, and the proper amount for postage must be added to the cost of the seeds.

Parcel Post rates to Cuba and the West Indies—The parcel post rate to Cuba, Bahamas and Jamaica is 45 cents, for Haiti 61 cents, and for the Dominican Republic 58 cents for the first pound or fraction thereof. If the net weight of an order is 1 pound or more, 1 pound must always be added to cover weight after being packed for shipment. In each case add 22 cents per pound for each additional pound or fraction thereof to the above indicated pound rates. Packages of merchandise going to Cuba, Haiti or Dominican Republic, weighing 8 ounces or less are subject to the postage rate of 1½ cents for each 2 ounces or fraction of 2 ounces. To the Bahamas and Jamaica, packages weighing 8 ounces or less are subject to the postage rate of 3 cents for each 2 ounces or fraction thereof, with a minimum charge of 15 cents.

WARRANTY AND LIMITATION OF LIABILITY—"We warrant that the seeds, bulbs and plants we sell are, at the time of delivery, as described on the container within recognized tolerances; but we limit our liability on this warranty to the amount of the purchase price of such seeds, bulbs and plants. We give no other or further warranty express or implied." The Kilgore Seed Company.

Notice of any defect or variation resulting from the failure of any seed to be as represented must be given within a reasonable time and not later than 15 days from the time such defect or variation becomes apparent and is observed by the purchaser. THE KILGORE SEED COMPANY. Section 578.26, Florida Statutes.

Air express shipments can now be made from Miami or Tampa, Florida, to any point in the West Indies.

Return of seeds for exchange—To safeguard our customers and ourselves, seeds may not be returned if seed package has been broken. In case of error on our part, unbroken packages may be returned within 10 days after shipment and we will gladly make correction at our expense.

Stock or lot number on all Kilgore's seed packages—Every lot of seed distributed by the Kilgore Seed Company will indicate (on one end of the front of the envelope or on the back of the label on canned seed, or with large seed like beans, peas, and corn on a tag attached to the bag), a stock number or lot number. This stock number is a code reference to our records of the history, performance and production of that particular lot of seed, by means of which we can trace it back to the growers' field where it was produced and to our trial plantings. It is recommended that every commercial grower of vegetables retain the empty packages or tags with the stock numbers. Correspondence pertaining to the performance of Kilgore's Bred-Rite seeds should include the specific lot number of the seed under consideration to aid investigation.

Postal, Express, and Sales Tax information—The parcel post weight limit for first and second zones is 40 pounds; 20 pounds in the third to eighth zones. Any weight up to 40 pounds in the first and second zones and 20 pounds in the third zone will usually go just as cheaply by parcel post as by freight or express.

In sections of Florida that are not easily accessible by freight or express, parcel post service will be found not only more satisfactory, but more economical. If on a rural route, delivery will be made to your door by mail.

Most Florida points are in the 1st (within 50 miles from Plant City, Fla.) or 2nd (within 50 to 150 miles from Plant City, Fla.) Postal Zone. Jacksonville north, Live Oak north and west, and points on the east coast south of West Palm Beach are in the 3rd Zone. (Within 150 to 300 miles from Plant City, Fla.) Puerto Rico and Virgin Islands are in the 6th Zone for surface mail; 8th Zone for Air Parcel Post. Surface mail 28c for 1st lb., 12c for each additional lb. Air Parcel Post 80c lb.

1st and 2nd ZONES				3rd ZONE			
1 lb.	.23	11 lbs.	.63	1 lb.	.23	11 lbs.	.75
2 lbs.	.27	12 "	.67	2 lbs.	.29	12 "	.80
3 "	.31	13 "	.71	3 "	.34	13 "	.85
4 "	.35	14 "	.75	4 "	.39	14 "	.90
5 "	.39	15 "	.79	5 "	.44	15 "	.96
6 "	.43	16 "	.83	6 "	.49	16 "	1.01
7 "	.47	17 "	.87	7 "	.54	17 "	1.06
8 "	.51	18 "	.91	8 "	.60	18 "	1.11
9 "	.55	19 "	.95	9 "	.65	19 "	1.16
10 "	.59	20 "	.99	10 "	.70	20 "	1.21

If in doubt about your zone rate, obtain it from your postmaster, and be sure to send enough money to cover necessary postage, if the items ordered are not indicated "postpaid" in this catalog. Do not have your order delayed or reduced by failing to add necessary money for postage. If you send too much money, it will be returned to you. It will be found plainly stated above each item whether or not it is postpaid. Also be sure to include with your remittance the State Sales Tax on non-exempt merchandise. (See Page 2).

When you order your seeds and supplies to come by express or freight we can send them collect, so you may pay the transportation charges on delivery. Give your express or freight station, if it is not the same as your post office.

SALES TAX INFORMATION

The Florida Revenue Act of 1949 (Sales and Use Tax) imposes a 3% tax on the retail sales of many commodities. Numerous agricultural supply items are exempt for the tax, and for your guidance in determining tax exempt items shown in this catalog, we list below those items that are not subject to the tax.

EXEMPT ITEMS—Beekeeper's Supplies (except Hive Bodies and Frame).

Cans and Jars
Crate Materials
Fencing—For farm use
Feeds—For farm animals
Grafting Wax and Rubber Strips
Hotkaps
Label Paste
Peat Moss
Poultry and Stock Remedies
Paper Pots—Plant Bands and Flats

Seeds—All Vegetable, Field Crop, Pasture
Grass Seeds, Onion Sets, Seed Potatoes
Seedbed Cloth
Seed Inoculants
Seed Treating Compounds
Starter Solutions and Root Inducing Materials
Terra-Lite Tractors
Weed Killers

EXEMPT ONLY IF USED FOR COMMERCIAL AGRICULTURAL PURPOSES

Dusters
Farm Tools
Fertilizers
Insecticides and Fungicides
Soluble Plant Foods

Fumigants
Garden Tools
Raffia and Twine
Rubber Brands
Sprayers
Tarpaulins

All flower seed and other merchandise listed in this catalog is subject to 3% State Sales Tax except sales to Dealers, Government Agencies, Schools, Churches and Charitable organizations provided Tax Exemption Certificate has been obtained from the State Comptroller. All orders from institutions or organizations claiming tax exemption must clearly indicate on order sent us, the certificate serial number and date of certificate.

THE RATE OF TAX APPLICABLE TO NON-EXEMPT SALES IS AS FOLLOWS:

- (1) On single sales of 10c or less no amount shall be added for taxes.
- (2) On single sales in amounts for 11c to 35c, both inclusive, 1c shall be added for taxes.
- (3) On sales in amounts from 35c to 65c, both inclusive, 2c shall be added for taxes.
- (4) On sales in amounts from 65c to \$1.00, both inclusive, 3c shall be added for taxes.
- (5) On sales in amounts of more than \$1.00, 3% shall be charged upon each dollar of price, plus the above bracket charges upon any fractional part of a dollar in excess of even dollars.

SPECIAL NOTICE—In order to avoid delay in filling your order, on all non-exempt items, be sure to include with your remittance the proper amount of sales tax as indicated above. We are not permitted by law to assume the Sales Tax.

FLORIDA HOME VEGETABLE GARDENS IN 1957

Florida is one of the most important commercial vegetable producing states of the Union. Florida home vegetable gardens offer almost no competition to commercial vegetables which are grown mainly for shipment to northern markets. Many Florida families who do not get fresh vegetables from their own gardens, seldom get them, at least in sufficient quantity for good health, for the market. Florida people, city residents as well as rural folks, can help solve their food problems by producing vegetables in the home garden.

Florida climate and soils are favorable for producing the finest vegetables in the United States. In Flor-

ida fresh vegetables can be grown every month in the year, especially during the fall, winter and spring months when they cannot be grown in other sections of the United States. In addition to producing fresh vegetables in your Florida home garden, plans should also be made for preserving vegetables from the garden by canning or freezing for home use, which will mean not only better health for the family, but also a material saving in cost. For best results, order your seeds from the Kilgore Seed Company — Florida's Leading Seedsmen, Mail Order Department, Plant City, Florida.



Location and plan of the vegetable garden

A level, well drained piece of ground, convenient to the house should be selected for the garden. It should be located where it can be irrigated if necessary and away from hedges, large trees or the shade of buildings. On the farm it should be well-fenced to protect it from rabbits, chickens and other animals.

After selection of a location, and determining the size of the garden, a list of vegetables should be made. In selecting varieties, primary consideration should be given to the likes of the family, also to the size of the plot available. Different vegetables give variety to the diet.

Making a seedbed

Small seeds such as lettuce, cabbage, collard, pepper, eggplant, tomato, etc., should be sown in a seedbed and the young plants transplanted to the field or garden. A shallow box or flat may be used for this purpose, or a small plot in the garden may be used as a seedbed for starting plants for transplanting. The soil should be in perfect condition. Make seedbeds four to six inches high, to prevent overflow in case of heavy rains. In preparing seedbed soil it is desirable to mix in "Terra-Lite" (Vermiculite) in order to increase germination of seed and prevent damping off of seedlings. (See page 74). Fertilize with 2 to 3 pounds per 100 sq. ft. of a commercial fertilizer such as Vigoro or Vertagreen (page 67), applied at least a week or ten days before sowing seeds. Sow the seed in rows about six inches apart. Very fine seed like celery, should not be covered with soil, but may be pressed into the soil with a light roller or with a wide board. Larger seeds like tomato, pepper, and cabbage may be covered, not over $\frac{1}{4}$ to $\frac{1}{2}$ inch deep.

We recommend the use of burlap or white muslin as a cover over the seedbed, stretching it on a three- or four-foot wide tent-shaped or slanting frame built over the seedbed, with the sides about 10 to 12 inches from the ground to provide ample circulation of air under the cover. If plants are being started in late

A map or plan of the garden should be made, showing the various vegetables to be grown, and their position in the garden, indicating length of rows and the space between rows. In case of a small, backyard garden, entirely worked by hand, the rows should be close together, i.e., the smaller figure in column 5 of our chart on page 4, so that large quantities of vegetables may be produced on a limited space.

The planting data given on page 4 of this catalog will help greatly in preparing a plan since seed requirements, planting depths, distances and time of maturity are given.

summer for fall planting, the seedbed cover will serve as a shade to protect seedlings from the hot sun, also to break the force of heavy rains. If plants are started in winter for a spring crop, this seedbed cover will protect seedlings from frost or freezes. It may be also advisable to use ground covers of burlap or old fertilizer bags. When seeds begin to germinate these ground covers should be removed immediately to prevent spindly plants. After the seed is sown, it is desirable to sprinkle every evening or oftener if necessary with cool water by means of hand sprinkling pots.

Success in getting transplanted plants to grow satisfactorily depends very much upon how the plants are grown in the seedbed. For a week or ten days before transplanting, the plants in seedbed should be gradually hardened. This means leaving the cover off the plants for a week or so before transplanting, and water should be withheld from the plants during this period. When plants are about six inches high they are ready to transplant into the field or garden, preferably on a cloudy day or toward evening. Water the plants thoroughly several hours before removing from seedbeds, and again immediately after setting. A starter solution is helpful (see page 67). To grow good vegetables, a large supply of moisture and fertilizer is essential until the crop is ready to harvest.

Preparation of the soil for the vegetable garden

The soil in the garden should be well prepared. Grass and weeds removed with all roots. A little extra time spent in preparing the soil for planting will save many hours of hoeing later. It is best to spade or plow, and apply commercial fertilizer to the soil ten days to two weeks before planting in order to avoid burning the seed or young plants. A sample of the soil should be tested, either by yourself with an inexpensive soil test kit (page 75) or by your county agent, and if found too acid or sour apply Hardwood Ashes or Hydrated Lime; if too alkaline or sweet apply Aluminum Sulphate and Manganese Sulphate (see page 68).

Dairy Organic Compost is excellent for fertilizer and land conditioner, will not burn, is odorless and contains no weed seed. Use at the rate of 20 to 25 lbs. per 100 sq. ft. broadcast. See page 67. If available, leaf mold or black muck or hammock soil are ideal materials to work into soil. If none of these materials is available, a two or three inch layer of horticultural grade peat moss (page 74) may be worked into the garden soil. This material is clean and easy to handle, and is wonderfully effective for breaking up hard soil, making it mellow, or for adding humus to sandy soil, making it more retentive of moisture and fertilizer materials. After the peat moss is worked into the soil, an inch

or two inch layer of Pulverized Sheep Manure, or Dairy Organic Compost, along with a liberal application of a good garden fertilizer such as Vigoro or Vertagreen (see page 67) should be used, applying one of these fertilizer materials broadcast at the rate of 4 to 6 lbs. per 100 square feet. If this material is raked into the surface soil a week or ten days before planting, and if well watered in case of no rain, in order to speed the incorporation and decomposition of the fertilizer or manure, you will have the foundation for an excellent vegetable garden. If the fertilizers are applied just before planting, they are very apt to burn the young seedlings as they emerge from the seed.

After the vegetable plants are well started in the garden, fertilizer should be applied lightly every two or three weeks on either side of the row of vegetables, but not in contact with the leaves or stems to avoid burning.

Note:—All materials indicated above, such as Pulverized Sheep Manure, Dairy Organic Compost, Peat Moss, Terra-Lite, Super Phosphate, Vigoro, Vertagreen, Wood Ashes, Lime, Aluminum Sulphate, Manganese Sulphate, etc., are available at all Kilgore seed stores, and in our mail order department at Plant City, Florida.

For further information on Florida home gardening write the Florida Agricultural Experiment Station, Mailing Department, Gainesville, Florida, for a free copy of Bulletin 131 entitled "The Florida Home Garden," and also for Circular 104 entitled "Vegetable Garden Production Guide."

Reference Table for Florida Vegetable Gardens

Many mistakes may be avoided by following directions given in this table. The time requirements indicated for crops to reach edible size vary, of course, according to the season, varieties and soil conditions. Allowance also must be made for the relative estimate of the yields, due to climatic and soil conditions. However, the aver-

ages indicated in the table will be found to be fairly accurate for Florida growing conditions on the basis of many years of observation and experience. **For information on "when to sow vegetable seeds in Florida" see page 5.**

Kind of vegetable Months to Sow Seed (Inclusive)**	Number Running feet of row for a family of 3 or 4 people (Includes for canning)	Seeds or plants to provide for a family of 3 or 4 people	Depth to cover in inches	Space between rows in feet	Set plants from seedbed or thin out to inches in row	Approximate time required to produce crop, from seed planted in garden or from transplanting plants to garden (days)
Beans (Bush) 1-4, 9-10	100*	1/2 lb.†	1 1/2—2	2—2 1/2	3—4	48 to 56
Beans (Pole) 3-5	50	1/4 lb.	1 1/2—2	3—4	9 or 3 ft. hills	63 to 70
Beans (Bush Lima) 3	100*	1/2 to 1 lb.†	1 1/2—2	3	4—5	65 to 75
Beans (Pole Lima) 3	50	1/2 lb.	1 1/2—2	3—4	9 or 3 ft. hills	90
Beet 1-3, 9-12	100*	1 oz.†	1/2—3/4	1 1/2—2	3—4	56 to 68
Broccoli 8-12	40	1 pkt. or 25 plts.	1/4—1/2	3	18—24	55 to 60 from plts.
Brussels Sprouts 1-2, 8-12	25	15 plants	1/4—1/2	2 1/2—3	18—24	90 from plants
Cabbage 1-3, 8-12	60	50 plants	1/4—1/2	2 1/2—3	12—16	62 to 98 from plts.
Chinese Cabbage 1-2, 8-12	40*	1 pkt. or 50 plts.†	1/4—1/2	2 1/2—3	8—12	70 from seed
Cantaloupe 1-4	50	1 pkt.†	1/2—3/4	4—5	24 or 4 ft. hills	75 to 95
Carrot 1-3, 9-12	100*	2 pkts.†	1/4—1/2	1 1/2—2	2—3	72 to 95
Cauliflower 1-2, 8-12	50	30 plants	1/4—1/2	3—4	18—24	55 to 62 from plts.
Celery 7-12	50	150 plants	1/8	3	4—5	105 to 130 from plts.
Collard Every Month	50*	1 pkt. or 35 plts.†	1/4—1/2	2 1/2—3	14—18	50 to 52 from plts.
Corn (Sweet) 1-3	200*	1/4 lb.†	1—2	2 1/2—3	12—15	65 to 80
Corn (Roasting Ear) 1-3	200*	1/4 lb.†	1—2	3—3 1/2	15—18	68 to 87
Cowpeas (Table) 3-8	150	1/2 to 3/4 lb.	1—1 1/2	3—4	5—6	68 to 85
Cucumber 1-3, 8-9	50	1 pkt.	3/4	5—6	12 to 24	40 to 50
Eggplant 1-3, 6-8	50	17 plants	1/2	4—4 1/2	18 to 36	80 to 85 from plts.
Endive (Escarolle) 1-2, 8-12	40	pkt. or 50 plts.	1/4—1/2	1 1/2—2	9	90 to 95 from seed
Garlic 8-11 (bulbs or sets)	30	1 lb.	1	1 1/2	4	90
Kale 1-3, 8-12	25	1 pkt.	1/4—1/2	2 1/2	20—24	70
Kohl Rabi 1-3, 9-12	25*	1 pkt.†	1/4—1/2	1 1/2—2	6—8	50
Leek 1-2, 9-12	50	1 pkt. or 120 plts.	1/2	1 1/2	5—6	140 from seed
Lettuce (Head) 1-2, 8-12	75*	1 pkt. or 90 plts.†	1/4—1/2	1 1/2—2	10—15	66 to 84 from seed
Lettuce (Leaf) 1-3, 8-12	50*	1 pkt. or 60 plts.†	1/4—1/2	1 1/2—2	10—12	46 from seed
Mustard 1-12	40*	1 pkt.†	1/4—1/2	2	8—10	40 to 45
Okra 3-9	75	1 oz.	1—1 1/2	3—4	10—15	50 to 60
Onion Seed 8-11	80	1 pkt. or 400 plts.	1/2	1 1/2	2—4	110 to 150 from seed
Onion Sets 1-2, 8-12	40	1 pt.	1	1 1/4—1 1/2	1—2	30 to 60
Parsley 1-2, 9-12	30	1 pkt.	1/4—1/2	1 1/2—2	10—12	90
Parsnip 1-2, 9-12	50	1 pkt.	1/2	2 1/2—3	4	120
Peas (English) 1-2, 9-12	100*	1/2 lb.†	1 1/2—2	3	1—2	50 to 53
Pepper 1-2, 6-8, 11-12	40	24 plants	1/2	2 1/2—3 1/2	10—16	60 to 80 from plts.
Potatoes (Irish) 1-3, 8-12	200	12 lbs.	1—4	3—3 1/2	9—12	75 to 90
Pumpkin 3-5	50	1/2 oz.	1	8—10	5 ft. hills	100 to 118
Radish 1-4, 8-12	25*	1 pkt.†	1/2	1—1 1/2	1—2	21 to 25
Rhubarb 9-11	25	pkt. or 15 plts.	1/2—3/4	2 1/2—3	18—24	90 from plants
Roselle 3-4	25	1 pkt.	1/2—3/4	5—6	24—36	175
Rutabaga 1-2, 8-12	50	1 pkt.	1/4—1/2	2	5—6	90
Spinach—Savoy 1-2, 8-12	50*	1 pkt.†	1/2—3/4	1 1/4—1 1/2	6	40
Spinach—New Zealand 3-7	25	1 pkt.	3/4—1	3	20	60
Squash (Bush) 1-3, 8-9	50	1 pkt.	1	3—4	2 ft. hills	42 to 52
Squash (Running) 1-2	50	1 pkt.	1	6—8	3 or 4 ft. hills	60 to 115
Swiss Chard 1-3, 9-12	25	1 pkt.	1/2—1	2	12—15	58 to 60
Tomato—Ground 1-3, 8-9, 12	150	50 plants	1/2	3 1/2—5	24—36	70 to 90 from plts.
Tomato—Staked 1-3, 8-9, 12	75	50 plants	1/2	4	16—20	70 to 90 from plts.
Turnip 1-4, 7-12	50	1 pkt.†	1/4—1/2	1 1/4—1 1/2	4—6	35 to 50
Watermelon 1-4	75	1 pkt.	1/2—1	8—10	6 to 8 ft. hills	85 to 100

* Make two or more plantings at different times during the season. Number of feet of row indicated is for each planting.

† Number of feet and amount of seed or number of plants indicated are for each planting.

** Months indicated are for central and south Florida. In north Florida, some tender kinds will not stand the winter cold, and this fact should be taken into consideration with reference to time of sowing seed.

THE KILGORE SEED COMPANY—A Florida Institution

GENERAL OFFICES AND MAIL ORDER DEPARTMENT, PLANT CITY, FLORIDA

Fourteen Kilgore seed stores serving Florida, located at:

Belle Glade
Fort Myers
Gainesville
Homestead

Miami —
600 N.W. Fifth St.
Ocala
Pahokee

Palmetto
Plant City
Pompano Beach
Sanford

Vero Beach
Wauchula
West Palm Beach
1310 Belvedere Road

What and When to Sow Vegetable Seeds in Florida

JANUARY and FEBRUARY

Beans	Corn, Sweet	Onion Seed	Rutabaga
Beet	Cucumber	Onion Sets	Spinach
Brussels	Eggplant	Parsley	Squash
Sprouts	Endive	Parsnip	Swiss Chard
Cabbage	Kale	Peas, English	Tomato
Cantaloupe	Kohl-Rabi	Pepper	Turnip
Carrot	Leek	Potatoes	Watermelon
Cauliflower	Lettuce	Radish	
Collard	Mustard	Romaine	

MARCH

Beans	Corn, Sweet	Okra	Squash
Beet	Cucumbers	Potatoes	Swiss Chard
Cabbage	Kale	Pumpkin	Tomato
Cantaloupe	Kohl Rabi	Radish	Turnip
Carrot	Lettuce	Roselle	Watermelon
Collard	Mustard	Spinach	

APRIL

Beans	Mustard	Radish	Turnip
Cantaloupe	Okra	Roselle	Watermelon
Collard	Pumpkin	Spinach	

MAY

Pole Beans	Mustard	Pumpkin	
Collard	Okra	Spinach	

JUNE

Collard	Mustard	Pepper	Spinach
Eggplant	Okra	Rutabaga	Turnip

JULY

Celery	Mustard	Rutabaga	
Collard	Okra	Spinach	
Eggplant	Pepper	Turnip	

AUGUST

Broccoli	Collard	Lettuce	Romaine
Brussels	Cucumber	Mustard	Rutabaga
Sprouts	Eggplant	Okra	Spinach
Cabbage	Endive	Onion Sets	Squash
Cauliflower	Garlic	Pepper	Tomato
Celery	Kale	Radish	Turnip

SEPTEMBER

Bush Beans	Collard	Okra	Rutabaga
Beet	Cucumber	Onion Seed	Spinach
Broccoli	Endive	Onion Sets	Squash
Brussels	Garlic	Parsley	Swiss Chard
Sprouts	Kale	Parsnip	Tomato
Cabbage	Kohl Rabi	Peas—English	Turnip
Carrot	Leek	Radish	
Cauliflower	Lettuce	Rhubarb	
Celery	Mustard	Romaine	

OCTOBER

Bush Beans	Celery	Mustard	Rhubarb
Beet	Collard	Onion Seed	Romaine
Broccoli	Endive	Onion Sets	Rutabaga
Brussels	Garlic	Parsley	Spinach
Sprouts	Kale	Parsnip	Swiss Chard
Cabbage	Kohl Rabi	Peas, English	Turnip
Carrot	Leek	Potatoes	
Cauliflower	Lettuce	Radish	

NOVEMBER

Beet	Collard	Mustard	Rhubarb
Broccoli	Eggplant	Onion Seed	Romaine
Brussels	Endive	Onion Sets	Rutabaga
Sprouts	Garlic	Parsley	Spinach
Cabbage	Kale	Peas, English	Swiss Chard
Carrot	Kohl Rabi	Pepper	Turnip
Cauliflower	Leek	Potatoes	
Celery	Lettuce	Radish	

DECEMBER

Beet	Collard	Onion Seed	Romaine
Broccoli	Eggplant	Onion Sets	Rutabaga
Brussels	Endive	Parsley	Spinach
Sprouts	Kale	Parsnip	Swiss Chard
Cabbage	Kohl Rabi	Peas, English	Tomato
Carrot	Leek	Pepper	Turnip
Cauliflower	Lettuce	Potatoes	
Celery	Mustard	Radish	

What and When to sow grasses, clovers, and field crop seeds in Florida

JANUARY and FEBRUARY

Alfalfa	Bahia Grass	Rape
Clover, Crimson	Oats	Rye

MARCH

Beans, Velvet	Italian Rye, Sudan, Bahia	Peanuts
Benne		Rape
Corn, Field	Indigo	Rice
Crotalaria	Hegari	Egyptian Wheat
Grasses	Millet	Sorghum
Bermuda, Carpet,	Peas	Sunflower

APRIL

Beans, Velvet	Grasses	Peanuts
Beggarweed	Bermuda, Carpet,	Rice
Benne	Italian Rye, Sudan, Bahia	Egyptian Wheat
Chufas	Indigo	Sorghum
Clover, Alyce	Hegari	Sunflower
Corn, Field	Millet	
Crotalaria	Peas	

MAY

Beans, Velvet	Grasses	Peas
Beggarweed	Bermuda, Carpet,	Peanuts
Benne	Italian Rye, Sudan, Bahia	Sesbania
Chufas		Egyptian Wheat
Clover, Alyce	Indigo	Sorghum
Corn, Field	Hegari	Sunflower
Crotalaria	Millet	

JUNE

Beans, Velvet	Crotalaria	Peanuts
Beggarweed	Grasses	Rice
Benne	Bermuda, Carpet, Sudan, Bahia	Sesbania
Chufas		Egyptian Wheat
Clover, Alyce	Indigo	Sorghum
Corn, Chicken	Hegari	Sunflower
Cowpeas	Millet	

JULY

Beans, Velvet	Cowpeas	Hegari
Beggarweed	Grasses	Millet
Chufas	Bermuda, Sudan and Bahia	
Corn		

AUGUST

Cowpeas	Bermuda, Sudan and Bahia	Millet
Grasses		Sunflower

SEPTEMBER

Clover, Sweet	Grasses	Lupines
Kenland Red	Bermuda, Bahia	Rape
		Rye

OCTOBER

Clover, Sweet	Grasses	Lupines
Kenland Red	Bermuda, Italian Rye, Carpet, Bahia	Oats
		Rape
		Rye

NOVEMBER

Clover	Grasses	Oats
Crimson, Sweet, White Dutch, Ladin, Kenland Red	Bermuda, Italian Rye, Carpet, Bahia	Rape
	Lupines	Rye

DECEMBER

Clover	Grasses	Oats
Crimson, Sweet, White Dutch, Ladin, Kenland Red	Italian Rye, Carpet, Bahia	Rape
	Lupines	Rye

See pages 38 to 47 for complete cultural directions on field corn, grasses, clovers and field crops in Florida.
See page 48 when to sow flower seed in Florida.

BEANS

Our bean seed is highly germinating, guaranteed western-grown stock, from the best western growing areas, all carefully milled, graded and fitted for the critical grower.

See page 34 for details of the Kilgore method of handling seeds

Culture of Green and Wax Podded Bush Beans. Beans are grown on all types of soil in Florida, and are not very particular with reference to soil requirements. Beans are a very quick maturing crop.

Spring planting in the southern and central part of the State begins January 1 and continues until April. Fall planting starts in August. For a fall crop, plant from the middle of August until the middle of October; for a main crop on the Florida east coast plant from October to December. Sow seed in drills, 2 to 2½ feet apart, dropping a bean every 2 to 2½ inches, and cover 1½ to 2 inches deep. On muck soils apply 300 to 600 pounds of a 3-8-8 fertilizer mixture per acre. On sandy loam apply 1200 to 1600 pounds of a 4-7-5 fertilizer per acre, making one application. The fertilizer put out from a week to ten days before planting. There is

always a risk of the fertilizer burning and killing the vitality of the seed when both are put in at the same time.

For the control of insects and diseases of beans see page 53. In order to control soil diseases and get a better stand, it is always desirable to treat bean seed before planting. (See pages 52 and 62). This often means the difference between success and failure in getting a stand. For control of watery soft rot or white mold use Granular Aero Cyanamid.

Inoculate bean seed with Nitragin or A.B.C. Inoculant. This hastens maturity, increases yields and enriches the soil. (See page 69).

The number of days in parentheses after each variety represents average time to maturity.

Green Podded Bush Beans

Plant ½ lb. to 100 feet of row at each planting

Plant 3 pecks (45 lbs.) to 1 bushel (60 lbs.) per acre. In the garden make four plantings at different times.

Write for special prices on larger quantities



Contender (Mosaic Resistant) Bush Beans

CONTENDER (Mosaic Resistant). (45 days.) A consistently early, high yielding, attractive appearing, stringless bean, adapted for shipment to fresh markets. Plant is very vigorous and produces pods similar to Black Valentine, although the Contender pods will average three-fourths of an inch longer, are slightly heavier and thicker, are round to oval, and have a tendency to curve slightly.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$3.85; 60 lbs. (bu.) \$14.40

ASGROW BLACK VALENTINE. (48 days.) The plant is large, vigorous, erect and very prolific. The oval pods are nearly straight, medium green, entirely stringless, and of fine quality and flavor. They retain their color and texture in shipping long distances. Very attractive in appearance.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$4.15; 60 lbs. (bu.) \$15.60

GLADES (53 days.) Fully resistant to common bean mosaic and the variant strain NY 15. The plant is vigorous, tall and upright, holding the pods well off the ground. The pods are thick-flat to flat oval in shape, averaging longer and narrower than Plentiful and a darker green color. Compared with Asgrow Black Valentine the plants are taller, and the pods longer and slightly wider. Very attractive pods, a good shipper and heavy yielder.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25; pk. \$4.90; bu. \$18.60

SEMINOLE. (53 days.) A round podded bush bean released by the Everglades Experiment Station of Florida A.E.S. The plant is somewhat sprawling, with nu-

merous leaves that protect the pods from wind-scars. Seminole has yielded well in widely scattered trials. The pods are similar to Tendergreen in length, slightly lighter in color. Pods are stringless, fiberless and of good flavor, texture, quality and appearance.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$5.60; 60 lbs. (bu.) \$21.30

TENDERGREEN. (50 days.) This variety is popular due to its excellent shipping quality. Plants are large, erect, and thick-stemmed. The pods average about five inches in length, are attractive in appearance, being nearly straight, round, fleshy, dark green, stringless, and of excellent quality.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$4.40; 60 lbs. (bu.) \$16.50

TENDERLONG 15 (Improved Tendergreen Type). (50 days.) Developed for disease resistance; longer, straighter pods than Tendergreen. The plant is an erect, compact bush, holding the pods well off the ground. Highly productive and resistant to common bean mosaic and several other virus diseases, one form of powdery mildew; with considerable tolerance to Halo blight. The pods, averaging six inches, stringless and fiberless and of excellent flavor and medium dark green color.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$4.40; 60 lbs. (bu.) \$16.50

RESISTANT TENDERGREEN (50 days.) Similar to regular Tendergreen in bush and pod type, but resistant to common bean mosaic, and will produce longer pods and higher yields than regular Tendergreen when the disease is present.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25; pk. \$4.40; bu. \$16.50



Tenderlong 15

Green Podded Bush Beans (Continued on next page)

BEANS

Green Podded Bush Beans (Cont'd)

PLENTIFUL. (50 days.) Plants are spreading in growth and should be given a little wider spacing in the row than most bush beans. The pods are long, flat, straight, and of medium green color. Heavy yielder.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$4.15; 60 lbs. (bu.) \$15.60


WADE (B1515) (Mosaic Resistant) (48 days.) The vigorous plants are tall and hold the pods well off the ground. The heavy foliage helps to protect the pods from wind-scars. It is a fleshy, round podded bean of the Tendergreen type, but powdery mildew and mosaic resistant, it has a consistently higher yield and produces long, straight pods. The pods are of a dark green color and hold their green color exceptionally well in transit and on the market. Stringless and little fiber. Good for market as well as for processing. Because of its long bearing season, high quality and disease resistance, it is highly recommended for home gardens and is a good shipper.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25; Postpaid
Not prepaid, 15 lbs. (pk.) \$4.90; 60 lbs. (bu.) \$18.60

WAX PODDED BUSH BEANS

Plant 3 pecks (45 lbs.) to bushel (60 lbs.) per acre.
In the garden plant ½ lb. to 100 feet.

Write for special prices on larger quantities.



Cherokee (Valentine Wax)

CHEROKEE (Valentine Wax). (50 days.) Plant large, vigorous, erect and prolific. Early and an enormous yielder of beautiful, attractive appearing, golden-yellow, long, thick oval pods, nearly straight, entirely stringless and of excellent quality. Considered the best wax bean for Florida and the South.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$4.30; 60 lbs. (bu.) \$16.20

GREEN PODDED POLE BEANS

Plant 2 pecks (30 lbs.) per acre.
In garden ¼ pound to 50 feet.

Culture of Green Podded Pole Beans. Pole beans are later in maturing than bush beans, but bear for a longer season and are extremely productive, yielding more than bush beans, and also produce bigger pods. They are somewhat more tender than bush beans if harvested young. Pole beans should be supported by wires or by good strong poles to keep them off the ground. Poles should be 8 to 9 feet above ground, and set firmly in the ground, in rows 3 to 4 feet apart, and 2 to 3 feet apart in the row. In order to keep pole beans bearing over a long season, it is important to keep them closely picked. Inoculate bean seed with Nitragin or A.B.C. Inoculant. (See page 69.) Pole beans are more susceptible to mildew, rust and other foliage diseases than are bush beans, and therefore should be sprayed or dusted quite often. (See page 53.)

Write for special prices on larger quantities.

FLORIGREEN (Rust Resistant). A new pole bean developed by Dr. A. P. Lorz, horticulturist at the Florida Experiment Station, Gainesville, Fla., and Dr. J. M. Walters, Pathologist of the Gulf Coast Station, Bradenton, Fla.

Florigreen is highly resistant to rust. Also resistant to common and southern bean mosaics. The pods resemble U. S. No. 4 or 191, but a little darker. They are long, straight and retain a fresh dark green color, which is highly desirable for fresh market, stringless at the proper stage for harvesting. Florigreen has consistently outyielded U. S. No. 4 and 191 in experimental trials. This new pole bean looks very promising. All bean growers should give it trial.

Seed supply limited this season.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25; (pk.) \$5.65; (bu.) \$21.60

U. S. NO. 4 (No. 191) (WHITE KENTUCKY WONDER). (63 days.) A high yielding type of white-seeded Kentucky Wonder, resistant to certain forms of rust. Plant is vigorous, with dark green leaves. The pods are dark green, oval, straight, long, averaging from 8 to 9 inches, and attractive in appearance. Stringless in the marketable stage. Productive and very desirable for market or home use. Grown most extensively on the west coast of Florida for shipment to southern and northern markets.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$4.45; 60 lbs. (bu.) \$16.80

KENTUCKY WONDER (OLD HOMESTEAD). (65 days.) This pole bean is early and very productive over a long season. Pods are very long, ranging from eight to ten inches, curved, nearly round. Pods are meaty, very brittle, and of good quality if harvested young. The pods hang in large clusters from the top to the bottom of the vine.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$4.45; 60 lbs. (bu.) \$16.80

McCASLAN. (65 days.) A vigorous grower and heavy producer, unlike other varieties, producing from the ground up over a long season. The pods are medium green, eight inches long, large, flat, slightly curved, thick and meaty, stringless, brittle, fine grained, and of excellent quality if harvested young. Because of the high yield of large, attractive pods, this variety has become the leading pole bean on the lower east coast of Florida for shipment to northern markets.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid
Not prepaid, 15 lbs. (pk.) \$4.60; 60 lbs. (bu.) \$17.40



McCaslan pole beans as grown from Kilgore's high vitality seed in the Homestead section.

LIMA BEANS

BUSH LIMA BEANS

Fordhook—Plant 1 bushel (60 lbs.) per acre.

Henderson's—Plant 1 to 2 pecks (15 to 30 lbs.) per acre.

In the garden make two plantings at different times. With Fordhook plant 1 pound to 100 feet or with Henderson's plant ½ pound to 100 feet.

Culture of Bush Lima Beans. In general, cultural requirements of Bush Lima Beans are similar to those given for green and wax podded bush beans, but they need more room. Lima beans are not so hardy as string beans, and the ground should be warm when planting the seed. Fertilizer should be applied at least ten days before planting. In order to control soil diseases and get a better stand, it is always desirable to treat lima bean seed before planting (see pages 52 and 62). Plant in rows three feet apart, dropping a bean every 3 to 4 inches in the row, covering an inch deep.

Write for special prices on larger quantities.

FORDHOOK 242. (72 days.) It is an excellent bearer, producing a heavy, early yield, and is especially suitable to Florida because of its ability to make a heavy set and yield under high temperature conditions. The plants are as tall as regular Fordhook, grow more erect and have a more open appearance. Pods average 3 to 4 inches long by 1 and 1½ inches wide and about ½ inch thick, and contain 3 to 4 thick beans. Green shell beans are very slightly smaller than regular Fordhook, have light green seed coats, and are of delicious, finely flavored edible quality.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid

Not prepaid, 15 lbs. (pk.) \$4.70; 60 lbs. (bu.) \$17.70

FORDHOOK-CONCENTRATED. (75 days.) Developed for concentration of pods on the plant, making it easy to harvest, and for concentration in time of reaching maturity. The plants are stocky, without tendrils and very prolific. Pods are dark green, very large, with thick fibrous walls giving them body for long distance shipments. Pods contain 3 to 4 large, very thick, plump beans of excellent quality.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid

Not prepaid, 15 lbs. (pk.) \$4.70; 60 lbs. (bu.) \$17.70



Jackson Wonder (Old Florida Bush Speckled Butter Bean)

HENDERSON'S BUSH LIMA (Baby Lima). (65 days.) An early variety, hardy and a sure cropper. The plants are small and very productive. The pods are small in size, very short and narrow, thin, flat, straight to slightly curved. The seeds are smaller and not as plump as Fordhook. This variety matures about ten days earlier than Fordhook.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid

Not prepaid, 15 lbs. (pk.) \$3.70; 60 lbs. (bu.) \$13.80

Butter or Green Shell Beans

Used principally as Green Shell Beans, like Lima Beans

Old Florida Pole—Plant 1 peck (15 lbs.) per acre.

Jackson Wonder Bush—Plant 1 to 2 pecks per acre.

French's Horticultural—Plant 1 to 2 pecks per acre.

In the garden plant ¾ pound Bush to 100 feet,

½ pound Pole to 100 feet.

Write for special prices on larger quantities.

FLORIDA POLE SPECKLED BUTTER BEAN (Old Florida Pole). (75 days.) Culture similar to Green Podded Pole Beans (See page 7). Plant is vigorous, and very prolific over a long bearing season. Pods, borne in clusters, are of medium size, flat, containing an average of 3 medium-sized, flat seeds. Seeds are of buff color spotted with reddish brown. A dependable, prolific variety for Florida.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid

Not prepaid, 15 lbs. (pk.) \$4.45; 60 lbs. (bu.) \$16.80

JACKSON WONDER (Old Florida Bush Speckled Butter Bean). (65 days.) Culture similar to Bush Lima Beans. This variety is similar to Old Florida Pole, but produces a bush or dwarf plant. Plant is small to medium in size, spreading, with semi-runners resembling somewhat Henderson's Bush Lima. Pods are flat, containing 3 or 4 beans. The seeds are small, flat and of light buff color. Early and very prolific, drought-resistant, a sure cropper and of good quality.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid

Not prepaid, 15 lbs. (pk.) \$3.70; 60 lbs. (bu.) \$13.80

FRENCH'S HORTICULTURAL (68 days.) This is the most attractive bean of the speckled Cranberry type making long pods (6 to 8 inches), straight very heavily splashed with bright red or deep carmine, which gives them a very striking and attractive appearance. Strong, erect, sturdy bush plants, which produce short runners. Very prolific. The shelled beans are pinkish buff color, mottled and streaked. Can be used as a dry shell or green shell bean. Of fine flavor.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.25, Postpaid


Not prepaid, 15 lbs. (pk.) \$6.40; 60 lbs. (bu.) \$24.60

BEET

Plant 6 pounds of seed per acre.
In the garden make three plantings at different times.
Plant 1 oz. to 100 feet of row at each planting.

Culture. Successful culture demands good soil, to which has been applied 600 to 1,200 pounds per acre of good standard vegetable fertilizer. Plant seeds in rows 1½ to 2 feet apart and when 3 inches high thin to 3 or 4 inches apart in the row. In Florida, sow from latter part of September to March.

Prices quoted are postpaid.
Write for special prices on larger quantities.



Asgrow Wonder

ASGROW WONDER (Early Blood Red Market). (56 days.) One of the earliest and most uniform beets ever introduced. Roots very attractive in appearance, slightly flattened globe, smooth, with small tap root, of dark red color outside. Flesh of deep dark red, tender, and excellent quality.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.40
5 to 25 lbs. \$1.30 per lb.

CROSBY'S EGYPTIAN (Improved). (60 days.) This standard variety produces beets of uniform flattened globe shape, very much flattened at the top, of deep red color outside, with a deep purple color with little zoning inside. Sweet and tender.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.40
5 to 25 lbs. \$1.30 per lb.

DETROIT DARK RED. (68 days.) A globe-shaped beet, of dark red color, outside as well as inside, without white zoning; sweet and of excellent quality.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.

SWISS CHARD

Plant 4 pounds per acre.

In the garden plant one packet to 25 feet of row.

Culture. Plant like beets in the fall, winter, or spring, in rows 2 feet apart, and thin to 12 or 15 inches in the row.

FORDHOOK GIANT. (68 days.) The thick, fleshy, dark green leaves are much crinkled or savoyed, and the pearly white stems are very broad, thick, and fleshy. A very vigorous, hardy grower.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.

SUGAR AND STOCK BEET

Culture. Plant 4 to 6 pounds per acre in rows 2½ to 3 feet apart, and thin to 6 inches apart in the row.

SUGAR BEET, KLEIN WANZLEBEN. (85 days.) Valuable for sugar making and stock feeding.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

STOCK BEET, MAMMOTH LONG RED. (80 days.) An enormous yielder and one of the best for dairymen.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

BRUSSELS SPROUTS

Sow ¼ pound in seedbed to plant an acre.

In the garden sow 1 packet in seedbed.

Sow seed and cultivate same as cabbage.

LONG ISLAND IMPROVED. (90 days from setting out plants.) Plants grow to a height of two feet. Stems are

thickly set with small firm, cabbage-like heads, 1½ inches in diameter, maturing small heads successively. Cooked and served like cabbage.

Pkt. 10c; ½ oz. 25c; oz. 45c; ¼ lb. \$1.40; 1 lb. \$5.00
5 to 25 lbs. \$4.90 per lb.

BROCCOLI

Sow ¼ to ½ pound in seedbed to plant an acre.

In the garden 1 packet to 40 feet or row, or 1 packet in seedbed for 25 plants to set 40 feet of row.

Culture. In Florida, seed may be sown in seedbeds from September to January. Plants should be set in rows 3 feet apart, and 1½ to 2 feet apart in the row. When the main head is cut, thick stems form at the base of leaves on the stalk below. At the end of each branch a medium-sized green head develops. These small heads should be cut with 4 to 5 inches of stem, and tied in bunches for market.

EARLY GREEN SPROUTING OR CALABRESE. (55 days from plants.) This is the old standard early variety of Broccoli which produces tall, erect, bushy plants, with an attractive dark, bluish-green color to the head. Forms a large, compact central head. After the central head is cut the side shoots send out small heads over a long season. An exceptionally high yielder.

Pkt. 10c; ½ oz. 20c; oz. 30c; ¼ lb. 85c; 1 lb. \$3.00
5 to 25 lbs. \$2.90 per lb.

TEXAS 107. (50 days from plants.) Early maturing variety. In numerous Florida trials in different sections of the State it consistently ranked top in yield. Produces a medium size main head with a great number of good size side shoots. Individual bud formation in head and sprouts, of medium size and very tight and compact, slightly lighter green in color than Early Green Sprouting.

Pkt. 10c; ½ oz. 20c; oz. 30c; ¼ lb. 85c; 1 lb. \$3.00
5 to 25 lbs. \$2.90 per lb.

WALTHAM 29. (60 days.) Low growing, heavy, short compact plant, which produces a large, central head. The plant is heavily branched with dark green, tight buds that are of medium size, evenly distributed. Both the plant and head have a most attractive dark blue-green color. It has ability to come through periods of hot weather without the heads opening up. Because of its uniformity, a large portion of the crop matures at one time. Its excellent dark green color, fine head and bud texture and high yield make it a valuable addition for Florida. Of excellent quality and appearance for fresh market and for freezing.

Pkt. 10c; ½ oz. 25c; 1 oz. 40c; ¼ lb. \$1.15; 1 lb. \$4.00
5 to 25 lbs. \$3.90 per lb.



Broccoli, Waltham 29

CABBAGE

Sow 6 ounces in seedbed to plant an acre or 12 ounces per acre in field.
For the garden, sow 1 packet in seedbed for 50 plants to set 60 feet of row.



A typical plant of Marion Market (Yellows Resistant) Cabbage

Culture. Cabbage is a heavy feeder and should be grown rapidly. For maximum yields on sandy loam soils 1500 to 2000 lbs. per acre of a 4-7-5 fertilizer mixture should be applied in the row a few days prior to setting plants. In addition, two or three side dressings of Nitrate of Potash or Nitrate of Soda should be made at three to four week intervals. Where planted in seedbeds, which is the desirable and common practice, plant seed thin in drills 6 inches apart, cover $\frac{1}{4}$ to $\frac{1}{2}$ inch deep, using 8 ounces of seed for an acre of plants, and do not make the soil in seedbeds as rich as soil where plants are to be set. When seed is planted in the field, use 12 to 16 ounces of seed per acre. Make rows $2\frac{1}{2}$ to 3 feet apart and thin to 12 to 18 inches apart in row, depending on variety and size of head you wish to produce. For a winter crop of cabbage in Florida seed is sown in September and early October. For an early spring crop plant in November or early December. For the control of cabbage insects and diseases see page 53.

To control downy mildew disease in cabbage seedbeds (see page 53). For the prevention of downy mildew and other cabbage diseases in the field. (See page 53).

Our cabbage seed is grown in disease-free territory on the Pacific coast where Black Rot has never been found by plant pathologists who inspect our seed growing fields each year.

The number of days in parentheses after each variety indicates the relative number of days to marketable heads from setting out plants. From the time seed is sown until the plants are ready for setting, thirty-five days are usually required.

Prices quoted are postpaid.
Write for prices on larger quantities.

EARLY ROUND DUTCH. (71 days.) A hardy, and dependable variety of round headed cabbage. Heads are firm and of high quality. Plants vigorous, small, and compact, with erect leaves and short stems. Suitable for close planting, but must be fertilized heavily if planted close to produce size. Withstands cold weather. Especially suited for muck land production.


Pkt. 10c; oz. 40c; $\frac{1}{4}$ lb. \$1.25; 1 lb. \$4.50
5 to 25 lbs. \$4.40 per lb.

MARION MARKET (Yellows Resistant). (75 days.) This is a late maturing disease-resistant strain of Copenhagen Market, highly resistant to Yellows. Larger and coarser in growth than Copenhagen Market but makes somewhat heavier yields, with large, firm, solid heavy heads. Will produce a normal crop and heavy yield on Yellows infested soil where non-resistant varieties will fail.

Pkt. 10c; oz. 35c; $\frac{1}{4}$ lb. \$1.00; 1 lb. \$3.50
5 to 25 lbs. \$3.40 per lb.

MIDSEASON MARKET. (75 days.) The heads are of medium size, hard, globe-shaped on short stems. Leaves are dark green, thick and leathery, making this variety hardy and resistant to cold damage, and also making it a good shipper in bags, the heads not being bruised so readily as other varieties. Of crisp texture and good quality.

Pkt. 10c; oz. 35c; $\frac{1}{4}$ lb. \$1.00; 1 lb. \$3.50
5 to 25 lbs. \$3.40 per lb.



Golden Acre

Pkt. 10c; oz. 35c; $\frac{1}{4}$ lb. \$1.00; 1 lb. \$3.50
5 to 25 lbs. \$3.40 per lb.

GOLDEN ACRE. (64 days.) An extremely early, small to medium-sized round headed cabbage of the Copenhagen Market type, but considerably earlier than most stocks of Copenhagen Market. Our strain of this variety heads up very uniformly and is exceptionally well bred.


COPENHAGEN MARKET (Early Strain No. 86). (68 days.) Our early strain of Copenhagen Market has become very popular in all the large cabbage producing sections of Florida. It is short-stemmed, compact with very solid, round, medium-sized heads, having few outer leaves, thus facilitating close planting, very uniform in shape, and matures practically all at the same time. This uniform maturity is desirable from a marketing standpoint.

Pkt. 10c; oz. 35c; $\frac{1}{4}$ lb. \$1.00; 1 lb. \$3.50
5 to 25 lbs. \$3.40 per lb.



Copenhagen Market. (Early Strain No. 86.)

BADGER MARKET (Yellows Resistant). (72 days.) This new yellows resistant cabbage is showing much




Badger Market

promise on Florida muck soils. On sandy soils it is inclined to be too small unless heavily fertilized. In season it is only a few days later than Early Copenhagen Market No. 86. It is extremely uniform as to type and time of maturity. It makes a uniform, small, compact, round head, of attractive dark blue-green color, very firm and hard, even on loose muck, with a short to medium core. It is very resistant to splitting, and can be left in the field for ten days to

two weeks after it is mature and still show no signs of splitting. It possesses excellent edible quality. This new variety is strongly recommended for Florida muck and the better sandy loam soils where it makes heads two and one-half to three pounds in weight, just the right size for northern markets.

Pkt. 10c; oz. 50c; ¼ lb. \$1.50; 1 lb. \$5.50
5 to 25 lbs. \$5.40 per lb.

EARLY GLORY. (72 days.) A medium-early, round headed type. Somewhat similar to Copenhagen Market, but slightly later, larger, with a darker green color and a heavier yield. The heads are protected by large, thick, overlapping wrapper leaves, thus resisting cold damage. Heads are solid, of medium size, and of excellent shipping and market quality. An improved early maturing strain of the old Glory of Enkhuizen.



Early Glory

Pkt. 10c; oz. 35c; ¼ lb. \$1.00; 1 lb. \$3.50
5 to 25 lbs. \$3.40 per lb.

RED ACRE. (72 days.) This variety fills a long felt need in being the only red cabbage that matures very early, only a few days later than Early Copenhagen Market. The plant is small, compact and makes good-sized round, hard, deep red colored heads. Very uniform as to type. Especially desirable for muck land production in Florida, but also does exceedingly well on fertile sandy loams.

Pkt. 10c; oz. 40c; ¼ lb. \$1.15; 1 lb. \$4.00
5 to 25 lbs. \$3.90 per lb.

ROUND RED DUTCH. (80 days.) A medium early variety, similar to Red Acre. Is a little more vigorous but with larger head at maturity. It is possible to cut Round Red Dutch just as early as Red Acre and get as much or more yield per acre. The heads are round, solid, with a deep red color.

Pkt. 10c; oz. 40c; ¼ lb. \$1.15; 1 lb. \$4.00
5 to 25 lbs. \$3.90 per lb.


JERSEY WAKEFIELD. (62 days.) Very early and resistant to cold weather. The small, medium-sized, conical-shaped, pointed heads are very firm, of finest quality, and especially desirable for home use.

Pkt. 10c; oz. 35c; ¼ lb. \$1.00; 1 lb. \$3.50
5 to 25 lbs. \$3.40 per lb.

CHARLESTON WAKEFIELD. (70 days.) The heads are larger and broader at the base, not quite so pointed and later in maturing than Jersey Wakefield. Of uniform size, very solid and of resirable market quality. Resistant to cold weather and premature seeding.

Pkt. 10c; oz. 35c; ¼ lb. \$1.00; 1 lb. \$3.50
5 to 25 lbs. \$3.40 per lb.

SAVOY-CHIEFTAIN. (90 days.) Savoy cabbage is commonly called "curly leaf" or "curly" cabbage. It has a distinct flavor which is favored by many people. The plant is of large growth. The large, rich, dark green leaves are uniformly wrinkled or savoyed and curled. The heads are medium in size, hard and stand a long time without bursting. The best variety of the savoy type for Florida.



Savoy

Pkt. 10c; oz. 40c; ¼ lb. \$1.15; 1 lb. \$4.00
5 to 25 lbs. \$3.90 per lb.

CHINESE CABBAGE

Also known as Celery Cabbage

Plant 1 pound per acre.


In the garden make two plantings at different times.

1 packet to 40 feet of row at each planting.

Culture. This is a cool-weather plant and should not be grown for a late spring or summer crop, because it bolts to seed in hot weather. Seeds should be planted in 2½ to 3 foot rows where the crop is to mature, thinning out as the plants develop to stand 8 to 12 inches in the row. Chinese Cabbage makes excellent cole slaw.

MICHIHLI. (70 days.) From seed planting to edible stage. Sure header, maturing uniformly into long, slender, very firm, well blanched, crisp, tender, and sweet heads. It is slightly more vigorous and darker green than other strains, the heads are more uniform, and slower to bolt to seed. The heads become 19 to 20 inches tall, 3½ to 4 inches thick, tapered near the tip.


Pkt. 10c; oz. 35c; ¼ lb. \$1.00; 1 lb. \$3.50
5 to 25 lbs. \$3.40 per lb.



Michihli (Celery Cabbage)

CANTALOUPE

Plant 1 to 2 pounds per acre.
In the garden plant 1 packet to 50 feet of row.



No. 45 (Powdery Mildew Resistant)

Culture. General cultural requirements, diseases and insects and their control, similar to those of cucumber. (See page 16.) Plant in hills 4 to 5 feet apart, 5 to 6 seeds to a hill. Thin out to two plants per hill when 4 inches high. Plant in 5 foot rows, thinning to 2 feet in the row. Apply 600 lbs. fertilizer 10 days before planting, side dress with 400 to 600 lbs. at vining. For earlier melons, use **Hotkaps**. (See page 75.) For insect and disease control (See page 54).

Prices quoted are postpaid.

Write for special prices on larger quantities.

NO. 45, POWDERY MILDEW RESISTANT (Hale's Best type). (80 days.) Attracting much favorable attention in Florida because of its high yielding, good shipping and holding quality. Vines moderately large and produce a very heavy crop. Highly resistant to powdery mildew, but not to downy mildew. Medium sized fruits (about 3½ lbs.) are short-oval, indistinct ribbing and well netted. Flesh thick, firm, sweet and of good quality, light orange color. Can be picked almost mature for local markets or distant shipping.

Pkt. 10c; oz. 25c; ¼ lb. 75c; 1 lb. \$2.50;
5 to 25 lbs. \$2.40 per lb.

SMITH'S PERFECT (Florida Golden Dew). (90 days.) A strong, vigorous vine growth, withstanding wind and cold damage. Very resistant to downy mildew disease commonly called rust. Also resists aphids. A very good commercial type. Fruits average 4 lbs. Round in shape, slightly flattened at stem and blossom ends, free of ribbing, a tough rind; makes a good shipper and market melon. For market, it is best to harvest them a little on the green side just as they begin to turn yellow. The flesh is firm, very thick, with a small seed cavity, of attractive deep orange or salmon color, and has a delicious flavor.

Pkt. 10c; oz. 25c; ¼ lb. 75c; 1 lb. \$2.50;
5 to 25 lbs. \$2.40 per lb.

HALE'S BEST, JUMBO STRAIN. (78 days.) A large fruited strain of Hale's Best, grown very extensively in Georgia and other southern states. Fruits are oval in shape, showing some trace of ribbing, with heavy coarse netting. Average about 4 to 5 lbs. Flesh thick, relatively small seed cavity, deep orange color.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00;
5 to 25 lbs. \$1.90 per lb.

HALE'S BEST, No. 36. (75 days.) An improved strain of Hale's Best, developed for uniformity in shape and complete netting. Fruits are round in shape, with very heavy, coarse netting, and average about three pounds. Flesh thick, small seed cavity, deep salmon color, very sweet, fine quality. Excellent shipper.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00;
5 to 25 lbs. \$1.90 per lb.

GEORGIA 47 (Mildew Resistant) (New). (72 days.) This new downy mildew and powdery mildew resistant variety is very early, exceedingly prolific, with firm, thick flesh of golden orange color, very sweet. Fruits small, and have a tendency to produce an unattractive swelling at blossom end. For home garden or local markets this variety is highly recommended.

Pkt. 10c; oz. 35c; ¼ lb. 75c; 1 lb. \$2.50
5 to 25 lbs. \$2.40 per lb.

RIO GOLD. A new shipping type cantaloupe, with extreme resistance to both the downy and powdery mildew and to aphids. Comparable in size to Imperial 45, has a greenish-yellow rind with well raised net. Flesh is deep and a bright orange color. Of excellent flavor and sweetness. A very productive melon and valuable for its resistance.

Pkt. 10c; oz. 25c; ¼ lb. 75c; 1 lb. \$2.50;
5 to 25 lbs. \$2.40

CAULIFLOWER

Sow 6 ounces in seedbed to plant an acre.

In the garden sow one packet in seedbed for thirty plants.

Culture. Starting plants in well prepared seedbeds where conditions can be provided for continuous rapid growth. Plant in seedbeds from Sept. to Dec. and cultivate the same as cabbage, spacing 3 to 4 feet between rows and 18 to 24 inches in row. Cauliflower needs rich soil and plenty of moisture. When head develops to the size of an apple, draw outer leaves over head and tie with soft cord to produce attractive, white, solid heads. Our cauliflower seed is grown in black rot free territory. The number of days after each variety indicates time to produce edible heads from setting out plants.

Prices quoted are postpaid.

Write for special prices on larger quantities.

SUPER SNOWBALL. (55 days.) An early maturing strain. Snowball type, in which curled formation of leaves over head protects it from sun and from freezing. Dwarf, compact plants are strong, vigorous. Heads are large, smooth, solid, pure white and very uniform; Recommended as an early winter crop, to mature in cold weather. Susceptible to "whiptail" and should be grown in alkaline soils with high pH to avoid trouble.

Pkt. 25c; ½ oz. \$1.00; oz. \$1.50; ¼ lb. \$4.00;
1 lb. \$15.00

SNOWDRIFT. (62 days.) One of the surest headers known, especially well adapted to withstand the hot, humid climatic conditions in Florida. Vigorous plants grow larger and taller than Snowball, are more compact and upright in growth with long leaves which protect head or curd in its early growth. Large, snow-white attractive heads are very deep. Heavy and firm. An ideal variety for early spring production in Florida. Very free of leafy and ricey heads. Very resistant to "whiptail" disease.

Pkt. 25c; ½ oz. \$1.00; oz. \$1.50; ¼ lb. \$4.00;
1 lb. \$15.00



Snowdrift

CARROT

Plant 4 pounds of seed per acre. In the garden make three plantings at different times. Plant 2 packets to 100 feet of row at each planting.

Culture. Sow in good, deeply worked, fine soil, enriched with 800 to 1200 pounds per acre of a good vegetable fertilizer, in drills $1\frac{1}{2}$ to 2 feet apart, and cover seed $\frac{1}{4}$ to $\frac{1}{2}$ inch deep. When 2 inches high, thin out to 2 to 3 inches apart. Carrots can be planted in Florida from August to March.

Prices quoted are postpaid.


Write for special prices on larger quantities.

DANVERS HALF LONG. (75 days.) Our stock of this variety is half stump or semi-pointed, from 6 to 7 inches in length. The shoulder at the crown is from $1\frac{1}{2}$ to $1\frac{3}{4}$ inches thick, and tapers uniformly to a thickness of about $\frac{3}{4}$ -inch just where the root begins to taper to the tip. The tops are a little larger than those of Chantenay, and the crown is flat without a depression. The roots are smooth and attractive in appearance.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

LONG IMPERATOR. (77 days.) The roots average about 10 inches in length, with a shoulder diameter of $1\frac{1}{4}$ to $1\frac{3}{4}$ inches, smooth, uniformly tapered to a semi-blunt end, deep rich orange color, both exterior and interior with comparatively inconspicuous core. The flesh is of rich orange color, extending to the center of the root, with an indistinct core; fine grained, tender and of excellent quality. The tops are of medium size, but strong enough for good bunching. For shipping quality, attractiveness and popularity in the market. Long Imperator is superior to other long-rooted varieties.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 60c; 1 lb. \$2.00
5 to 25 lbs. \$1.90 per lb.



Long Imperator Carrot

Developed to meet the need for a better market type.



Comparison of root characteristics of carrot varieties. From left to right: Imperator, Nantes (Coreless), Danvers Half Long, Red Core Chantenay.

RED CORE CHANTENAY. (72 days.) The roots are tapered, less stump rooted, and slightly longer than the old Chantenay, averaging about 6 inches. The interior is solid orange-colored without the usual lemon color in the core. The flesh is very tender and sweet, which with the excellent outside color and shape combines to make this variety valuable for the market.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

NANTES. (70 days.) Cylindrical, roots about $5\frac{1}{2}$ to 6 $\frac{1}{2}$ inches long and 1 to $1\frac{1}{2}$ inches thick, very stump rooted with a small rat tail. The color is clear orange-red and the core is rather undefined, being of the same color as the flesh so that it is called "Coreless." The tops are small and short, but heavier than ordinary stocks of Nantes, and the root is very fine-grained, tender, delicate, sweet and of highest edible quality. An excellent high quality variety for the home garden and local market.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

LONG ORANGE. (86 days.) Our stock is the true thick-shouldered strain of Long Orange, $1\frac{3}{4}$ to 2 inches thick at the shoulders. 10 to 12 inches long, and tapering in a straight line to the tip. The tops are rather large. Grown in sections of Florida where a smooth, reddish-colored, long-rooted carrot is desired.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

YELLOW BELGIAN (Stock Carrot). (95 days.) The giant among carrots. Grown exclusively for stock, very large, 12 inches long and $2\frac{1}{2}$ inches thick at shoulder. Pale yellow skin and flesh. Hardy, heavy yielder and very nutritious for stock.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

CELERY

Sow ½ pound in seedbed to plant an acre.

For the garden sow 1 packet in a well protected seedbed for 150 plants to set 50 feet of row. Prices quoted on celery seed are postpaid. Write for special prices on larger quantities.

Culture of Celery. First sowings in Florida are made the latter part of June and continue to December. The main sowings for a winter crop are usually made in July or early August, and for the spring crop, grown principally on muck, they are usually made in October and early November. Celery is always started in protected seedbeds. The soil must be put in perfectly fine condition and made into beds 4 to 6 or more inches high, to prevent overflow in case of heavy rains. Sow the seed broadcast or in rows 6 inches apart, using 1 pound of seed to 125-150 yards of bed 3 feet wide. The seed should be barely covered with soil.

We recommend use of burlap or white muslin as a cover over the seedbed stretching it on a tent-shaped or slanting frame built with the sides about 12 inches from the ground to provide ample circulation of air. This cover will serve as shade and break the force of heavy rains. After the seed is sown, sprinkle with a very fine spray in order to keep the soil cool and moist, which will increase the germination of the seed during the hot summer months. With surface irrigation, water should be circulated continuously around the beds.

When plants in seedbed are 6 inches high, transplant to the field in double rows 3 to 3½ feet apart, with 7 inches between the double rows, and set the plants 4 to 5 inches apart in the row. The plants may be set in single rows 3 feet apart, allowing 4 to 5 inches between plants in the row. The Pascal varieties should always be planted at least 6 inches in the row. To grow fine celery, a large supply of moisture and fertilizer is essential from the time the seed is sown in the seedbeds until the crop is ready for harvest.

For the control of insects and diseases of celery see page 53. For the control of pink rot of celery use Granular Aero Cyanamid.

Good control of black heart in celery has been obtained by spraying weekly with 10 lbs. Calcium Nitrate dissolved in 100 gal. water per acre.



Summer Pascal

The number of days in parentheses after each variety indicates the relative number of days to marketable stage or harvest from setting out plants. It usually requires about seven or eight weeks to grow plants large enough to be set in the field.



Typical plant of Florida Green Pascal Celery ready for packing in Howard crates.

FLORIDA GREEN PASCAL. (125 days.) A very vigorous, compact-growing variety, with broad, semi-round, medium-long, thick, very meaty, smooth, attractive ribs. The ribs or petioles will average from 7½ to 8 inches to the first joint and plants will range from 20 to 23 inches overall. The ribs or stalks are of ideal edible quality, being tender, brittle, stringless and crisp. Even the outermost stalks possess a rich and nutty flavor. Our Florida Pascal strain has been developed for smooth glossy ribs, a taller, more compact growth, and better heart formation than other strains of Green Pascal.

This stock is not subject to premature seeding, and can be grown at any time during the fall, winter or spring in Florida. It is important not to cut this variety too soon, as it requires 2 to 3 weeks more time in the field than varieties of the Golden type, but will out-yield the Golden type. The yield and quality are materially cut down if harvested too soon. Set plants of this variety at least 6 to 7 inches apart in the row for large sizes and heavy yields. The seeds of Green Pascal are larger than Golden type celery, and therefore 25 to 50% more seed by weight is required to get the same stand.

Pkt. 15c; ½ oz. 40c; 1 oz. 75c
¼ lb. \$2.50; 1 lb. \$9.00

SUMMER PASCAL (Waltham Strain). (115 days.) This is an early maturing Green Pascal variety. A rapidly growing type of celery with long, upright compact ribs or petioles, making a very attractive appearing celery for the markets. The ribs are thick and semi-rounded, but not quite as smooth as our strain of Florida Green Pascal, although the Waltham strain will mature about a week earlier, and makes a high yield of large sizes. This strain must be cut promptly when ready or it will tend to go down, in the field after maturity.

Pkt. 15c; ½ oz. 40c; 1 oz. 75c
¼ lb. \$2.50; 1 lb. \$9.00

CELERY

FLORIDA GOLDEN NO. 15. (115 days.) One of the best varieties for a mid-winter crop in Florida, because it withstands cold weather better than most varieties of the Special or Golden Plume type. Selected for uniformly long, thick, round ribs, large full hearts of rich golden yellow color, and for heavy yields of good sizes. The ribs average eight to ten inches in the first joint, rounded and exceptionally thick. The plants range from 21 to 24 inches over all, making an attractive pack. It is intermediate between the Old Golden and Special or Golden Plume types, possessing the desirable characteristics of both types. Recommended especially for a Florida mid-winter crop.

Pkt. 15c; ½ oz. 35c; 1 oz. 65c; ¼ lb. \$2.25
1 lb. \$8.75



Part of typical row of Florida Golden No. 15 Celery

COLLARD

Sow ½ pound in seedbed to plant an acre, or plant 2 pounds per acre in field.
For the garden make two sowings at different times. Sow one packet in seedbed for 35 plants for each planting to set 50 feet of row.

Culture. Collard is an old standby for winter greens all over the South. May be started in seedbeds or direct in the field, in rows 2½ to 3 feet apart, with plants 14 to 18 inches apart in the row. Use about 1200 pounds of fertilizer per acre before setting plants. Collard can be grown every month of the year in Florida, but it grows best and produces the highest edible quality during cool weather. This vegetable should be included in every Florida family garden, as it is very hardy, easy to grow and will furnish an abundance of greens, high in health-producing vitamins.

The number of days in parentheses after each variety indicates the time from setting out plants to harvest. It requires about a month in the seedbed to get plants from seed sowing to transplanting stage.

Prices quoted are postpaid.
Write for special prices on larger quantities.

GEORGIA 912. (50 days.) A non-heading type of the cabbage family, very extensively grown for greens in home and commercial gardens in Florida.




Georgia 912

GEORGIA 912 is a distinct strain of Georgia type collard, developed at the Associated Seed Growers' South-eastern Breeding Station at Bradenton, Fla. through in-breeding and progeny testing. Compared to the standard Georgia type it is slower to bolt or to seed; has more compact plant habit, with shorter petioles and broader leaf blades, and has more savoyed leaves. Very uniform plant type and a heavy yielder.

Because of the more compact plant habit, broader and more savoyed leaves, good uniformity of stock, cold hardiness and long standing habit, Georgia 912 should find a place for commercial shipping and processing, as well as for general use in collard growing areas.

Pkt. 10c; oz. 15c; ¼ lb. 40c; 1 lb. \$1.20
5 to 25 lbs. \$1.10 per lb.



Vates

VATES. (55 days.) A very large, vigorous, rather low growing plant, not subject to wind damage, with very large, broad, heavy, thick leaf blades, the leaf blade developing from the base of the short stem or petiole. The leaves are smooth with an attractive deep, dark green color, the stems and leaves never turning yellow or purplish even in the coldest weather or on poor soils. It retains this deep green color in shipment and on the markets. A heavy yielder, slow to bolt to seed in cold or hot weather. Uniform as to plant type.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.30
5 to 25 lbs. \$1.20 per lb.

LOUISIANA SWEET (Dark Green Strain). (50 days.) The plant develops a deep, compact, rosette center with a large mass of thick, tender, dark green leaves entirely free of purple or red color. This new variety is highly recommended for commercial growers and shippers. The tall growing plant is hardy and slow to go to seed.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c; per lb.



A field and typical fruits of Marketer (Early Green Market) Cucumbers

CUCUMBER

In the garden plant 1 packet to 50 feet of row.
Plant 2 pounds per acre.

Culture. Rich, moist, sandy loam soil is the best for growing cucumbers. The soil should be well fertilized with a good vegetable fertilizer applied at the rate of 1200 lbs. per acre a week or ten days before planting; 600 to 1000 lbs. is generally used as a side-dresser just before the plants start to run. Fall plantings are made in late August and in September. Spring planting in Florida begins early in January and continues through February. Plant in rows 5 to 6 feet apart, 6 inches apart in the row, and thin to 1 or 2 feet between every two plants in the row. Sometimes it is difficult to get a stand early if the ground is wet and cold, as the seed will rot before it germinates. Therefore seed should be treated. (See page 52).

If young plants are very badly damaged with frost or cold, it is best to plant over. When young plants are stunted they never make a normal crop and usually not much earlier than if they were planted over. A row of oats planted between every two rows of cucumbers makes a good windbreak.

It is very necessary to have hives of bees in or near cucumber fields to ensure pollination and a good set of fruits. The bees should not be placed in the field until the cucumbers begin to blossom, in order to provide food for the bees so they will remain in the cucumber fields. Without proper pollination the cucumbers are deformed, and a large percentage of bottlenecks and nubbins are produced.

For the control of insects and diseases of cucumbers see page 54. To control angular leaf spot treat seed with a mercuric chloride. (See page 52.) The best control for downy mildew, sometimes called "rust," is Nabam spray or Zineb Dust. (See page 54.) The most effective and safest control for aphids and some other cucumber insects is Isotox spray or Kilgram dust. (See page 54).

For an earlier cucumber crop and a protection against beetles, use **Hotkaps**. (See page 75).

For dill to make cucumber pickles see Page 20.

Prices quoted are postpaid.

Write for special prices on larger quantities.

The number of days after each variety represents the time from seed planting to first picking under average spring or fall conditions in Florida.

MARKETER (Early Green Market). (60 days.) The leading variety for yield and quality. Holds its dark green color much better than most varieties, when grown in late spring or early fall. It averages about 8 inches in length, and is especially desired by the eastern and southern markets where a medium size cucumber is preferred. The fruits are exceptionally uniform in size, shape and color, with practically no jumbos or oversize even when not harvested for several days. It seems to stand heat of late spring better than other varieties. Exceedingly prolific, throwing an exceptionally large number of female blossoms per plant. It will average over 90 per cent fancies. Buyers pay a premium for this cucumber, when other varieties go begging at lower prices.

Pkt. 10c; oz. 25c; ¼ lb. 65c; 1 lb. \$2.25
5 to 25 lbs. \$2.15 per lb.



Ashley (Mildew Resistant)

ASHLEY. (55 days.) (Mildew resistant.) A new downy mildew resistant cucumber developed at Clemson Truck Experiment Station, South Carolina. Vigorous vines and a heavy producer. Fruit dark green color, about 8" long, tapering at the stem end. This new cucumber looks promising and should be given a trial by growers.

Pkt. 10c; oz. 30c; $\frac{1}{4}$ lb. 85c; 1 lb. \$2.75
5 to 25 lbs. \$2.65 per lb.

SANTEE (Downy Resistant). (50 days.) Similar to Palmetto, but earlier and more prolific, and darker green color. Attractive cylindrical shape with blunt ends. Similar to but darker green than Cubit. Quite resistant to downy mildew.

Pkt. 10c; oz. 30c; $\frac{1}{4}$ lb. 85c; 1 lb. \$2.75
5 to 25 lbs. \$2.65 per lb.

PUERTO RICO NO. 39 (Mildew Resistant) (Special Long Strain). (50 days.) This variety is highly resistant to downy mildew or blight. Will hold up under adverse weather conditions, when other varieties go down with mildew disease, making it especially valuable as a home garden variety and market cucumber. Although highly resistant to downy mildew it is not immune, and should be sprayed or dusted regularly with a good fungicide like Dithane. The vines are rather compact in growth, hardy and very productive. As originally introduced the fruits of this variety were rather short, but we have selected for long fruits until with our special strain the fruits will average about 7½ inches in length with dark green color.

Pkt. 10c; oz. 30c; $\frac{1}{4}$ lb. 85c; 1 lb. \$2.75
5 to 25 lbs. \$2.65 per lb.



Puerto Rico No. 39 (Mildew Resistant)

PALOMAR. (55 days.) A downy mildew resistant cucumber developed by crossing Palmetto and Marketer. Vines are heavy and vigorous. Extra early and a heavy yielder. Fruit about 8½" long, dark green color. Some fruits are slightly curved.

Write for prices.



Stays Green (Pickling Type)

STAYS GREEN (Pickling Type). (40 days.) This variety should be grown only for pickles, not for a slicing or as a fresh market cucumber. It is very early and exceedingly productive of short, dark green, warty cucumbers most desirable for dill pickles.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

PALMETTO (Mildew Resistant). (60 days.) Developed by Dr. W. C. Barnes of the Clemson College Truck Experiment Station, Charleston, S. C., from a cross of Puerto Rico 40 and Cubit in order to develop a good market type possessing the downy mildew resistance of the Puerto Rico 40. This variety produces vigorous vine growth, and is highly resistant to downy mildew, the most serious cucumber disease of the South. Palmetto will survive and produce a good crop, when other varieties may go down with mildew under cool, wet weather conditions.

Pkt. 10c; oz. 30c; $\frac{1}{4}$ lb. 85c; 1 lb. \$2.75
5 to 25 lbs. \$2.65 per lb.



Palmetto (Mildew Resistant) Cucumbers

SWEET CORN

(For Field and Roasting Ear Corn, See Pages 38 and 39)

10 pounds seed required per acre. In the garden make two plantings. Plant $\frac{1}{4}$ lb. to 200 feet of row at each planting.

Our Hybrid Sweet Corn seed is now graded and sized for better, more accurate stands and more uniform maturity.

Culture. Rows should be $2\frac{1}{2}$ to 3 feet apart with the plants 10 to 18 inches depending on type and season. Sweet corn requires much more fertilizer than field corn. Use about 1000 lbs. per acre of a good commercial fertilizer mixture high in potash. It is claimed that potash helps the tips of the ears to fill better and improves the edible quality of sweet corn. When corn is about knee high apply 200 lbs. per acre of sodium nitrate or its equivalent. This fertilizer practice will greatly increase the yield. Early plantings are made in Florida from January 1st until March. Ship in special corn crates, five to six dozen ears per crate. For control of corn ear worms and leaf blight see page 54.



Calumet Sweet Corn. Note kernel covering over tip of ears.

CALUMET. (76 days.) This hybrid variety has received the high ratings for yield, uniformity and attractive appearance, and appears to have some resistance to ear worms. The plant is $6\frac{1}{2}$ to 7 feet, slightly taller than Ioana, is relatively resistant to drouth, and almost suckerless, with large dark green leaves. The ears are long, averaging about 8 inches, cylindrical, well filled to the tips and with tight husks of good length. Kernels are light yellow, medium wide, very deep, usually in 12 to 14 rows. Very uniform in maturing and a high yielder. Makes a very attractive pack of from 5 to 6 dozen per crate usually 5 to $5\frac{1}{2}$ dozen. One of the finest of the many hybrids recently introduced.


$\frac{1}{4}$ lb. 20c; 1 lb. 80c; 2 lbs. \$1.35, postpaid

Not prepaid, 10 lbs. \$4.70; 25 lbs. \$11.50; 100 lbs. \$45.00

GOLDEN CROSS BANTAM. (73 days.) Golden Cross Bantam is a hybrid sweet corn that has done well as a high quality, yellow-kerneled, early variety in Florida. The small, short stalk ($5\frac{1}{2}$ to 6 feet) is very sturdy and vigorous, Suckers quite badly. Ears 6 to 7 inches long, 10 to 14 rows. Kernels are attractive, yellow in color. Uniform, higher yielding. Recommended for home use, local markets and for shipping to select markets.

$\frac{1}{4}$ lb. 20c; 1 lb. 80c; 2 lbs. \$1.35, postpaid

Not prepaid, 10 lbs. \$4.10; 25 lbs. \$10.00; 100 lbs. \$39.00



Sixtypak

SIXTYPAK (XP-112) (74 days.) This new hybrid developed by Associated Seed Growers Inc., has been thoroughly tested (under breeding number XP-112) with other leading varieties of hybrid sweet corns in all important corn growing sections of Florida. In every test the "Sixtypak" produced a higher yield than any other variety.

The plants are very uniform, about six and one-half feet tall with heavy dark green foliage, a high percentage of them producing two fancy ears per stalk. The ears are very attractive, $7\frac{1}{2}$ to 8 inches long, well filled to the tip, with an average of 14 rows of golden yellow kernels. The husk is dark green, making a very attractive five dozen pack.

Sixtypak is one of the most promising hybrid sweet corns introduced for Florida. Seed supply limited.

$\frac{1}{4}$ lb. 25c; 1 lb. 80c; 2 lbs. \$1.45; 10 lbs. \$4.90;
25 lbs. \$12.00; 100 lbs. \$47.00

IOANA. (74 days.) A very productive, yellow-kerneled, hybrid. Vigorous plants are highly resistant to drouth. Medium tall, 6 feet, sturdy, with large, broad and numerous dark green leaves. Ears $7\frac{1}{2}$ to 8 inches long, 12 to 14 rowed, well-filled with medium wide, attractive light yellow kernels of fine quality and flavor, making it highly resistant to worms. Husks retain green color for a long time, making for fresh, attractive market appearance.

$\frac{1}{4}$ lb. 20c; 1 lb. 80c; 2 lbs. \$1.35, postpaid

Not prepaid, 10 lbs. \$4.10; 25 lbs. \$10.00; 100 lbs. \$39.00



Ioana

EGGPLANT

Sow $\frac{1}{4}$ pound in seedbed to plant an acre.

For the garden sow one packet in seedbed for 17 plants to set 50 feet of row.

Culture. For a spring crop, plant seed in November, December, and January; for fall crop, plant in June, July, and August, in muslin or burlap-covered seedbeds. Set in rows 4 to 5 feet apart and $2\frac{1}{2}$ to 3 feet between the plants. Eggplant being a long season crop requires several fertilizer applications. Best results are obtained by applying half or more of a complete fertilizer at planting time and the remainder when the crop is one-third to one-half grown. The initial application should be made in two bands each located 2 or 3 inches below and 3 inches to the side of the plant row. Subsequent application should be drilled close to the plant row.

Top-dressing applications of nitrogen or a combination of nitrogen and potash throughout the season is necessary to hold up size and color of fruit. For the control of insects and diseases of eggplant see page 54.

The number of days in parentheses after each variety indicates the time to first marketable fruits, from setting out plants in the field or garden. It usually requires six to eight weeks to produce plants for field setting.

Prices quoted are postpaid.

Write for special prices on larger quantities.

FORT MYERS MARKET. (83 days.) We introduced this variety, which is today one of the leading varieties grown in the South. Our breeding department has made special selections to improve the variety for uniformity in type of plant, size, shape and color of fruit. Produces immense yields of uniform, large-sized fancy fruits over a long period. The plants are of the high-bush type, with very tall, vigorous growth. The fruits are slightly elongated, cylindrical in shape, and they possess the intense deep black color so much in demand among buyers and in the markets.

Pkt. 10c; oz. 65c; $\frac{1}{4}$ lb. \$2.15; 1 lb. \$7.50
5 to 25 lbs. \$7.40



Florida Market (Cook's Strain)

FLORIDA MARKET (Cook's Strain) (phomopsis resistant). (75 days.) Resistant to tip over, leaf blight and fruit rot, all diseases caused by phomopsis fungus which have become so severe in the state as to make eggplant production unprofitable in some sections. It makes a very vigorous growing, sturdy bush, holding the fruits well off the ground. It is extremely early and a heavy yielder, bearing over a long season. This makes it a very desirable variety for south Florida where it can be planted in early fall for production throughout the winter and spring months. The attractive fruits are cylindrical in shape and develop to a large size. They are of an attractive, dark color. After several years of selection on the original Florida Market variety, we have in this **Cook's strain** a great improvement in color, shape and uniformity.

Pkt. 10c; oz. 75c; $\frac{1}{4}$ lb. \$2.25; 1 lb. \$8.00
5 to 25 lbs. \$7.90 per lb.



A field of Fort Myers Market Eggplant at harvest time.

ENDIVE



Improved Florida Deep Heart Escarolle

Sow $\frac{1}{2}$ to 1 pound in seedbed to plant an acre, or 1 to $1\frac{1}{2}$ pounds per acre in the field.

For the garden sow one packet in seedbed for 50 plants to set 40 feet of row, or plant one packet to 40 feet of row.

Culture. Plant from August to February in seedbeds or in the fields or garden the same as lettuce. Set in rows $1\frac{1}{2}$ to 2 feet apart and 9 inches apart in the row. Soil and fertilizer requirements are the same as for lettuce. (See page 21).

The number of days indicated after each variety is the number of days from seed planting in the field or garden to edible or harvest stage. For insect control see page 54.

Prices quoted are postpaid.

Write for special prices on larger quantities.

ESCAROLLE—IMPROVED FLORIDA DEEP HEART. (90 days.) One of the best and most prolific types to grow commercially. In great demand on northern markets, and well adapted to Florida soil and climatic conditions. The leaves are large, broad, thick, slightly crumpled, of medium green color, with thick, nearly white midribs. Our improved stock of this variety is of the Full-Heart type, developing an upright growth, producing a very heavy, thick matted plant with large, deep, well balanced creamy white hearts.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

RUFFEC GREEN CURLED. (Green Ribbed). (95 days.) Sometimes called "Chicory." A hardy vigorous sort, with deep green, curled, finely cut leaves, giving a mossy appearance. The midrib is thick, broad, green, tender and fleshy. A dense mass of deeply divided leaves forms the head which blanches readily to a beautiful creamy white.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

DILL

A packet will sow 10 ft. of row; an ounce 50 ft. of row.

DILL—LONG ISLAND MAMMOTH. (65 days.) Partially matured seed clusters and flowerheads are useful for flavoring dill pickles. A tall growing plant, 2 to $2\frac{1}{2}$ feet high, vigorous, with very finely divided leaves. Seed clusters are borne in umbels or clusters, have a strong aromatic odor, and mildly pungent flavor.

Pkt. 10c; oz. 35c; $\frac{1}{4}$ lb. \$1.00; 1 lb. \$3.50

GARLIC (90 days)

Plant 1 pound to 30 feet of row.

Garlic sets come in the form of bulbs which are divided into so-called sets or cloves, which are planted the same as onion sets, from August to November, in rows 15 to 20 inches apart, setting the cloves 3 to 4 inches apart in the row, being sure not to cover the tips of the cloves. Garlic is a southern crop and does well in Florida.

$\frac{1}{2}$ lb. 45c; 1 lb. 85c; 5 to 25 lbs. 75c per lb.; postpaid

KALE

Plant 1 ounce to 100 feet of row, or 1 packet to 25 feet of row. **Culture.** Kale is grown for greens. Plant in Florida from August to April, in rows $2\frac{1}{2}$ feet apart, and for large plants thin from 20 to 30 inches apart in the row. The plants are extremely hardy, withstanding cold and heat.

DWARF BLUE SCOTCH. (70 days from seed sowing to edible stage.) This is the most desirable hardy variety for Florida and for the market. It forms a handsome, dwarf, spreading plant with many long, heavy, finely curled, attractive, blue-green, plume-like leaves. Our stock is exceptionally well-bred, uniform and very attractive. This vegetable ranks high in vitamin content.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

KOHL RABI

For the garden make short row plantings at different times. Plant 1 packet to 25 feet of row.

Culture. Grow in row $1\frac{1}{2}$ to 2 feet apart, thinning from 6 to 8 inches apart in the rows. Planting every ten days will give a succession of tender, fleshy bulbs until hot weather. Start fall planting September, and successive plantings can be made until early spring.

WHITE VIENNA. (50 days from seed sowing to edible stage.) The fleshy stems, or so-called bulbs, are formed above ground and are very light green, tender and delicate, and of best quality if used when about 2 to 3 inches in diameter, before fully grown. Prepare by boiling the same as turnips. They make a delicious and nutritious dish.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 60c; 1 lb. \$2.00
5 to 25 lbs. \$1.90 per lb.

LEEK

Sow $1\frac{1}{2}$ lbs. in seedbed to plant an acre, or plant 3 lbs. per acre in the field.

For the garden sow 1 pkt. in seedbed for 120 plants to set 50 feet of row, or plant 1 pkt. to 50 feet of row.

Culture. Leek seed may be planted any time during the fall or winter directly in the field, or preferably in seedbeds, in good rich soil. When plants in seedbeds are six to eight inches high, transplant to rows 18 inches apart, and 5 to 6 inches apart in the row, or thin out to this spacing in the field. The long, thick stems when boiled and served with a butter or cream dressing are delicious. The stems are eight to ten inches long and $1\frac{1}{2}$ inches thick. When marketed they are tied in bunches with twelve Leeks to a bunch.

LARGE AMERICAN FLAG. (140 days from seed planting in field or garden to full edible stage.) The heaviest yielding and the best variety for Florida growing conditions. It makes larger and thicker, more fleshy stems than other varieties.

Pkt. 10c; $\frac{1}{2}$ oz. 25c; oz. 45c; $\frac{1}{4}$ lb. \$1.40; 1 lb. \$5.00
5 to 25 lbs. \$4.90 per lb.

LETTUCE

Sow ½ to 1 pound in seedbed to plant an acre, or plant 2 to 3 pounds per acre in field. For the garden make two plantings at different times. Sow 1 packet in seedbed for 60 to 80 plants to set 50 to 75 ft. of row at each planting.

Culture. First sowings are made in Florida the beginning of September and continued until March. Low, damp soil is best suited for this crop. Lettuce must be grown rapidly in order to give it size, tenderness, crispness and high edible quality. Therefore, the land must be highly fertilized and moist. In Florida about one ton per acre of 4-7-5 fertilizer is commonly used for the Boston, Romaine and leaf types, where 800 to 1200 pounds per acre of a 4-8-8 fertilizer is recommended for the Iceberg type. Broadcast the seed on good moist seedbeds that have been thoroughly prepared, or sow in rows six inches apart and use a light roller to press seed into soil. A pound of seed sown broadcast plants a bed three feet by a hundred yards. Place burlap or well-washed old fertilizer sacks over the beds until seeds sprout. Seedbeds should be kept thoroughly moist at all times after the seed is planted. In hot weather germination may be hastened and improved by placing lettuce seed in moist cloth in the ice box for several days before sowing. In level culture, lettuce plants are usually set 15 by 15 inches in the field, or in rows 1½ to 2 feet apart, and 10 to 15 inches in the row, or in double rows 16 inches apart, with 3½ to 4 feet from center to center of double rows, and 14 inches in the rows. Plants should be watered as set in field. For insect control see page 54.

Rusting of head lettuce and Romaine is usually due to boron deficiency. To prevent or stop this trouble, it is recommended to spray with 1 lb. borax to 50 gallons of water applied at the rate of not more than a total of 4 pounds borax per acre in two applications. Too much borax is toxic.

The number of days in parentheses after each variety represents the time required to develop marketable heads from the planting of seed.

Prices quoted are postpaid.

Write for special prices on large quantities.

EIFFEL TOWER COS. (Kilgore's Mammoth Romaine) (73 days.) Tall, vigorous and makes large heads with correspondingly heavy yields. It can be left in the fields for 10 days to two weeks after it is ready to harvest without danger of bolting to seed. It forms firm, solid heads with the leaves well capped over the top, and is very uniform.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.10
5 to 25 lbs. \$2.00 per lb.

PARRIS ISLAND COS (Mosaic Resistant). (68 days.) This is a new Cos or Romaine type resistant to lettuce mosaic, and very resistant to premature seeding. Similar to Dark Green Cos, but a shade lighter green, with the leaves of somewhat heavier texture. Of value wherever mosaic is a limiting factor in the growing of Cos type lettuce. Uniform and attractive.

Pkt. 10c; oz. 25c; ¼ lb. 85c; 1 lb. \$2.75
5 to 25 lbs. \$2.65 per lb.

GREAT LAKES (No. 118). (83 days.) This strain 118 has been developed for uniformity of size and type, and is the very best strain for Florida. The plants are large, vigorous and have a distinctive, deep grassy-green, attractive appearance. The heads are relatively large, compact, firm and solid, with abundant wrapper leaves which cover the heads, protecting them from cold and heat injury. Great Lakes will stand high temperatures, is much less subject to bottom rot, tip burn and slime rot, and does not bolt so readily as other varieties of this type. It offers a possibility of prolonging the Florida lettuce growing season in late spring, and in early fall.

Pkt. 10c; oz. .35c; ¼ lb. \$1.10; 1 lb. \$3.75
5 to 25 lbs. \$3.65 per lb.

SIMPSON, BLACK-SEEDED. (46 days.) The best large, very early, nonheading or leaf variety. The leaves are very broad, much blistered and crumpled, and excessively frilled on the borders, light yellowish-green in color, very tender, of fine delicate texture, and of excellent quality.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00
5 to 25 lbs. \$1.90 per lb.

WHITE BOSTON (Ocoee Strain). (75 days.) This strain has been selected under Florida conditions for the development of a large framed lettuce similar to Big Boston. Produces large heads and heavy yields.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00
5 to 25 lbs. \$1.90 per lb.

BIG BOSTON. (75 days.) The heads are of cabbage or butterhead type, large sized. The outer leaves are broad, almost smooth, but wavy at the edge, medium light green with a tinge of brown on outer leaf margins.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00
5 to 25 lbs. \$1.90 per lb.

MIGNONETTE. (67 days.) An early maturing, high quality, small crisp head lettuce, large tinged with white hearts, not grown for commercial shipment, but useful for the Florida home gardener and for local markets. High edible quality, being sweet-flavored, crisp and tender.

Pkt. 10c; oz. 35c; ¼ lb. \$1.00; 1 lb. \$3.50
5 to 25 lbs. \$3.40 per lb.

BIBB. (60 days.) A very small, early maturing, loose butterhead type of lettuce, with thick, tender leaves of deep glossy green color, rich in vitamins, rich, creamy yellow interior head possessing superior crispness and flavor. Recommended for home gardens. Set plants only 6 inches apart because of their small size.

Pkt. 10c; oz. 35c; ¼ lb. 65c; 1 lb. \$2.20
5 to 25 lbs. \$2.10 per lb.



Great Lakes Lettuce (No. 118)

M U S T A R D

Plant 4 pounds of seed per acre.
In the garden make two plantings at different times.
1 packet to 40 feet of row at each planting.



Chinese Broad Leaf Mustard



Florida Broad Leaf Mustard

Culture. General cultural directions similar to those for cabbage (see page 10), but is always planted direct in the field or garden. Mustard is one of the quickest growing greens. Sow seed August to March in good rich garden soil in rows 24 inches apart and thin from 8 to 10 inches in the row. Heavy fertilization with a fertilizer mixture high in nitrogen or ammonia, and moist soil are conducive for quick growth, high yield and good edible quality. Does best when grown in cool weather of late fall, winter and early spring. Makes a nutritious dish, high in food value, when boiled for greens.

The number of days after each variety indicates the time from seed to harvest.

Prices quoted are postpaid.

Write for special prices on larger quantities.

FLORIDA BROAD LEAF. (40 days.) Tall and upright, large, thick, broad, smooth, round or oval, medium green leaves (8 to 10 inches long and 8 inches wide.) The leaves have a large, distinct, flattened, whitish stem and midrib. It produces more leaf growth than other varieties. Plants stand a long time without bolting to seed, and do well in hot weather. Of excellent edible quality, the flavor being very mild. Fine for greens and salads.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

CHINESE BROAD LEAF. (40 days.) A hardy and vigorous grower, producing large, broad, elongated or oblong leaves (9 to 10 inches long and 6 inches wide) which are slightly crumpled and of medium dark green color. The leaves have a small, rounded midrib and differ in this respect as well as in shape of leaf from the Florida Broad Leaf. Tender and of excellent quality, with mildly pungent flavor, when boiled for greens.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

FORDHOOK FANCY. (42 days.) A virgorous, upright grower, producing large, bright greens, very deeply fringed leaves, attractive in appearance. A very heavy yielder. The plants stand relatively long without bolting to seed. Of good edible quality, crisp and tender in texture, and mild in flavor. Our stock of this variety is uniformly true to type.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

TENDERGREEN (MUSTARD SPINACH). (45 days.) Although not a true mustard, it is like a quick growing mild mustard, with a spinach flavor, and remains tender even in dry, hot weather. An all-season plant, it withstands heat and may be grown throughout the spring and summer. The smooth, oblong leaves are dark green in color, with small white midribs. Relatively slow to bolt to seed.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.



Fordhook Fancy Mustard

OKRA

Plant 8 pounds per acre.

In the garden plant 1 ounce to 75 feet of row.

Okra does wonderfully well all over Florida where the young seed pods are used in different ways and are highly, esteemed. In addition to being served alone, pods are used in soups and stews to which they give body. It is a healthful vegetable, and should be planted in every Florida garden. We have done a great deal of work on our seed stocks of this valuable Florida vegetable.

Each year, we make special selections for stock seed from only the best, earliest maturing plants. Our seed growing fields are carefully rogued so as to eliminate any slightly off-type plants. We are confident we have the best strains of okra seed obtainable anywhere.

Culture. Okra can be grown in Florida any time from February until September, but it is a warm-weather plant and will not do much until the ground becomes warm. For early planting when the ground is cold always treat okra with Spergon or Arasan. (See pages 61 & 62.) Plant in rows 3 to 4 feet apart and sow thinly in the drill. Okra should be cut every second day. If this is not done some of the pods become too old for market and the young plants will stop bearing pods. Plants should bear for several months after harvesting starts. Okra does best on sandy loam soil with a fair amount of fertility. On sandy loam soils, 500 to 700 lbs. per acre of fertilizer at time of seeding, and two or three side dressings of nitrate of potash during the bearing period are desirable. Okra is very susceptible to nematodes causing root knot disease, and therefore it is best to plant on land not previously used for growing crops that are subject to attack by nematodes. For the control of root knot nematodes use "D-D". (see page 64). For insect control see page 55.

Prices quoted are postpaid.

Write for special prices on larger quantities.

CLEMSON SPINELESS. (55 days.) Plants are semi-dwarf, growing to a height of 4½ feet on fairly good soil and produce uniform, straight, large, thick, ridged pods of the Perkins' Long Green type, of good length, averaging 3 to 3½ inches when ready for market, green in color, and entirely free from spines, thus making it easy to pick without discomfort. Pods are thicker than Perkins' and fill the crates quickly. Plants are close jointed, and produce an enormous yield. Our strain of this variety is highly bred, uniform, and one of the best, the fields showing practically no variation in plant type of pods. Very well adapted for home gardens and local markets, but does not stand long distance shipments as well as Perkins'. In greatest demand on southern markets.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.



Clemson Spineless Okra



A typical commercial field of Perkins' Spineless Okra at Plant City, Florida

PERKINS' SPINELESS. (53 days.) Pods similar in size, shape and appearance to Perkins' Long Green, pods and plants are practically spineless. Early, productive, makes a semi-dwarf plant 4½ to 5 feet tall. Selected for uniformity in higher yields of dark green, long slender pods desired in the markets. This **Perkins' Spineless** should not be confused with Clemson Spineless, being an entirely different and distinct variety, the pods having the long, slender shape and dark green appearance of Perkins' Long Green. Perkins' Spineless can be sold in any market for Perkins' Long Green.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

DWARF STALK LONG GREEN POD (Perkins' Long Green type). (50 days.) This improved strain of the Perkins' Long Green type is a big improvement in uniformity and yield over ordinary strains of this variety. Developed by several years of careful painstaking selection. Plants average 3½ to 4 ft. tall, close jointed and branching, and loaded with pods from ground up. Pods are long, usually about 3½ to 4 inches when ready for market, straight, slender, tapered, ridged, deep dark green and meaty. Retain their tenderness and green color for a long time in transit and on the markets.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

ONION SETS and SEED

For dry bulbs sow 1½ lbs. in seedbed to plant an acre, or plant 3 lbs. per acre in the field. For the garden sow 1 packet in seedbed for 350 to 450 plants to set 80 feet of row or plant 1 packet to 80 feet of row.

ONIONS FROM SEED

Culture. For large mature bulbs in Florida, first plantings are made the end of August and continued until November. September and October are the best months to plant. The seed is generally planted in the open field, where the onions are to grow, in rows 18 inches apart, and when 4 to 5 inches high they are thinned out from 3 to 4 inches apart, depending on variety and size of bulbs desired. Those taken out can be used to transplant any possible skips. Onion seed can also be planted in seedbeds and transplanted to the field later. This is a safe and sure method, and is becoming more popular in Florida. Transplant when plants are small. The larger the plants when transplanting, the more splits will develop. It is also important to avoid checking plants in growth when transplanting to avoid splits. Do not work soil to onions after they start bulbing. Onions do best in sandy loam or muck soil, and require at least a ton of fertilizer mixture per acre for sandy soils or 1,000 to 1,200 on muck.

The number of days after each variety indicates the time from planting the seed to large mature bulbs under average Florida conditions.

Prices quoted are postpaid.

Write for special prices on larger quantities.

EXCEL (YELLOW BERMUDA 986). (100 days.) Produces more onions of high marketable quality ten to fourteen days earlier than regular Yellow Bermuda. It is practically free of bolters, doubles and splits, and color defects, very uniform in maturity, and in size, color and shape of bulbs, the shape being a deep or thick-flat, the color a light yellow. Crisp and mild in flavor, and keeps better in storage than other Yellow Bermudas.

Pkt. 10c; ½ oz. 20c; 1 oz. 30c; ¼ lb. 85c; 1 lb. \$3.00
5 to 25 lbs. \$2.90 per lb.

GRANEX HYBRID (Yellow). (120 days.) The first short-day F¹ hybrid onion adapted to the South and most nearly fits the requirements for an onion suitable for growing in Florida. Very early, extremely uniform for type, color and maturity, and does not have the large, thick neck but dries down to a small neck. Practically no splits or bolting to seed. In Texas trials, it has out-yielded both its Grano type and Excel type parents. Yields of 600 to 900—50-lb. bags of U.S. No. 1 onions per acre have been reported. The bulb of Granex is short top-shaped or slightly flat, mild in flavor, yet keeps unusually well. On the basis of earliness, total yield, percentage of marketable bulbs, attractive appearance and general all-around good quality, Granex is strongly recommended.

Sold out until about September, 1957



From left to right—Texas Early Grano, Granex Hybrid, Excel.

ONION SETS



Sets for green onions or early maturing bulbs.

Plant 8 to 12 bushels of sets per acre. In the garden 1 pint of sets to 40 feet of row.

A bushel weighs about 32 lbs.

Set out in rows 1½ feet apart and 1 to 2 inches apart in rows for green onions. Plant from September to March.

Prices quoted are f.o.b. Plant City except pints and quarts which are postpaid. 1 bushel weighs about 32 lbs.

	Pint	Quart	Peck	Bushel
White Sets30	.50	1.15	4.00
Yellow Sets30	.50	1.15	4.00

Available from September to March

TEXAS GRANO 502. (130 days.) This strain of early Grano or Babosa was developed by the Texas Agricultural Experiment Station for greater uniformity of maturity and earliness. It is fully ten days earlier, more uniform in shape, and slightly smaller in size than regular Grano. Bulbs have light straw-colored skin, with pure white flesh, are globe- or top-shaped, slightly tapering to the root. Mild in flavor with a sweet taste.

Sold out until about September, 1957

CRYSTAL WAX BERMUDA. (110 days.) Early maturing, flat in shape, of waxy white color, attractive in appearance, and of mild flavor.

Pkt. 10c; ½ oz. 20c; oz. 30c; ¼ lb. 85c; 1 lb. \$3.00
5 to 25 lbs. \$2.90 per lb.

RED CREOLE. (150 days.) Very productive, of medium to large size, semi-globe shaped, quite resistant to thrips. Strong in flavor, but one of the very best keepers. Grows slowly and is late in maturing.

Pkt. 10c; ½ oz. 20c; oz. 35c; ¼ lb. \$1.00 1 lb. \$3.60
5 to 25 lbs. \$3.50 per lb.

NEBUKA (Long White Bunching). (60 days to green onions.) Makes a long stem, much longer than ordinary bunching onions, pure white in color, and without a bulb. Resistant to pinkroot, thrips and smut.

Pkt. 10c; ½ oz. 20c; oz. 35c; ¼ lb. \$1.00 1 lb. \$3.60
5 to 25 lbs. \$3.50 per lb.

For best results use Kilgore's "Bred-Rite" Seeds



Paramount (Improved Moss Curled)

PARSLEY

Plant 1 packet to 150 feet of row, or 2 pounds per acre.
In the garden plant 1 packet to 30 feet of row.

Culture. Parsley can be planted in the fall or spring, in drills 18 inches apart, and thinned to 6 inches in the row. Seed is slow to germinate and it usually takes from two to three weeks to produce a satisfactory stand.

PEAS

ENGLISH OR GARDEN

Our pea seed is high germinating, guaranteed western-grown stock, all carefully fitted for the critical grower.

Plant 1 bushel (56 lbs.) per acre in single rows; 1½ bushels (84 lbs.) per acre in double rows.

In the garden make three plantings at different times.
Plant ½ pound to 100 feet of row at each planting.

Although there are hundreds of distinct varieties of English or Garden Peas, we list only two which we have found particularly well adapted to Florida growing conditions.

Culture. English peas are very hardy and are not commonly attacked by insects. Peas can be grown on almost any kind of soil and with about the same cultivation as beans. Planting in double rows, six to eight inches apart, is a good plan, because this enables the plants to support each other. Single rows should be spaced two and a half to three feet apart, and double rows three to three and a half feet apart from center of double rows. Plant seed 1 to 2 inches apart in the row, and 1½ to 2 inches deep. By working the soil up against the stems of the plants they will bear longer. Use a high grade fertilizer, at the rate of 1,000 to 1,200 pounds per acre. To control soil diseases, and to get better stands, more vigorous growth and increased yields, treat seed with **Spargon Seed Protectant** at the rate of two ounces per bushel of seed (see pages 52 and 62.) Being a legume, the seed should be inoculated with Nitragin. (See page 69.)

The number of days after each variety indicates the relative time from planting to first picking. This time will vary in different sections and in different seasons.

Write for special prices on larger quantities.

HUNDREDFOLD (Kilgore's Winner). (53 days.) This variety was introduced by us and it has become very popular. It is one of the finest large early peas for Florida sandy loam soils, and in cool seasons is also adapted to Florida muck soils. It is a surer cropper and

PARAMOUNT (Improved Moss Curled). (90 days from seed plantings to first harvest.) Plant very vigorous in growth, usually 12 inches tall and spreading 20 inches when given sufficient room, with very long, stout stems for bunching. This variety has been bred and selected for uniform, attractive leaves of very dark green color.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00
5 to 25 lbs. \$1.90 per lb.

PARSNIP

Plant ½ ounce to 150 feet of row.

In the garden plant 1 packet to 50 feet of row.

Culture. Parsnips do best on deep, rich, sandy soil. Sow seed in Florida from September to December, in rows 1½ to 2 feet apart; cover with one-half inch of soil. When plants are well up, thin to 4 inches apart in row. Parsnip seed germinates very slowly. The Parsnip is a winter vegetable with more starch and sugar than other root crops. To prepare, cook roots one hour in water, scrape off the skin. Slice or chop coarsely or mash, and season with milk, pepper and salt, or slices may be fried after seasoning. A delicious and nutritious dish.

HOLLOW CROWN. (120 days from seeding to harvest.) Roots 10 to 12 inches long 2½ to 3 inches in diameter at shoulders.

Pkt. 10c; oz. 20c; ¼ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

a better yielder than other early large podded varieties. The plants are vigorous in growth, eighteen to twenty inches in height, with heavy, very dark green vine and foliage, producing a great abundance of large (4 to 4½ inches long) broad, deep green, pointed pods containing usually eight large dark green peas of fine quality. A very desirable sort for home gardens and for shipping to markets desiring a large podded pea of high quality.

¼ lb. 20c; 1 lb. 60c; 2 lbs. \$1.10, Postpaid

Not prepaid, 14 lbs. (pk.) \$3.75; 56 lbs. (bu.) \$14.00

LITTLE MARVEL. (52 days.) The most popular pea for Florida, because of its good quality, earliness and high yield of well filled pods. The most desirable early variety, especially well adapted to Florida muck soils where it is a sure cropper with well filled pods, even in warm seasons. The vines are vigorous, stocky, dark green in color, growing about 1½ feet tall, and loaded with straight, round, blunt, plump pods about 3 inches long, well filled with 7 to 8 tender peas. A heavy yielder and sure cropper, valuable for home garden and for early shipping.

¼ lb. 20c; 1 lb. 60c; 2 lbs. \$1.10, Postpaid

Not prepaid, 14 lbs. (pk.) \$3.75; 56 lbs. (bu.) \$14.00



Little Marvel, a heavy yielding, early maturing, sure-cropper, especially well adapted to Florida.

PEPPER

Sow $\frac{1}{2}$ to 1 pound of seed in seedbed to plant an acre.
Sow 2 to 3 lbs. per acre when drilled in field.
For the garden sow 1 packet in a protected seedbed for
24 plants to set 40 feet of row.

Culture. Peppers, like eggplants, are usually started in cloth-covered seedbeds. For a crop in central Florida sow the seed in June, July and August; for a long season winter crop on the east coast plant in August, September and October. For an early spring crop in Florida west coast sections sow seed in November and in north Florida in December or January.

High temperature is needed to germinate pepper seed satisfactorily, and it is difficult to get a good stand when the ground is wet and cold. Hammock land, or rather moist soil, is best suited to this crop, although any medium good sandy loam soil, properly fertilized, will grow good peppers. Make rows three feet apart and set plants 10 to 16 inches in the row. As this crop is in bearing for a long time, it is most profitable to make several applications of fertilizer, using 400 pounds per acre about every three weeks until at least a ton has been used.

For the control of insects and diseases see page 55.


The number of days in parentheses after each variety indicates the relative time from setting out plants to picking of marketable green peppers. It requires six to eight weeks to produce plants ready for setting.

WONDER GIANT. (74 days.) This variety of the thick meat Florida Giant type is popular, especially on the west coast of Florida where it has been grown the past few seasons. The vigorous growing plant is more spreading and does not grow tall as Florida Giant. Developed and selected for uniformity in plant and fruit type and productivity. The blocky fruits borne upright on the plant are mostly 4-lobed, with sweet, thick flesh, of deep glossy green color, making an attractive pack.

Pkt. 10c; oz. 60c; $\frac{1}{4}$ lb. \$1.90; 1 lb. \$7.00
5 to 25 lbs. \$6.90 per lb.

YOLO WONDER (Mosaic Resistant Pendent California Wonder Type). (70 days.) This new variety of thick meat pepper possesses a combination of two desirable qualities not found in any other California Wonder type pepper, namely high resistance to tobacco mosaic disease, and in addition it produces almost 100% pendent fruits. With the fruit hanging down they are protected to a great extent by the large, deep green leaves, and hence do not sunburn so badly as other strains of thick meat pepper which produce upright fruits exposed to the sun. A rather short bush, which makes it well adapted to muck soil. The fruit is rather short and blocky, with a tendency to be indented at the blossom end, resembles California Wonder. Yolo Wonder is better adapted for spring growing than fall on sandy soil.

Pkt. 10c; oz. 60c; $\frac{1}{4}$ lb. \$1.90; 1 lb. \$7.00
5 to 25 lbs. \$6.90 per lb.



Yolo Wonder



Harvesting Kilgore's Florida Giant Pepper at Plant City

Prices quoted are postpaid.

Write for special prices on larger quantities.

FLORIDA GIANT. (75 days.) This variety, introduced by us, produces an abundant yield of large, heavy, thick walled, fleshy, dark green fruits, over a long period, but requires heavy and frequent fertilization, and a rich, moist soil. The tall plants are compact but very vigorous and upright in growth. Slightly later than World Beater, they are heavy producers, often bearing six to eight large fruits at one time. The fruits are longer than most thick-meated peppers, about 4 to 4½ inches long, and 3 to 3½ inches in diameter, usually with four lobes, are smooth and well filled at the blossom end. Makes a very attractive pack and carries well to distant markets.

Pkt. 10c; oz. 60c; $\frac{1}{4}$ lb. \$1.75; 1 lb. \$6.50
5 to 25 lbs. \$6.40 per lb.



California Wonder

CALIFORNIA WONDER. (75 days.) An outstanding medium-late variety. Plant is vigorous, upright, prolific. The thick meated, smooth, large, heavy fruits average about four inches in length and about the same in diameter. This is an excellent shipping type, holding up well in transit and on the market, and is in great demand by truckers. Solid thick walls and very sweet and mild. Blocky in shape. Color an attractive, glossy dark green. Requires frequent fertilizing and continuous moisture to hold the size of the fruit up over a long harvesting period.

Pkt. 10c; oz. 60c; $\frac{1}{4}$ lb. \$1.75; 1 lb. \$6.50
5 to 25 lbs. \$6.40 per lb.



Early Calwonder

EARLY CALWONDER. (65 days.) An early maturing strain of California Wonder, maturing 7 to 10 days ahead of other strains, and producing fruits over a long season. The plant is not so tall and erect in growth as regular California Wonder, but is very vigorous and prolific. The fruits usually are four-lobed and attractive in appearance, being smooth, dark green, uniform and thick meated. This early strain is rapidly growing in popularity.

Pkt. 10c; oz. 60c; ¼ lb. \$1.90; 1 lb. \$7.00
5 to 25 lbs. \$6.90 per lb.

WORLD BEATER (Thick Wall Strain). (70 days.) This strain of World Beater Pepper was developed in our breeding station at Bradenton, Florida, for a more uniform, thicker walled, heavy yielding strain. In numerous trial plantings in different sections of Florida we have found this to be a heavy yielder of uniform, true to type, smoother, thicker walled fruits, which are larger in size and carry better to distant markets than regular World Beater which usually are very thin-walled. In some sections it has brought a premium on the market. The plants are erect in growth and hold the large heavy fruits well off the ground.

Pkt. 10c; oz. 60c; ¼ lb. \$1.75; 1 lb. \$6.50
5 to 25 lbs. \$6.40 per lb.




World Beater Thick Wall Strain

ITALIAN "EL" (Green Neapolitan). (65 days.) This green-fruited strain of Neapolitan is in increasing demand on northern Italian markets where it is often preferred to the regular yellow-fruited Neapolitan because of its peculiar sharp taste. Fruits are slightly larger and thicker fleshed than the regular yellow Neapolitan. The vigorous growing plants produce a heavy yield of light green fruits over a long season.

Pkt. 10c; oz. 60c; ¼ lb. \$1.90; 1 lb. \$7.00
5 to 25 lbs. \$6.90 per lb.

NEAPOLITAN (Italian). (50 days.) This is an early, medium large, prolific variety, growing in popularity on some northern markets where it is in demand among Italian people. This plant is wonderfully productive, carrying as many as forty fruits on a single plant. The fruits are upright in growth, averaging 4 inches long and 2 inches in diameter, three-lobed, yellowish green, becoming bright red when ripe. The flesh is mild and sweet.

Pkt. 10c; oz. 60c; ¼ lb. \$1.90; 1 lb. \$7.00
5 to 25 lbs. \$6.90 per lb.



Neapolitan (Italian)

ANAHEIM CHILI. (80 days.) A popular, rather long, mildly pungent variety, not as hot as the small fruited Long Thin Cayenne. The fruits are meaty, 6 to 7 inches in length and 1 to 1½ inches in diameter at stem end, green, ripening to a bright red. Often dried for winter use.

Pkt. 10c; oz. 55c; ¼ lb. \$1.65; 1 lb. \$6.00
5 to 25 lbs. \$5.90 per lb.

HUNGARIAN YELLOW WAX. (60 days.) A very early, hot sort, valuable for canners. Plants are rather dwarf in growth, and exceedingly prolific. Fruits pendent, medium slender, long, tapering, smooth and very hot or pungent. Color waxy yellow, changing to bright crimson at maturity.

Pkt. 10c; oz. 65c; ¼ lb. \$2.15; 1 lb. \$7.50
5 to 25 lbs. \$7.40 per lb.

LONG THIN CAYENNE. (70 days.) This favorite hot variety produces large and productive plants. Fruit 4 to 5 inches long, ½ inch thick, slender, tapering, pendent, very hot or pungent, deep green in color, ripening to a brilliant red. Used largely for canning and in pickles.

Pkt. 10c; oz. 60c; ¼ lb. \$1.75; 1 lb. \$6.50
5 to 25 lbs. \$6.40 per lb.


SEED POTATOES

Plant 600 to 900 pounds (10 to 15 bu.) per acre in 3-ft. rows. In the garden plant 12 pounds to 200 feet of row.

Culture. Plant in rows 36 to 42 inches apart, dropping a seed piece every nine to ten inches in the row. The seed piece should be at least one to one and one-half ounces in size. It is best to cut seed within one day of planting. Fertilizer should be put in seven or eight days before planting. Use from 1,500 to 2,000 pounds per acre, depending on type of soil. A 4-7-5 fertilizer mixture is a good standard for potatoes. On rich Everglades muck from 200 to 500 pounds per acre of 0-10-12 or 0-8-16 mixture applied in one application at planting time is a common practice. To control blight and increase the yield, spray with Dithane D-14, using 2 quarts Dithane D-14, $\frac{3}{4}$ lb. zinc sulphate, to 100 gallons water. For control of insects and diseases see page 55.

RED PONTIAC (Dakota Chief) (State Certified). (80 days.) Packed in 100-lb. bags. This variety is the result of a cross, Red Bliss and Katahdin, made by the Michigan Experiment Station. In tests at the Sub-tropical Experiment Station in Homestead and elsewhere in Florida, it has greatly outyielded Red Bliss. The tubers of Red Pontiac are slightly longer, and have shallower eyes, but the color is a bright red like Red Bliss. Pontiac tubers are tougher, stand shipment better and are better keepers than Bliss.

Write Plant City or your nearest Kilgore store for price on seed potatoes stating quantity wanted.



Red Pontiac

PUMPKIN

Plant two pounds seed per acre.

In the garden plant $\frac{1}{2}$ ounce to 50 feet.

Culture. Plant 4 to 5 seeds in hills 8 to 10 feet apart one way and 5 feet apart the other way, when the weather gets warm. Can be planted in corn. Thin to two or three plants after well started. Give shallow and frequent cultivation until vines begin to run. Cut pumpkins when fully ripe.

Prices quoted are postpaid.
Write for special prices on larger quantities.

KING OF THE MAMMOTHS. (100 days.) Commonly known as a pumpkin, but is really the very large fruited Mammoth Chili squash, known as the giant among squashes. Fruits are globular, slightly ribbed, light yellow color mottled with orange. Flesh is solid, light yellow-orange, coarse, of fair quality. Grown mainly for stock feed. Fruits measure 20 to 22 inches across and weigh 60 to 80 pounds.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.



Small
Sugar

SMALL SUGAR. (110 days.) The best variety for general use, and particularly desirable for pies. Fruits are round, flattened at ends. Skin is hard, smooth, somewhat ribbed, of deep orange color. Flesh is thick, sweet and dry, of bright orange color, and very high quality. Makes delicious pumpkin pies. Fruits measure eight to ten inches and weigh six to eight pounds.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

CORNFIELD. (118 days.) Prolific and vigorous, usually planted in cornfields. Extensively grown for making pies, for canning and for stock feed. Fruits are large, round to slightly flattened, with a hard, smooth surface, slightly ribbed, deep yellow or orange color. Fruits measure 12 to 14 inches and weigh 15 to 20 pounds.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

RHUBARB

Sow 3 to 4 pounds in seedbed for an acre of plants, or plant 5 to 6 pounds per acre in field.

For the garden sow a packet in seedbed for 15 plants to set 25 feet of row.

Culture. Plant in the fall in rows $2\frac{1}{2}$ to 3 feet apart and thin to about 18 to 24 inches in the row. It is best to plant seed in seedbed and transplant to field or garden. Rhubarb does best in well drained, fertile, moist soil, rich in humus. In Florida it is necessary to plant Rhubarb each season.

VICTORIA. (90 days from plants set in garden to edible stalks.) This vigorous growing variety produces thick, fleshy stalks of the best quality and flavor, and with attractive red color. It is the best variety for Florida.

Pkt. 10c; oz. 30c; $\frac{1}{4}$ lb. 85c; 1 lb. \$3.00
5 to 25 lbs. \$2.90 per lb.

ROSELLE

Jamaica Sorrell or Florida Cranberry
(175 days from seed planting to ripe fruits.)

Plant 3 pounds per acre. In the garden plant one packet to 25 feet of row.

The red fruit is used when young for making jelly and jam. It is grown commercially for local markets, fruits being marketed in quart berry baskets.

Culture. Plant seed in field in March and April in Florida, when the ground is warm, spacing row 5 to 6 feet apart and thinning to about $2\frac{1}{2}$ feet apart in the row, leaving one strong plant at a place. Roselle is an all-season crop requiring from five to six months from seed to ripe fruits.

Florida Cranberry

Pkt. 10c; $\frac{1}{2}$ oz. 25c; oz. 45c; $\frac{1}{4}$ lb. \$1.40; 1 lb. \$5.00
5 to 25 lbs. \$4.90 per lb.

RADISH

Plant 6 to 8 pounds of seed per acre.

In the garden make four plantings at different times. 1 packet to 25 feet of row at each planting.

Our Radish Seed is graded and sized for more accurate planting, better stands and more uniform maturity. Plant Kilgore's graded radish seed for best results.

Culture. Plant in rows 12 to 18 inches apart and thin to 1 or 2 inches in the row. Successive plantings should be made every few weeks for a continuous supply. For best results, a rich, loose, moist sandy loam or muck soil is required. Radishes must be grown rapidly to be crisp, tender, and of best quality. Our radish seed stocks are produced each year from carefully selected transplanted roots. For insect and disease control see page 55.

Prices quoted are postpaid.

Write for special prices on larger quantities.

EARLY SCARLET GLOBE. (21 days.) We have a perfected stock of this early variety. It is the earliest of all varieties; roots are globe-shaped, with a medium top. They are crisp and tender, of uniformly attractive bright, scarlet color. In great demand on all markets.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

SCARLET TURNIP WHITE TIPPED (Sparkler). (22 days.) A superior strain of this popular early variety. The roots are nearly round, being slightly flattened on the underside, deep scarlet, shading to a white tip on the bottom. They are very solid, crisp, tender, and sweet. Attractive in appearance.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

BUCCANEER (Scarlet Globe Type). (21 days.) A new development of the Scarlet Globe type, developed for uniformity and attractive market appearance. The root is globe shaped, with a lustrous scarlet exterior, and with short top growth. Crisp-textured and a most uniformly attractive market radish.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.



Early Scarlet Globe Radish

LONG WHITE ICICLE. (23 days.) The leading long radish. The straight roots are from 5 to 6 inches long, cylindrical, and very smooth and waxy white throughout. The flesh is mild, crisp, tender and of fine quality.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

SPINACH


Virginia Savoy—Plant 12 to 15 pounds per acre in rows, or 20 to 25 pounds per acre broadcast.

In the garden make two plantings at different times. Plant one packet to 50 feet of row at each planting.

All prices quoted are postpaid.

Write for special prices on larger quantities.

Culture of Virginia Savoy Spinach. Successful for fall and winter planting. Does not grow well in hot weather. Requires a moist rich soil. Plant seed in rows 16 to 20 inches apart, and thin to 6 inches apart in the row.



Virginia Savoy

VIRGINIA SAVOY (Blight Resistant). (40 days.) Better for market than the smooth leaved varieties, because it does not pack as closely and thus is not as subject to heating as are the smooth leaved varieties, nor does not wilt or turn yellow as quickly before reaching the consumer. Developed for resistance to blight or mosaic disease. The plant is upright and compact in growth, with very thick, curly or savoyed dark green leaves.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. 90c
5 to 25 lbs. 90c per lb.

New Zealand—Plant 3 to 4 pounds of seed per acre. In the garden plant one packet to 25 feet of row.

Culture of New Zealand Spinach. As seeds are very hard and germinate with difficulty, soak them in warm water for 24 hours to hasten germination. Plant in rows three feet apart, and drop three or four seeds per hill about 20 inches apart. Thin to one plant per hill.

NEW ZEALAND. (60 days.) Entirely distinct from the true Spinach. The plant is spreading, with numerous side shoots. It grows to a height of 1 foot and spreads over a wide radius. Leaves are thick, light green, rather small, broad and pointed. It grows well in hot weather, and is sometimes called "Summer" or "Hot-weather Spinach." The leafy tips of the branches are removed and new growth follows, making possible many cuttings. It does best on moist, rich land.

Pkt. 10c; oz. 15c; ¼ lb. 40c; 1 lb. \$1.10
5 to 25 lbs. \$1.00 per lb.

SAGE

A popular garden herb used in flavoring or seasoning. Leaves should be cut as plant blooms, and should be dried quickly in the shade. Grows 18 inches tall.

Culture. Sage does well on a sandy soil. Plant seed in rows 15 to 18 inches apart. Thin to six inches in the row. The plant is a hardy perennial, and of easy culture in Florida.

Pkt. 10c; ¼ oz. 35c; 1 oz. \$1.00

SQUASH


Culture. This is one of the quickest and easiest crops that can be grown, and usually proves very profitable. The earliest varieties begin bearing in six or seven weeks from seed. Squashes can be grown on almost any kind of soil. Use from 800 to 1,000 pounds of fertilizer per acre.

For bush varieties, have rows at least 3 to 4 feet apart and hills 2 feet apart in the row. Plant 3 to 5 seeds in each hill, and thin out to two plants when 3 inches high. Bush squash in Florida is a good paying crop for fall or spring. Plant for spring, during January, February and March; for fall, from August until October. Mosaic disease has become a most serious disease of squash in many sections of Florida. See Sanitation on page 50. Insect and Disease Control on page 54.

The number of days after each variety represents the average time required from seed planting to edible or marketable fruits.

All prices quoted are postpaid.

Write for special prices on larger quantities.



Cocozelle Squash, or Long Slender Italian Vegetable Marrow

BUSH VARIETIES

Plant 3 to 4 pounds of seed per acre.

In the garden one packet to 50 feet.


COCOZELLE (Italian Vegetable Marrow). (50 days.) This is a true Italian Vegetable Marrow. When ready for market, the fruit is 10 to 12 inches long, slender, 1 to 1½ inches in diameter, cylindrical in shape, straight, smooth, dark green with lighter green stripes.

Our stock of this variety produces a bushy, open type of plant, with deeply cleft leaves, and a heavy yield of uniformly true-to-type fruits. We have devoted many years to the development of this superior strain, and we can say with full confidence that our Cocozelle is the earliest, most prolific, and uniformly true-to-type strain you can secure anywhere.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.

EARLY PROLIFIC STRAIGHTNECK. (42 days.) This development is ten days earlier and more productive than other strains of Straightneck Squash. The plants are vigorous, and the fruits are of brighter lemon-yellow color. Valuable for both home and market use. Because of its uniformity, earliness, high yield and attractive appearance this variety has grown greatly in popularity especially in northern markets. Makes a very attractive pack.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.




Early Yellow Summer Crookneck or Baby Crookneck

EARLY YELLOW SUMMER CROOKNECK (Baby Crookneck). (45 days.) A very early and prolific, light lemon-yellow color strain with an open type of plant growth. The most popular and most extensively grown of any variety of squash in Florida. Fruits are small with curved neck, and uniform. Its rich, bright, lemon-yellow color gives it a very good market appearance, and it is in demand on all southern markets. Our strain of this variety is very early and exceedingly prolific, and has been selected for uniformly shaped fruits with a pronounced crook. No better strain of this most popular variety can be secured anywhere for Florida planting.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.

BLACK ZUCCHINI (Kilgore's Improved Selection). (45 days.) The plant of this improved selection, instead of developing a large, vigorous growing bush like the regular Black Zucchini, forms a short, single semi-runner, and produces a heavy yield of fruit over a long period. The plant is an open type of growth, with deeply cleft leaves, allowing the air and sunlight to circulate and prevent fruit rot. The fruits are dark green in color, six to eight inches long at market stage. A very early, prolific, improved strain of Black Zucchini.

Pkt. 10c; oz. 20c; ¼ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.



Early Prolific Straightneck

SQUASH—BUSH VARIETIES (Continued)

WOOD'S EARLY PROLIFIC (Patty Pan Type). (42 days.) The earliest, meatiest, and most prolific of the patty pan type. It will average seven days earlier than Early White Bush, matures its fruits quickly. The plants are prolific, producing round, smooth, deep, thick fruits, slightly scalloped, and of creamy-white color, usually with a light greenish cast, being bulged at both stem and blossom ends. A very good variety for home use or for shipment.

Pkt. 10c; oz. 15c; $\frac{1}{4}$ lb. 45c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.

WHITE BUSH SCALLOP (EARLY WHITE). (48 days.) This is a popular type of squash for shipment to northern markets. The smooth fruits are of pale green color when very young, becoming white as they grow, are round and flat but deep, with ridged or scalloped edges, measuring up to five or six inches across by three inches thick when ready to market.

Pkt. 10c; oz. 15c; $\frac{1}{4}$ lb. 45c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.



White Bush Scallop

MAMMOTH WHITE BUSH (Patty Pan Type). (50 days.) A large strain of the Early White Bush, fruits being about one-third larger, measuring to 8 inches in diameter. It is slightly later in maturing and not quite so prolific as Early White Bush. The fruits are thick, tender, and of good quality.


Pkt. 10c; oz. 15c; $\frac{1}{4}$ lb. 45c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.

SQUASH - RUNNING VARIETIES

Plant two pounds per acre of running varieties
In the garden one packet to 50 feet.

Culture. Cultural requirements are similar to bush squash (see page 30), but running varieties require much more room, planting in rows 6 to 8 feet apart and hills 3 to 4 feet apart. Plant 3 to 5 seeds in each hill and thin to two plants when three inches high. Generally grown as a spring crop in Florida.

In these types of squash the vines run 10 to 12 feet. The fruits are left on the vines until fully grown with a hard rind, and are then harvested by cutting the stem. Running squash may be stored in a cool, dry place until desired for use, but must be handled carefully to avoid bruising. Unlike bush squash, fruits of the running varieties may be stored and used boiled, baked or for making squash pies.



Butternut

BUTTERNUT. (80 days.) Vines produce pear-shaped yellowish brown or deep buff-colored fruits. Fruits weigh $2\frac{1}{2}$ to 4 lbs., 8 to 10 inches long, and 3 to 4 inches in diameter at the largest diameter. The outer shell is smooth and hard, making this squash a wonderful shipper and keeper. The dry flesh is yellow, fine-textured, sweet, and of excellent flavor. Also excellent for making pies.

Pkt. 10c; oz. 25c; $\frac{1}{4}$ lb. 75c; 1 lb. \$2.50
5 to 25 lbs. \$2.40 per lb.

ALAGOLD (Improved African Squash). (100 days.) Resistant to most insects and diseases which commonly attack squash in the South. Average about 5 pounds in weight. The rind is of a dark green color, maturing to a deep yellow or buff color. A very good keeper.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 60c; 1 lb. \$2.00
5 to 25 lbs. \$1.90 per lb.

ROYAL ACORN (MAMMOTH TABLE QUEEN). (65 days.) To meet an increasing demand in Florida for a larger fruited strain of the Table Queen or Acorn Squash, we are offering this large-fruited strain. It is slightly later than regular Table Queen, not quite so prolific or as uniform in shape and color but the fruits are consistently larger and of a darker green color. In other respects this strain is similar to the regular Table Queen.

Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 50c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.

TABLE QUEEN (ACORN). (60 days.) Extremely early for a running squash. Makes an enormous yield over a long season. Fruits dark green, weighing about 2 lbs., with sides deeply ribbed, and with a point at the blossom end. Yellow-colored flesh is very fine-grained and sweet when baked like Hubbard squash. It is a very good keeper and desirable for home and market use, also for shipping.

Pkt. 10c; oz. 15c; $\frac{1}{4}$ lb. 45c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.




Table Queen (Acorn)

TOMATO

Sow $\frac{1}{2}$ pound in seedbed to plant an acre, or 1 pound per acre in field.
For the garden sow one packet in a seedbed for 50 plants to set 150 feet of row.
If plants are staked, sow one packet in seedbed for 50 plants to set 75 feet of row.

Culture. Tomatoes are grown on all kinds of soil, from a light sand to a muck. The land should be put in good condition. Lay off rows $3\frac{1}{2}$ to 5 feet apart, open a furrow, and drill in 1200 pounds per acre of high grade fertilizer such as 4-7-5, and mix thoroughly with the soil. It is good policy to let this stand for seven to ten days and then plant seed or set the plants. Seed may be planted direct in the field or in a cloth-covered seedbed. Plants started in a seedbed should be 6 to 8 inches high when ready to transplant to the field in 4 to 7 foot rows and $2\frac{1}{2}$ to 3 feet apart in a row, or if staked 16 to 20 inches in the row, rows 4 feet apart.

On the east coast of Florida, first plantings are made during August and continue until January. North and central Florida begin planting for a spring crop the middle to the latter part of December and continue to the middle of February. For a fall crop in central Florida, plantings are made in July and August.

For control of insects and diseases of Tomatoes see page 56. Spraying or dusting tomato plants with Dithane before the disease appears will help prevent the destructive late blight disease.

The number of days indicated after each variety represents the time required from setting of plants in the field to produce marketable fruits. It usually takes four or five weeks to produce plants for field setting.

*All prices quoted are postpaid.
Write for special prices on larger quantities.*




Jefferson (Wilt Resistant)

JEFFERSON (Fusarium Wilt Resistant). (76 days.) The plant is very vigorous and makes densely-leaved tall vines. It is more open and less subject to bunching than Rutgers. Produces well in the fall but is best suited as a spring tomato on wilt infested soils. The fruits are globe-shaped, ripening to a uniform, attractive dark red color, with a good flavor, thick walled with a small core, and relatively free from cracking and catfacing. This variety holds up under adverse conditions better than any other variety. A good shipper as a green wrap tomato. Jefferson also has become a favorite for pink harvest, has good carrying qualities and will stand rougher handling when shipped as a vine ripe than most other varieties.

**Pkt. 10c; $\frac{1}{2}$ oz. 35c; oz. 60c; $\frac{1}{4}$ lb. \$1.75; 1 lb. \$6.50
5 to 25 lbs. \$6.40 per lb.**

PRITCHARD (Scarlet Topper). (75 days.) Vines medium dense, compact, erect, self-topping. Fruit matures very early, large, smooth, meaty, globular in shape. This variety is a heavy cropper and therefore requires rich or liberally fertilized soil. Does especially well on Everglades muck.

**Pkt. 10c; $\frac{1}{2}$ oz. 35c; oz. 60c; $\frac{1}{4}$ lb. \$1.85; 1 lb. \$6.75
5 to 25 lbs. \$6.65 per lb.**



Homestead 24 (Wilt Resistant)

HOMESTEAD 24 (Wilt Resistant) (new). (70 days.) A selected strain of Homestead tomato. Our aim in developing this strain has been elimination of off types in the original Homestead leading to uniformity of plants and fruits and a better crop. Compared with Rutgers, the plants of Homestead 24 are similar but semi-determinate, earlier in maturing and resistant to Fusarium Wilt. Very free from puffiness and cracking, meaty and firm, excellent for green wrap. Homestead 24 is a heavy yielder and will make good crops even on wilt-infested soils. It has done well in the Homestead area, both on heavy marl soil and on the rockland, and seems to be adapted to all Florida soils they have been grown on.

**Pkt. 10c; $\frac{1}{2}$ oz. 40c; oz. 75c; $\frac{1}{4}$ lb. \$2.25; 1 lb. \$8.00
5 to 25 lbs. \$7.90**

RUTGERS. (82 days.) Our improved strain is uniform in plant growth and produces uniform smooth, thick wall, deep globe-shaped fruits which ripen to a bright red. The plant is vigorous and a heavy producer. Very firm and free of puffs. Fruits size up well out to the end of the vine. Recommended for a spring crop in Florida and is well suited for light sandy soils and also for shallow, dry pine land on the lower east coast of Florida. Not suitable for deep, wet glade lands where it makes too much vegetative growth.

**Pkt. 10c; $\frac{1}{2}$ oz. 30c; oz. 55c; $\frac{1}{4}$ lb. \$1.60; 1 lb. \$5.75
5 to 25 lbs. \$5.65 per lb.**



Rutgers Tomato

GROTHEN'S GLOBE. (70 days.) An early maturing, high yield variety especially well adapted for fall planting in Florida. The plants are of open growth with rather small foliage requiring rich, deep, moist land for best results, maturing fruits about 12 days earlier than Rutgers, and ripening to a beautiful red color. A heavy producer of smooth, deep globe-shaped, large fruits, sizing up well clear to the top of the plants, producing a high percentage of fancies. The fruits are firm and solid, and the interior ripens slowly, making this variety an exceptionally good shipper. Earliness and high yield of large fruits, of excellent shipping quality, are its main factors. It does not do so well on light sandy soils, but is especially recommended for a fall, winter and early spring crop in Florida and is well suited to deep, wet glades, hammock, and rich sandy loam soils.

Pkt. 10c; ½ oz. 30c; oz. 55; ¼ lb. \$1.65; 1 lb. \$6.00
5 to 25 lbs. \$5.90 per lb.



Kokomo (Wilt resistant Rutgers type)

WILT RESISTANT GROTHEN'S GLOBE. (70 days.) This variety has been developed from a cross of Pan American and regular Grothen's Globe. Wilt Resistant Globe possesses all the good qualities of the regular Grothen's Globe, and in addition it is very resistant to Fusarium Wilt, although slightly earlier and more vigorous than regular Grothen. In numerous comparative trials between Wilt Resistant and regular Grothen's Globe, even where wilt was not a factor, the Wilt Resistant strain yielded just as high as the regular strain. This Fusarium Wilt Resistant strain is especially recommended for old, wilt infested soils where the regular Grothen's Globe and other varieties cannot be grown successfully. Is not resistant to all wilts, but is highly resistant to the most serious wilt disease of tomatoes caused by the Fusarium soil fungus.

Pkt. 10c; ½ oz. 35c; oz. 65c; ¼ lb. \$2.15; 1 lb. \$7.50
5 to 25 lbs. \$7.40 per lb.



Grothen's Globe

MARGLOBE. (77 days.) This is an excellent strain of the old standard type Marglobe. It produces a vigorous plant resistant to nailhead rust. Fruit deep globe-shaped, smooth and thick walls, free of navel and blossom-end scars. Ripens to a beautiful deep scarlet-red color. Well adapted to and recommended for north Florida.

Pkt. 10c; ½ oz. 35c; oz. 60c; ¼ lb. \$1.75; 1 lb. \$6.50
5 to 25 lbs. \$6.40 per lb.

KOKOMO (Wilt resistant Rutgers type) (new). (80 days.) As a result of numerous trials in different sections of Florida during the past few years we are fully convinced that this new, Fusarium Wilt resistant variety of the Rutgers type is exceedingly well adapted to Florida growing conditions. It will do well wherever Rutgers does well. It is of the general Rutgers type, not as vigorous or bunchy in growth as Rutgers. Vines open up more, making it easier to spray or dust and to harvest. Kokomo sets a heavy crop of good size, smooth, deep globe-shaped, meaty, thick-walled fruit, is a good shipper. It makes a very high percentage of smooth, fancy fruits and it is claimed to set well in hot weather. Is particularly valuable where Fusarium Wilt is a serious crop hazard, under which conditions Rutgers cannot be successfully grown.

Pkt. 10c; ½ oz. 35c; oz. 65c; ¼ lb. \$2.15; 1 lb. \$7.50
5 to 25 lbs. \$7.40 per lb.

MANALUCIE (Wilt Resistant) (new). (82 days.) This new, vigorous growing, disease resistant variety was developed and recently released by the Gulf Coast Experiment Station at Bradenton, Fla. It possesses multiple resistance, being very resistant to four serious Florida tomato diseases, namely, Fusarium Wilt, gray leaf spot, early blight, and leaf mold. It is also mildly resistant to black spot stem canker caused by the Phoma fungus. Makes a very strong, vigorous growing plant with large leaves and large, strong stems. Fruits are late, but are large size throughout the harvest period. Smooth and relatively resistant to cracking. Firm and fleshy, and good edible quality. Very slow ripening and must be allowed to fully mature before harvesting. Good for pink harvest.

Pkt. 15c; ½ oz. 40c; oz. 75c; ¼ lb. \$2.25; 1 lb. \$8.00
5 to 25 lbs. \$7.90 per lb.

PONDEROSA (Brimmer). (88 days.) A large, pink-fruited variety. Fruits semi-flat, solid with few seeds, and of splendid edible quality. Recommended only for home gardens.

Pkt. 10c; ½ oz. 40c; oz. 75c; ¼ lb. \$2.50; 1 lb. \$9.00
5 to 25 lbs. \$8.90 per lb.

OXHEART. (90 days.) A smooth, thick, tender and almost seedless pink-fruited tomato of delicious flavor. An ideal variety for the home garden and for the local market.

Pkt. 15c; ½ oz. 65c; oz. \$1.25; ¼ lb. \$4.00; 1 lb. \$15.00
5 to 25 lbs. \$14.90 per lb.

What Will Your Harvest Be ?

The answer to that question depends principally on the kind of seeds you plant. Kilgore's Bred-Rite Seeds are Bred and Developed in Florida for Florida. This is why The Kilgore Seed Co. is referred to as "Florida's Leading Seedsmen."



Partial view of Experimental Farm and Breeding Station at Bradenton, Florida.

Samples of different stocks of Kilgore seeds are planted from time to time on an Experimental Farm, at Bradenton, Florida. In addition, there also are included in these trials, samples from other seed dealers and seed growers to compare with Kilgore stocks. This shows just how Kilgore's strains and varieties compare in earliness, yield, uniformity, trueness to type and adaptability with others growing side by side under identically the same conditions in Florida. It also enables the company to develop new or improved varieties of distinct merit for Florida conditions, which we offer to our customers as soon as a supply of seeds can be produced. Modern, scientific breeding methods are employed both in maintaining and improving seed stocks. New products are carefully tested under Florida conditions, and superior stocks are then sent to our seed production fields in the North and West. Careful notes are kept on all experimental farm results

which enable us to check up with the constant improvements which are being made in standard varieties of vegetables. Our principal objective is to breed, develop or select strains and varieties of vegetable seeds adapted to Florida growing conditions, which differ materially from those found elsewhere in the United States. A well trained and experienced plant breeder supervises this work.

In addition, the Kilgore Seed Company personnel is in constant contact with the scientific research workers of the Florida Agricultural Experiment Station and field laboratories throughout the state where plant pathologists, horticulturists and plant breeders are doing a splendid work in developing new and improved vegetable varieties especially adapted to the climate and growing conditions of Florida, which we introduce to Florida growers as soon as available.

HIGH VITALITY SEED MUST BE PLANTED TO SECURE A GOOD CROP

Kilgore's vegetable seeds are air-conditioned and tested.

This means higher vitality, better germination and stronger plants.

The Kilgore Seed Company, has in its central warehouse in Plant City, Florida, the most up-to-date dehumidifying machinery in order to maintain high vitality in all Kilgore seed. Surplus seed supplies in our stores are also kept in specially constructed dry rooms where the humidity is maintained at a uniformly low degree.

Many of our vegetable seeds, such as cabbage, celery, eggplant, onions, pepper and tomato are now Vigor-pak by Associated Seed Growers. Asgrow "Vigor-pak" seed must first have a high germination and vitality standard. The seed is scientifically pre-conditioned, treated and sealed in air-tight cans. The cans should be opened only as needed for planting. In unopened cans the seed will retain its vigor and

germination even under Florida conditions of high humidity and temperature.

In addition to keeping our seeds in artificially dried storerooms, we have a seed testing laboratory where samples of all our seed stocks are tested at regular intervals to further insure our customers against planting low vitality seed. In our seed testing laboratory, tests are made every six weeks of all stocks in our warehouse, mail order department and fourteen Florida stores. We know at all times the correct germination of all seed we have on hand. If any seed stocks are found to be dropping in germination they are immediately removed from our warehouse and stores and are destroyed.

We are taking all these precautions and giving our customers this extra service without any additional cost, in order to provide in Kilgore's Bred-Rite Seeds the best that money can buy. Play safe and get your seed from the Kilgore Seed Company, Florida's leading seedsmen.



VIGORPAK SEED. Your insurance against loss in germination and vigor • Loss from pests • Waste from mixture

TURNIP

Plant 2 to 3 pounds of seed per acre.
In the garden make three plantings at different times.
1 packet of turnip seed to 50 feet of row.

Culture. Plant in rows 15 to 20 inches apart. Thin early to one inch, because any over crowding will result in misshapen, slow growing, poor quality roots. Final thinning should be 4 to 6 inches apart. Thinnings may be used for turnip greens. For best quality, turnips should be grown rapidly in rich, moist soil. When turnips are grown for commercial canning and freezing, where only the tops are used and harvested with machinery, plant in double rows 4½ to 5 feet apart, 14 inches apart in the double row. 6 to 8 lbs. of seed is required, no thinning. Turnips can be planted every month in the year.

*Prices quoted are postpaid.
Write for special prices on larger quantities.*

TOKYO MARKET. (35 days.) Extremely early in maturity. Attractive, white, globe-shaped roots. Small foliage grows upright, and is somewhat aphid resistant. Very uniform, which makes it a good variety to grow for market. Exceptionally fine quality.

Pkt. 10c; oz. 15c; ¼ lb. 40c; 1 lb. \$1.25
5 to 25 lbs. \$1.15 per lb.

PURPLE TOP MILAN (EARLY FLORIDA). (35 days.) The roots are of medium size, smooth, flat, and white, with a bright purple top, and extremely early. Recommended for home garden and local market variety.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.



A typical root of Purple Top White Globe

WHITE GLOBE. (44 days.) A large, round turnip of fine quality. It has small tops, grows rapidly, and gets very large, producing a heavy yield per acre. Resembles Purple Top White Globe except for color.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00
5 to 25 lbs. 90c per lb.

SHOGIN (FOLIAGE) TURNIP. (50 days.) An aphid-resistant foliage turnip which resists the hot, dry weather. It produces an immense yield of upright growing foliage, excellent for freezing and canning. The roots are white, semi-globe shaped, but more largely planted for the tops which are large, thick, succulent, tender, and of finest mild flavor.

Pkt. 10c; oz. 15c; ¼ lb. 30c; 1 lb. 75c
5 to 25 lbs. 65c per lb.

PURPLE TOP WHITE GLOBE. (50 days.) Ours is a very uniform and well colored strain of this popular variety. Roots globe-shaped, flesh and skin are white, with purple or red at the stem end above ground. Fine grained and sweet, of superior quality. Grown extensively in Florida for shipment to northern markets.

Pkt. 10c; oz. 15c; ¼ lb. 30c; 1 lb. 75c
5 to 25 lbs. 65c per lb.

WHITE EGG. (45 days.) An early, oval or egg-shaped variety with clear white roots of medium size. The flesh is white, firm, fine grained and sweet.

Pkt. 10c; oz. 15c; ¼ lb. 30c; 1 lb. 80c
5 to 25 lbs. 70c per lb.

RUTABAGA

Plant 1 pound per acre.
In the garden plant one packet to 100 feet of row.

Culture. Rutabaga is best grown as a late fall and winter crop. Planted in rows 2 feet apart, and the plants thinned to about 5 to 6 inches in the row.

AMERICAN IMPROVED. (90 days.) The smooth roots attain a much larger size than turnips, slightly oblong, pale yellow with purple top, and with a very small neck. Flesh is yellow, solid, tender.

Pkt. 10c; oz. 15c; ¼ lb. 30c; 1 lb. 75c
5 to 25 lbs. 65c per lb.

WATERMELON

Plant 1 pound of seed per acre.

In the garden plant 1 packet to 75 feet of row for a family of 3 to 4 people.

Culture. Watermelons may be grown on nearly all kinds of Florida soils that have good drainage, but rolling, sandy pine land is preferred. In general, new land is best for this crop in order to avoid diseases. The field should be checked off, planting in hills 8 feet apart each way or 7 by 9 feet. Plant $\frac{1}{2}$ to 1 inch deep, and plants thinned to 2 or 3 plants per hill when they are still small and when all danger of frost is past, later thinning to 1 plant per hill. The most successful growers apply 600 to 800 lbs. of fertilizer mixture about a week before planting the seed. A second application of 400 to 600 lbs. of fertilizer or top dressing may be made if necessary when the vines begin to run. The first planting of watermelons for an early crop in south Florida starts early in December. In central Florida planting begins January 1st and extends to about February 15th. Plantings in north Florida are made 2 to 3 weeks later. For earlier melons use **Hotkaps**. (See page 75).

The leaf, stem and fruit disease of watermelons, anthracnose as well as gummy stem blight, the spores of which disease are frequently present on melon seed, are killed by treating the seed with Spergon Seed Protectant. (See pages 52 and 62.) If for disease and insect control see page 56.

For pollination of watermelons, see second paragraph under cucumber culture—page 16.

All our melon seed is now treated with Spergon to control seed-borne diseases and to give better stands, especially if planted early in cold ground.

The number of days after each variety indicates the time from seed planting to ripe fruit.

*All prices quoted are postpaid.
Write for special prices on larger quantities.*

FAIRFAX. (92 days.) This new variety was developed by the U.S.D.A. Vegetable Breeding Laboratory at Charleston, S. C. It has the general appearance, shape and color of Garrison but is distinguishable by a darker green stripe on the light green base. Also it has a hard rind in contrast to the soft rind of Garrison, thus making it a good shipping melon, but the flesh texture and edible quality (sugar content) do not compare with Garrison. It produces a melon up to 50 pounds, but will have an average weight of 30-35 pounds. Relatively resistant to hollow-heart and develops less "gourdneck" and blossom end rot than most long melons. Fairfax is resistant to both Fusarium Wilt and anthracnose. The vines are vigorous and hardy and show considerable capacity to withstand adverse weather.

**Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 55c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.**

GARRISON. (95 days.) We offer the genuine strain of this prized variety. Fruits are uniformly cylindrical in shape, very large, long with a thin, tender rind. The color is white and green striped, the green stripes usually being broken. Delicate flavor, sweet and tender meat, and grows to weigh 40 to 50 pounds, if the vines are pruned to one or two good shaped melons per plant, pulling off all other melons. Because of its high edible quality and large size it is popular and commands premium prices. It is not a shipping melon because of the brittle, easily broken rind, but is extensively for local markets.

**Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 55c; 1 lb. \$1.50
5 to 25 lbs. \$1.40 per lb.**



The Congo (Anthracnose-Resistant) Watermelon

CONGO (Anthracnose Resistant). (90 days.) This outstanding disease-resistant variety, resistant to both Fusarium Wilt and anthracnose diseases. The vines remain green longer than other varieties to protect the melons from sun burning. Congo has an exceedingly hard, tough rind which resists breakage and bruising in transit. In addition to these good qualities, it possesses many other desirable features. It is a heavy yielder of large size melons, under good growing conditions ranging from 32 to 40 pounds. Similar in size and shape to Garrison but having a darker green rind with darker stripes. It is free of the tendency to produce gourd necks. Flesh of attractive bright red color, of excellent quality and flavor, ripening well out to the rind, with small white seeds having light tan smears, and very attractive interior appearance. This variety does not need as heavy fertilization as other varieties, and little if any, side dressing with soda or nitrogen fertilizer should be used because heavy nitrogen may cause hollow heart or separation of the flesh. Melons of this variety should be harvested just before full maturity to avoid hollow hearts.

CONGO—FLORIDA STATE CERTIFIED—(Anthracnose Resistant). (90 days.) **CERTIFIED SEED SOLD ONLY IN 1 LB. SEALED BAGS** with the official Florida State Department of Agriculture certification tag attached to each sealed bag.

1 lb. \$1.75; 5 to 25 lbs. \$1.65 per lb.

***CONGO—VERIFIED (90 days.).**

**Pkt. 10c; oz. 20c; $\frac{1}{4}$ lb. 55c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.**

***VERIFIED WATERMELON SEED**

Associated Seed Growers, Inc. verifies that these seeds were: (1) grown in California, in an area free from anthracnose; (2) grown from seed stock tested and approved by the Asgrow Breeding and Research Department; and (3) are the product of a crop that was inspected by trained Asgrow fieldmen and found to meet the standards for freedom from disease and for trueness to variety type established by the Asgrow Quality Control Department.

**ASSOCIATED SEED GROWERS, Inc.
New Haven, Connecticut**

**OUR MELON SEED IS PRODUCED FROM SPECIALLY SELECTED STOCK SEED
WE HAVE SPECIALIZED IN WATERMELON SEED FOR MANY YEARS**

Florida Giant (Black Diamond or Cannon Ball)

FLORIDA GIANT (BLACK DIAMOND or CANNON BALL). (92 days.) The vines are of vigorous growth and produce large, nearly round, solid dark green-skinned melons, with thick rind and firm, rather coarse grained and fibrous, sweet, deep red, attractive flesh with small black seeds. Very hard, and withstands poor growing conditions better than most other varieties. Because of the dark green rind, it is in great demand, always having a fresh appearance. Has grown in favor among commercial melon growers in Florida. Also a desirable home garden variety.

**Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.30
5 to 25 lbs. \$1.20 per lb.**

BLACK DIAMOND—OKLAHOMA CERTIFIED. (92 days.) Our certified Black Diamond or Cannon Ball is produced for us by one of the best watermelon seed growers in Oklahoma and the fields are thoroughly inspected and certified by the Oklahoma Crop Improvement Association. No finer strain of seed of this popular variety can be secured anywhere at any price. Not a melon is shipped from our Oklahoma seed growing fields, and seed is saved from only the earliest maturing, largest sized and best shaped melons. **This certified seed is sold only in 1-lb. sealed bags** with the official Oklahoma Crop Improvement Association Certification tag attached to each sealed bag.

1 lb. \$2.00; 5 to 25 lbs. \$1.90 per lb.



Black Diamond

SUPER BLACK DIAMOND (Yellow Belly Strain). (92 days.) Our seed of this strain was harvested from only the largest, most uniform melons in the field. Melons are round in shape, like the regular Black Diamond, but of a darker blue-green color, bottom of the melon being a deep yellow. Seeds blackish brown little darker than the regular Black Diamond, and smaller. Rind somewhat thinner, and a little tougher. Meat very red, larger heart, and better quality. Anthracnose does not bother this type as much as the regular Black Diamond melon. The Super Black Diamond is a heavy bearer. Florida melon growers will be highly pleased with this new strain because of its attractive dark green rind and superior edible quality.

**Pkt. 10c; oz. 20c; ¼ lb. 55c; 1 lb. \$1.75
5 to 25 lbs. \$1.65 per lb.**

SUGAR BABY—A small icebox type watermelon weighing from 6 to 10 lbs. Early and prolific, perfectly round, with a thin hard tough rind, which is desirable for shipping. The melons are distinctly striped when immature, but become almost black when ripe. Inside appearance and flavor good. This melon is considered superior to other types of the icebox melons presently grown.

Seed supply limited.

**Pkt. 10c; oz. 30c; ¼ lb. 90c; 1 lb. \$3.00
5 to 25 lbs. \$2.90 per lb.**



CHARLESTON GRAY (Wilt and Anthracnose Resistant). (New). (92 days.) This is one of the most promising of the new modern varieties recently introduced. Developed by the U.S.D.A., Southeastern Breeding Station at Charleston, S. C. it is highly resistant to both Fusarium Wilt and Anthracnose. It makes a long, grey melon very slow to "sunburn", with a tough but not particularly thick rind, very striking in appearance when cut, with red meat and black seed. It is of delicious flavor and quality. A good yielder and very uniform in size, averaging under good growing conditions about 28 to 35 lbs. Will stay in the field in good condition for several days after it is ripe. This, along with its high resistance to the two most serious diseases of watermelons, makes it especially valuable for home garden use and to hold for a strong market. This new variety offers much promise as a high quality, disease-resistant, good shipping type melon for Florida.

CHARLESTON GRAY—*VERIFIED (92 days.)

**Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00
5 to 25 lbs. \$1.90 per lb.**

CHARLESTON GRAY—FLORIDA STATE CERTIFIED—(Wilt and Anthracnose Resistant). (92 days.) **THIS CERTIFIED SEED IS SOLD ONLY IN 1 LB. SEALED BAGS** with the official Florida State Department of Agriculture certification tag attached to each sealed bag.

1 lb. \$2.00; 5 to 25 lbs. \$1.90 per lb.

NEW HAMPSHIRE MIDGET (ICEBOX TYPE) (65 days.) This small icebox type melon, weighing 3 to



New Hampshire Midget

5 lbs. is attracting attention in Florida for home use, roadside stands, and local markets. It is the earliest of all watermelons making ripe fruits in from sixty to seventy days from seed. The vines lack vigor, but are prolific, setting many small melons, round to slightly elongated in shape with a light grey mottled rind, which is very thin and tender. Flesh is a beautiful deep red with small black seeds.

NEW HAMPSHIRE MIDGET—*VERIFIED (65 days.)

**Pkt. 10c; oz. 20c; ¼ lb. 55c; 1 lb. \$1.75
5 lbs. \$1.65 per lb.**

**Certified New Hampshire Midget sold in 1 lb bags only.
1 lb. \$1.75; 5 to 25 lbs. \$1.65 per lb**

FIELD AND ROASTING EAR CORN

(For Sweet Corn, See Page 18.)

Plant 7 pounds (½ peck) per acre or 56 pounds (1 bushel) to 8 acres.

To increase yields with field corn in Florida two things are essential, namely closer spacing and ample fertilizer. On good land 20 to 26 inch spacing in 38-inch rows is best with 500 lbs. per acre of fertilizer at planting time followed by side dressings of nitrate of soda up to 300 lbs. per acre about 5 weeks after planting.

The number of days indicated after each item represents the time from seed planting to harvest under average spring conditions in Florida.

Write for special prices on larger quantities.

All prices quoted are subject to change without notice.

FUNK'S G-737A HYBRID. (95 days.) This new yellow corn has given excellent results in numerous commercial trials in the Everglades muck and offers fine possibilities for other sections of the state. It consistently makes high yields (two ears per stalk) of attractive, medium large ears, well covered to the tips with hard, semi-flint yellow kernels, with extra heavy shuck, very weevil resistant and of excellent keeping quality. Makes strong, sturdy stalks which stand drouth better than many others.

¼ lb. 25c; 1 lb. 60c; 2 lbs. \$1.00; postpaid. Not prepaid. 14 lbs. (pk.) \$3.25; 56 lbs. (bu.) \$11.90



Typical ears of Funk's G-737A Hybrid showing heavy husks and excellent resistance to weather and weevil damage.

CUBAN YELLOW FLINT. (96 days.) This special strain is deep rooted, therefore can be given closer spacing than other varieties. If given close spacing it will yield as much as any prolific corn with two ears on 20% to 25% of the plants. Produces larger ears, longer grains, and heavier yields than other strains of this variety. Weevil resistant due to the long, tight, heavy and very tough husks. Seed is lemon to orange-yellow in color, with about 95% white cobs, 5% red cobs. Produces well in wet or dry seasons, and is recommended in all parts of Florida where extreme resistance to weevil damage is desired.

¼ lb. 15c; 1 lb. 45c; 2 lbs. 75c; postpaid
Not prepaid. 15 lbs. (pk.) \$1.65; 56 lbs. (bu.) \$6.00



Dixie 18 Hybrid Corn

DIXIE 18 HYBRID (Certified). (93 days.) This yellow hybrid field corn was developed at the Coastal Plain Experiment Station in south Georgia in cooperation with the U. S. Department of Agriculture. Dixie 18 is the best yellow hybrid corn for central and north Florida and south Georgia; a heavy yielder, averaging 2 ears per plant. Yields of 80 to 100 bushels per acre have been reported. The plants are relatively short, sturdy, and very vigorous, with a deep root system, resisting wind damage, lodging and drouth. The ears are uniform and borne low down on the plant. It makes a semi-hard corn and is very desirable for "hogging off". The grain is yellow and intermediate in texture being harder than Whatley. Has good resistance to weevil.

¼ lb. 15c; 1 lb. 45c; 2 lbs. 75c; postpaid
Not prepaid. 14 lbs. (pk.) \$2.25; 56 lbs. (bu.) \$8.50

IMPROVED GOLDEN DENT. (86 days.) A wonderful yielder, and does splendidly here in Florida. It is planted extensively for early use, but is too soft for a general crop. Has large, deep, broad grains, and is desirable for early feed.

¼ lb. 15c; 1 lb. 45c; 2 lbs. 75c; postpaid
Not prepaid. 14 lbs. (pk.) \$1.65; 56 lbs. (bu.) \$6.00

CORN-Roasting Ear

(For Sweet Corn, See Page 18)

Plant 7 pounds (½ peck) per acre. In the garden make two plantings at different times. Plant ¼ pound to 200 feet of row.

Write for special prices on larger quantities.

OKLAHOMA SILVERMINE. (82 days.) A heavy yielder of large ears. The cobs are covered with 14 to 18 rows, and run from 11 to 12 inches in length. Very resistant to worm damage. An excellent variety for shipping, with a heavy dark green husk that holds its color in shipment and on the market. This is the standard roasting ear variety grown in central Florida.

¼ lb. 15c; 1 lb. 50c; 2 lbs. 85c; postpaid
Not prepaid, 14 lbs. (pk.) \$2.00; 56 lbs. (bu.) \$7.00

SOUTHERN SNOWFLAKE. (80 days.) The ears are large, 10 to 12 inches long, well filled with 16 rows of large, broad kernels, and of very attractive appearance. The long tight husks protect the ears from worms.

¼ lb. 15c; 1 lb. 50c; 2 lbs. 85c; postpaid
Not prepaid, 14 lbs. (pk.) \$1.75; 56 lbs. (bu.) \$6.50

NORFOLK MARKET (Trucker's Favorite).

(76 days.)

¼ lb. 15c; 1 lb. 50c; 2 lbs. 85c; postpaid
(pk.) \$2.00; (bu.) \$7.00 f.o.b.



Beautiful lawns can be grown in Florida to enhance the appearance of your Florida home.

Lawn and Golf Course Grasses for Florida

All grass seed prices quoted are f.o.b. Plant City except lb. lots which are postpaid.

Write for prices on larger quantities.

Kilgore's Lawn Grasses are especially adapted to Florida conditions. Our grass seeds are recleaned, well-bagged, and of highest quality stocks.

All grass seed prices quoted are subject to change without notice. Bitter Blue-Stemmed St. Augustine and Centipede Grass are used only for lawns. The other grasses listed and described on this page are used for lawns, golf courses and for pastures. For information on "When to plant grass seeds in Florida" see page 5. For lawn and golf course fertilizers see page 67. For lawn grass pesticides and supplies see page 57 and 74.

BERMUDA GRASS

Unhulled—For lawns sow 2 to 4 pounds per 1000 square feet.

For pasture, sow 25 pounds per acre.

Hulled—For lawns sow 1½ to 2 pounds per 1000 square feet.

For pastures sow 15 to 25 pounds per acre.

Bermuda grass is generally considered best for maintaining a beautiful and perfect lawn. It is of fine texture, can be maintained green practically the year around, is a rapid grower and produces a lush, green carpet of turf though it requires careful attention in maintenance. Frost will turn it brown for a short period, but it soon regains its green color. It prefers loam or clay soil and will grow well on sand or muck. 20 to 30 pounds per 1000 square feet of a complete fertilizer such as Vigoro or Vertagreen should be applied twice a year. Frequent weeding is desirable. Although Bermuda grass will respond to watering, it will stand considerable drouth. A thorough watering every 5 to 7 days during periods of drouth should be sufficient. The grass may be clipped closely (¾ to 1 inch) and frequently. Bermuda will not grow in shade. It spreads by surface runners and by underground root stalks. Seed can be planted from March to November.

UNHULLED BERMUDA GRASS SEED

1 lb. 70c; 10 lbs. \$3.80; 25 lbs. \$9.00; 100 lbs. \$35.00

f.o.b. Plant City, except pound lots which are prepaid.

HULLED BERMUDA GRASS SEED

The almost watertight husks of the seed have been removed, facilitating quicker, surer results, the seed germinating more uniformly and in a much shorter time with proper moisture conditions.

1 lb. 85c; 10 lbs. \$5.10; 25 lbs. \$12.25; 100 lbs. \$48.00

f.o.b. Plant City, except pound lots which are prepaid.

PENSACOLA BAHIA GRASS FOR LAWNS

For Pasture See Page 41.

For Lawns sow 100 to 150 pounds per acre or 2 to 3 pounds per 1000 square feet. For pastures sow from 10 to 15 pounds per acre.

This variety of pasture grass is sometimes used for low maintenance lawns. It is of fine texture and will remain green most of the winter (longer than Centipede grass). Will tolerate dry or infertile soils, but makes an open, generally less attractive turf or lawn than regular lawn grasses. It is somewhat difficult to mow, especially when rapid growing seed shoots develop during the summer, but should not be mowed closer than 1½ to 2 inches. It requires fertilization in mid-spring and late summer, with a complete fertilizer such as Vigoro or Vitagreen. It is susceptible to damage from salt spray, but is relatively free of insect and disease troubles. Spring and early summer are the best periods for planting. Seeds are somewhat slow to germinate, sometimes requiring several weeks. The seedbed should be kept moist until germination occurs. For best distribution of seed, it should be mixed with dry sand. Sow seed at the rate of 2 to 3 pounds per 1000 square feet. "Common" Bahia should never be used for lawns.

1 lb. 70c; 10 lbs. \$3.80; 25 lbs. \$9.00; 100 lbs. \$35.00

(Lawn Grasses Continued on Page 40)

For further information on starting, fertilizing and care of Florida lawns write for a free copy of "Kilgore's Flower Guide for Florida." This valuable book also contains complete information on growing flowers in Florida. Also write the Florida Agricultural Experiment

Station, Mailing Department, Gainesville, Florida for a free copy of Bulletin 518 entitled "Lawns in Florida" and for Circular S-42 entitled "Insects and other Pests of Lawns and Turf."



COMMON RYE GRASS

For Lawns Sow 250 to 300 Pounds Per Acre.

Or 6 to 8 Pounds Per 1000 Square Feet.

For Temporary Winter Pasture Sow 50 Pounds Per Acre.

This is a valuable and desirable fall and winter grass for Florida, but it does not make a permanent lawn. It dies out in warm weather of late spring and early summer. For quick covering of lawns and golf courses in winter it is unsurpassed. Do not plant in Florida before October or later than March. It will make a vigorous, luxuriant growth throughout the fall, winter and early spring but must have plenty of water. A common practice in Florida is to seed Italian Rye grass on other brown lawns or in brown unsightly spots of old lawn in the fall and winter for a velvety green all-winter lawn. Plant $\frac{1}{2}$ lb. to 100 square feet right on top of old lawn after loosening the surface soil. Keep soil moist until seed sprouts.

1 lb. 45c; 10 lbs. \$1.25; 25 lbs. \$2.60; 100 lbs. \$9.25

CARPET GRASS

For Lawns Sow 100 to 150 Pounds Per Acre or 2 to 4 Pounds per 1000 Square Feet

For Pastures 25 Pounds Per Acre.

Carpet grass is particularly well adapted to low moist soils and does not require much fertilization, but needs plenty of moisture. This grass is not suited to drouthy conditions or to sweet soils with a high pH. It turns brown in the winter, is not at all salt tolerant nor will it stand much shade. Keep mowed to a height of 2 inches. Plant in the spring from March to July, or in fall from October to December.

1 lb. 65c; 10 lbs. \$3.30; 25 lbs. \$7.75; 100 lbs. \$30.00

BITTER BLUE-STEMMED ST. AUGUSTINE GRASS

Use one bushel of sprigs per 200 to 300 square feet.

This is one of the most popular lawn grasses for Florida. It is quick to establish, rather coarse in texture, but easy to mow, and remains green the year around. It should be cut at a height of 2 inches. Severe cutting back is quite harmful. It should be fertilized two or three times a year at the rate of 20 pounds per 1000 square feet with a complete fertilizer such as Vigoro or Vertagreen. St. Augustine is our best shade grass and will grow on alkaline as well as slightly acid soils. It is very salt tolerant and can be

grown along the beaches where a good soil base is provided. It can be grown by vegetative propagation only. Sprigs should be planted in well prepared soil, setting about 6 x 9 inches apart. For chinch bug control in St. Augustine lawns see page 57.

1 bu. \$2.50; 5 bu. \$2.40 per bu.; 10 bu. \$2.25 per bu.

Not prepaid. Available April to November. Shipped only by express direct from a Florida nursery.

CENTIPEDE GRASS

Use one bushel of sprigs per 300 to 400 square feet.

This is considered the best low maintenance grass for Florida. It is adapted to a wide range of soils including high sand hills, and will tolerate infertile acid soils, can get by on a minimum of fertilizer and water, and is relatively free of insect pest and diseases. It will not do well on alkaline or sweet soils but makes best growth on acid soils. A light spring and/or fall application of a complete fertilizer such as Vigoro or Vertagreen—10 to 15 pounds per 1000 square feet, should be sufficient for best results. **Do not over-fertilize.** The fertilizer should be watered in. In periods of extended drouth a thorough watering every 5 to 7 days should be sufficient, wetting the soil 4 to 6 inches deep each time. It is quick to establish and easy to mow to a height of 1½ inches. However, it turns brown during the winter, does not tolerate shade and salt spray as well as St. Augustine, and does not grow well in alkaline soils or in dense shade.

Centipede sprigs or root runners 3 to 5 inches long should be planted in rows 10 to 12 inches apart, 4 to 5 inches apart in the rows. Insert sprigs in the ground, leaving about an inch showing. Give frequent watering until plants are rooted firmly.

Centipede Grass Seed

1 oz. \$1.00; $\frac{1}{4}$ lb. \$3.25; $\frac{1}{2}$ lb. \$6.25; 1 lb. \$12.00

Postpaid

Centipede Grass Sprigs or Roots

1 bu. \$2.25; 5 bu. \$2.15 per bu.; 10 bu. \$2.00 per bu.

Not prepaid. Available April to November. Shipped only by express direct from a Florida nursery.

Write Mail Order Department, Kilgore Seed Co., Plant City, Fla., for instructive folder on "Centipede Grass Lawns."

PASTURE GRASSES and CLOVERS

All prices quoted are f.o.b. Plant City except 1 lb. and 2 lb. lots which are postpaid.
Write for prices on larger quantities.
All prices quoted are subject to change without notice.

To get better growth and increase soil nitrogen, it is essential to inoculate seed of all leguminous crops—clovers, Southern peas (Cowpeas), crotalaria, hairy Indigo, lupines, peanuts, sesbania, etc. (See page 69). Bermuda, Carpet and Italian Rye Grass are used for pasture as well as lawns. (See pages 39 and 40).



White Dutch Clover Pasture.

Photo courtesy of Florida Experiment Station

BAHIA GRASS FOR PASTURES

Sow 10 to 15 pounds broadcast per acre for pasture.

Bahia grass is established with seed planted $\frac{1}{4}$ to $\frac{1}{2}$ inch in depth, during any month of the year in Florida. Seeds often germinate slowly being covered by tough, waxy coat which does not allow water to enter for germination. Exposure to sun for a few days before planting hastens germination. Widely adapted. Will grow in dry soils because of deep root system. Makes one of Florida's best permanent pasture grasses.

Common Bahia. A low growing perennial, spreading by short, stout, woody runners which root heavily with large fibrous roots forming tough sod, even on drouthy, sandy soils. Leaf blades shorter and coarser than Pensacola but usually less than $\frac{1}{2}$ inch wide.

1 lb. 60c; 10 lbs. \$2.80; 25 lbs. \$6.50; 100 lbs. \$25.00

Pensacola Bahia. Narrow leaf, tall, erect-growing strain of Bahia grass, very hardy, claimed to stand cold weather conditions better than Common. Pensacola variety seed usually germinates quicker and stronger than Common. Succeeds on wide range of soils. Has heavy root system and withstands drouth better than other pasture grasses. For best results fertilize with 500 lbs. per acre of a complete fertilizer. Prevents erosion and provides very palatable pasturage. Withstands heavy grazing and is very resistant to trampling by livestock.

1 lb. 70c; 10 lbs. \$3.80; 25 lbs. \$9.00; 100 lbs. \$35.00

Argentine Bahia

Makes a semi-erect grass generally more vigorous than other Bahias, with leaves wider than Pensacola but slightly narrower than Common. Adapted to all kinds of soils, from high dry sand to low wet areas. Just as frost-resistant as Pensacola. For best results sow seed in late fall or early winter one-half to one-inch deep.

1 lb. 70c; 10 lbs. \$3.80; 25 lbs. \$9.00; 100 lbs. \$35.00

SWEET SUDAN GRASS

Sow 10 pounds in drills or 20 pounds broadcast per acre.

Resembles common Sudan in plant habit and yield, but is sweet and juicy, more productive and resistant to several foliage diseases which attack common Sudan. Grazing tests have indicated that cattle prefer Sweet Sudan to common Sudan. Sow from March to Sept.

1 lb. 40c; 10 lbs. \$1.30; 25 lbs. \$2.75; 100 lbs. \$10.00

BEGGARWEED (Florida Clover)

Sow 10 pounds per acre.

Perfectly adapted to Florida sandy soil. Makes fine pasture, nutritious hay. For best quality hay, cut crop when 3 to 4 feet high, or at beginning of blooming period. Sow from April to August. Cover seed lightly, inoculate seed with Nitragin. (See page 69.)

1 lb. \$1.15; 10 lbs. \$7.80; 25 lbs. \$19.00; 100 lbs. \$75.00

BENE (SESAME)

Sow 3 to 5 pounds per acre drilled.

Produces flowers, followed by seed pods, which scatter oily seeds in profusion, attracting quail and other game birds. Valuable for poultry feed. Plant from March to June, drilling seed thinly in rows 3 to 4 feet apart. We have only the native giant or tall variety growing to a height of from 4 to 6 feet, and rather spreading.

$\frac{1}{4}$ lb. 25c; 1 lb. 80c; 10 lbs. \$4.30; 25 lbs. \$10.25
100 lbs. \$40.00

ALFALFA-HAIRY PERUVIAN

Sow 20 pounds per acre.

A valuable forage and hay crop for Florida. Does well in central and south Florida where it is used extensively by dairy and cattle farmers. Does best on moist, dark sandy soils (not subject to flooding). Also does well on muck. Soil should be tested before starting seedbed and apply sufficient limestone several months in advance of seeding worked deep into soil to correct pH. The pH should be brought up to 6.5 or 6.8. Extra lime can be applied after the seedbed has been prepared if pH is too low. After plowing, firm seedbed by disking and cultivating. Broadcast 1000 pounds per acre of fertilizer high in phosphoric acid and potash plus 20 pounds of borax ahead of seeding. Inoculate seed with double recommended rate of inoculant. Use cultipacker type of seeder, and seed from September to January at rate of 20 pounds per acre, when soil is moist. After seeding apply 500 pounds of a phosphoric acid-potash fertilizer broadcast in 60 to 90 days.

1 lb. 70c; 10 lbs. \$4.10; 25 lbs. \$9.75; 100 lbs. \$38.00

ALYCE CLOVER

Sow 10 to 15 pounds per acre.

Used as cover crop, soil builder, for hay, and as pasture plant. Best time for planting is during spring. Must be planted on well drained soil free of contamination with root knot. Seed should be planted on well prepared ground and covered very lightly. Pack soil after seeding. Makes palatable hay. Cut for hay when it begins to flower. A good soil builder. Most economical fertilizer application is 200 to 400 pounds per acre of 0-16-16 or 0-14-14 mixture.

Write for prices.

CLOVERS and SOUTHERN PEAS (Cowpeas)

All prices quoted are subject to change without notice.

Prices quoted are f.o.b. Plant City except 2 lb. lots or less which are postpaid.

Write for prices on larger quantities, indicating variety and quantity desired.

Clovers (continued from page 41)

ANNUAL SWEET CLOVER

Sow 8 to 12 pounds per acre of inoculated seed alone, or 3 to 5 pounds per acre in mixed clover plantings.

A most valuable annual legume crop for grazing and soil improvement. Sow in Florida from Sept. through Jan., although from Oct. 15 to Nov. 15 sowing is best. Makes a rapid growth, and widely adapted, growing best on moist soils. Does not tolerate very close grazing. Grazing reaches peak during March and April. Liming soil heavily and applying 400 to 600 pounds of a fertilizer high in Phosphoric Acid and Potash at planting time is recommended.

HUBAM (Sweet). The best and most promising commercial variety for Florida.

1 lb. 50c; 10 lbs. \$1.95; 25 lbs. \$4.40; 100 lbs. \$16.50

FLORANNA (Sweet). This new variety developed by the Florida Agricultural Experiment Station from hardy strains found volunteering is best adapted to Florida soil and climatic conditions. It outyields and comes through drouth and cold weather better than other varieties.

1 lb. 60c; 10 lbs. \$2.50; 25 lbs. \$5.75; 100 lbs. \$22.00

WHITE DUTCH CLOVER (Louisiana grown)

Sow 4 to 6 pounds per acre alone, or 2 to 3 pounds per acre in a mixture.

Recommended principally for mixture in perennial pastures. Furnishes abundant grazing throughout the year in the South. Also useful for lawns, especially in mixtures. Does well with Bermuda Grass. Succeeds best on moist ground and during wet seasons. Sow in the late fall and early winter. Fertilize with 400 to 500 lbs. per acre of 0-14-10 mixture.

1 lb. \$1.50; 10 lbs. \$11.30; 25 lbs. \$27.75; 100 lbs. \$110.00

LADINO CLOVER (Certified)

This is a large form of White Dutch Clover. It grows rapidly and spreads by creeping stems that root at the nodes. Sow in late fall or early winter. Used for soil improvement, pasture and hay. Does best on fertile, moisture retaining soils.

1 lb. \$1.20; 10 lbs. \$9.00; 25 lbs. \$21.85; 100 lbs. \$86.50

CERTIFIED KENLAND RED CLOVER

Sow 4 to 6 pounds per acre.

This clover is rapidly gaining in popularity in Florida as a luxuriant hay and pasture crop for cattle and hogs. It has a longer active growing season than some other legumes. Fertilize with 400 to 600 lbs. per acre of a 0-14-10 or 0-10-10 fertilizer mixture just before planting time. Seed from September to December. Grows well with Bermuda and Bahia grass. For hay cut when in full bloom. It makes good growth in April, May and June when White Dutch and Crimson clover don't do so well.

1 lb. \$1.10; 10 lbs. \$7.30; 25 lbs. \$17.75; 100 lbs. \$70.00

CULTURE OF SOUTHERN PEAS

Peas may be sown any time from March to September. Table varieties are sown in drills three to four feet apart. The amount of seed per acre depends upon the size of seed, which varies with varieties. Fertilize up to 300 lbs. per acre. Inoculate seed with Nitragin. (See page 69.) It is important to treat pea seed with Spergon at the rate of 2 oz. per bushel when planted before the ground gets warm in spring. (See pages 52 and 62.) For insect control see page 54.

Sow 20 to 45 pounds per acre in drills with table varieties.

In the garden $\frac{1}{2}$ to $\frac{3}{4}$ pounds to 150 foot row.

Sow 1 to 1 $\frac{1}{2}$ bushels per acre broadcast with field varieties.

A bushel weighs 60 pounds.

TABLE VARIETIES OF SOUTHERN PEAS

TEXAS CREAM 40 (Improved Bush Conch.) (62 days.)

This variety of Conch pea is very early. A bunch type, the stocky vigorously upright growing plants making a bush about 24 inches tall. Some short runners of tendrils may develop as with most table peas. Pods



Thin Shelled Purple Hull (Dixie Queen)

are oval, medium green, straight, and closely filled with light green, tender peas. Seeds white.

$\frac{1}{4}$ lb. 20c; 1 lb. 60c; 2 lbs. \$1.10

15 lbs. (pk.) \$3.65; 60 lbs. (bu.) \$14.00

CALIFORNIA BLACKEYE NO. 5 (Early Wilt-Resistant Ramshorn Blackeyes). (65 days.) It is resistant to nematodes (root knot) and cowpea wilt. Very early and prolific, making a great number of 10 to 12 inch long pods on an upright vine. Uniform in size, growth and maturity, and flavor.

$\frac{1}{4}$ lb. 20c; 1 lb. 55c; 2 lbs. \$1.00

Write for prices on larger quantities.

GIANT WILT-RESISTANT RAMSHORN BLACK-EYES. (70 days.) A vigorous heavy yielding pea. About 10 days later than the Early Wilt-Resistant Ramshorn Blackeye. The peas are much larger than the regular Blackeyes and can be used both green and dried.

$\frac{1}{4}$ lb. 20c; 1 lb. 55c; 2 lbs. \$1.00

Write for prices on larger quantities.

ALABAMA CROWDER. (70 days.) The very vigorous bush is of the semi-runner type. The large, straight pods average about 8 to 9 inches in length and are of very attractive appearance, which has created a demand on the market for this variety, selling for higher prices than other Crowder peas. A sure cropper, heavy yielder, and well adapted to Florida growing conditions.

$\frac{1}{4}$ lb. 20c; 1 lb. 55c; 2 lbs. \$1.00

15 lbs. (pk.) \$3.75; 60 lbs. (bu.) \$14.00

THIN SHELLLED PURPLE HULL (Dixie Queen). (75 days.) Vines make vigorous branching growth, and do best when planted in three-foot rows. Very productive. Pods ripen to a deep purple color, are very large, long, slightly curved, well filled with large seeds of high edible quality.

$\frac{1}{4}$ lb. 20c; 1 lb. 60c; 2 lbs. \$1.00; pk. \$3.75; bu. \$14.00

FIELD VARIETIES OF COWPEAS

Due to the unstable condition of the market, we are unable to make price quotations on field varieties of cowpeas.

BRABHAM. Is resistant to wilt and root knot, makes fine, vigorous growth of vines, is early, and when planted in March, April, or May good cuttings of hay can be made. Also a very good soil builder.

1 lb. 45c; 2 lbs. 80c; pk. \$1.50; bu. \$5.50

MIXED. The advantage claimed for sowing Mixed Cowpeas is that they produce an exceedingly heavy growth of vines for forage or soil improvement.

1 lb. 40c; 2 lbs. 75c; pk. \$1.25; bu. \$4.50

FIELD SEED

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.

For information on "When to Plant Field Seed in Florida" see page 5.

CHUFAS

Plant 1 to 1½ pecks (10 to 15 lbs.) per acre.

(90 days.) This is a species of ground nut, easily grown, a useful hog fattening crop. Plant from April to August. Cultivation same as for bunch peanuts. Can be left in the ground until time to turn in hogs who do harvesting. On sandy loam soil best adapted for Chufas, plant in 2-ft. rows with plants 12 inches in the row, or in 3-ft. rows with plants 6 inches in the row.

1 lb. 50c; 2 lbs. 90c; 10 lbs. (pk.) \$2.25
40 lbs. (bu.) \$8.00

CHICKEN CORN. (Shallu or Egyptian Wheat)

Sow 8 to 10 pounds per acre for grain.
25 to 30 pounds for cover crop.

This crop is being grown extensively and very successfully as a summer green manure or cover crop in many sections of Florida. Grows rapidly and makes a great amount of vegetable matter to plow under in a relatively short time. Not a legume. If allowed to go to seed for a grain crop like other sorghums, makes large, loose, bushy heads covered thickly with small grains, which if left standing, shatter, and chickens gather them. Grain is of high feeding value and can be fed to chickens, hogs and cattle as desired. For grain sow seed thinly in 3 to 4 foot rows, leaving 3 to 4 plants every 3 feet of row. Plant from March to August.

1 lb. 50c; 10 lbs. \$1.55; 25 lbs. \$3.40; 100 lbs. \$12.50

CROTALARIA

Sow 10 to 20 pounds of seed per acre.

Crotalaria is particularly desirable as a cover crop for groves, truck and farm lands, does well on all soils except muck. Is a legume, thus adding a large amount of nitrogen to soil. Is especially valuable because it is immune to nematodes causing root knot disease. Furnishing abundance of green material for humus.

Sow broadcast March to June, on well prepared land.

Crotalaria—Early Spectabilis. This vigorous growing, long leaved variety is not so fibrous as others, the growth is much heavier, and the stems are hollow. As a result, it rots much quicker when plowed under. Spectabilis does well on high ground, but is best adapted to the better grades of sandy soils. Leaves, stems and seed of this Crotalaria plant, green or dry, are poisonous to poultry and livestock.

Write for prices.



Crotalaria—Early Spectabilis
Valuable for adding organic matter and nitrogen to the soil.

HEGARI (Higear)—See Sorghum, page 46

INDIGO (Indigofera hirsuta)

Sow three to five pounds per acre in 30 inch drills,
6 to 10 pounds broadcast.

A three purpose legume, used as a summer cover crop, for hay and for grazing. Used extensively as a cover crop by citrus growers in south Florida, as a grazing crop or for hay by cattlemen and as a cover crop by general farmers. Its feeding value compares fairly well with alfalfa, making a palatable and nutritious hay. It makes an ideal summer cover crop for citrus groves, harboring no troublesome insects, and reseeds itself year after year. It's very resistant to drouth. The plants grow four to seven feet in height, and produce an abundance of hay and grazing. If grazed or harvested for hay before the stems become woody a second or third growth may be expected.

Seed may be planted not over an inch deep, from the middle of March to the later part of May, but early seedlings are preferable. Heavier seedings are desired for green manure or forage, lighter seeding for seed. On poor soil 300 to 500 lbs. per acre before seeding of a 0-10-10 or 0-14-10 fertilizer mixture is desirable. On extremely acid soils apply 1000 lbs. of lime per acre. It will grow on a wide range of soils, but well drained sandy loam soils are best. It is sensitive to cold, is killed by first hard frost and is best suited to the section of Florida from Gainesville south, although the early strain can be grown successfully in north Florida and as far north as central Georgia. Highly resistant to root knot, but in general it is reasonably free from disease and insect injury.

COMMON INDIGO Write for price

EARLY INDIGO Write for price

BLANKET INDIGO

Sow 3 to 5 pounds of seed per acre.

This is a summer annual legume, different from Hairy Indigo in growth—ordinarily 18 to 24 inches. Excellent for sowing with other permanent pasture grasses. Also recommended for sowing in groves. Plant from early spring to the end of June. Offers great possibilities as a pasture or hay crop in Florida—especially for hay in late summer—September or early October. Prefers soils of low fertility, especially low in nitrogen. A good soil builder. Will not stand excessive moisture. It is highly palatable and will stand close grazing.

Write for prices.

LUPINES

50 to 60 pounds with yellow and 50 to 90 pounds with blue per acre broadcast, or preferably in drills with a grain drill.

Lupines are adapted to north and central Florida as a winter legume cover crop. They are not affected by cold and because of their heavy yield make an ideal green manure crop. Plant from late September to December, preferably in late September or October, in well prepared soil, using between 250 and 300 lbs. of Superphosphate per acre.

Bitter Blue Lupine. This variety has rapidly grown in popularity as a soil builder. Normal growth is about waist high and normal weight of green matter 20-30 tons per acre. It produces about three times as much growth as Winter Peas, Hairy Vetch, etc. It protects the soil from erosion and restores humus and plant food removed by peanuts and other crops. Peanut and corn yields following Bitter Blue Lupine have been greatly increased. A good winter cover or green manure crop, but poisonous to livestock.

1 lb. 40c; 10 lbs. 90c; 25 lbs. \$1.75; 100 lbs. \$6.00

(Continued on next page)

FIELD SEED (Continued)

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.

LUPINES (Continued)

SWEET BLUE LUPINE. This variety is the pasture and feed counterpart of Bitter Blue Lupine. Both plant and leaf are sweet to the taste and are relished by all kinds of livestock. Makes about the same growth and tonnage as Bitter Blue Lupine. Stock can be turned in on Sweet Blue Lupine as soon as it is about a foot high. The plants have a high protein value averaging approximately 18% of the dry matter.

1 lb. 50c; 10 lbs. \$1.45; 25 lbs. \$3.10; 100 lbs. \$11.50

SWEET YELLOW LUPINE. (White seeded.) Seems to be more palatable and therefore is more readily grazed by cattle than the Sweet Blue strains, and will recover more rapidly than the Blue after grazing. The sweet strains of Yellow Lupine tend to be more tender and less resistant to excessive cold than the Blue, and thus the Yellow Lupine is more particularly adapted to the southern portion of the Lupine belt than to the more northern areas.

1 lb. 60c; 10 lbs. \$1.90; 25 lbs. \$4.25; 100 lbs. \$16.00

PEARL or CATTAIL MILLET

Sow 8 to 10 pounds per acre in rows 3 feet apart.
Sow 20 to 25 pounds per acre broadcast for pasturage.

This is one of the best yielders of green forage and hay, and continues to grow and produce through the entire season if cut frequently enough for hay to prevent its going to seed. Relished by all kinds of stock, and they eat it greedily. Plant in early spring as soon as frost danger is past in March, and can be planted from then on until September 1. Sow thinly in rows three feet apart or for pasturage, sow broadcast 20 to 25 lbs. per acre.

1 lb. 45c; 10 lbs. \$1.50; 25 lbs. \$3.25; 100 lbs. \$12.00

OATS

Sow 5 pecks (40 lbs.) per acre in drills; broadcast 2 to 2½ bu. per acre. A bushel weighs 32 pounds.

Plant in drills 1½ feet apart and cover seed 2 to 3 inches deep. If the seed is broadcast, use tooth harrow. The best time to plant oats in Florida is October and November, but can be planted to January. Land should receive 300 to 400 lbs. per acre of complete fertilizer at planting, followed by a side-dressing of 150 to 200 lbs. Nitrate of Soda or its equivalent after the crop is half grown.

VICTORGRAIN 48-93. (120 days.) An early high yielding cold resistant variety, with strong, stiff, straws that will withstand wind and storms. Resistant to most races of smut and rust, also tolerant to strains of blight. Victorgrain 48-93 is an improvement over the original strain and is highly recommended for Florida.

1 lb. 35c; 2 lbs. 50c; pk. 60c; bu. \$1.65



Victorgrain 48-93

SOUTHLAND (Disease Resistant). (125 days.) Is a vigorous growing oat with broad, coarse leaves and a strong upright growth habit. Produces a large amount of very early forage. A spring type oat and is not cold-hardy, and where not injured by cold it produces an abundance of green pasture. Adapted to the Gulf Coast region and to north central Florida. Resistant to Victoria blight and to most races of Crown rust, it is susceptible to two races of this rust which have now become prevalent in Florida. It suffers little from stem rust, but is quite susceptible to Culm rot. Yields of forage and grain have been high under good cultural conditions. Grain is heavy and plump, free of hairs, white in color and of high quality. A good general purpose oat for Florida.

1 lb. 35c; 2 lbs. 50c; 8 lbs. (pk.) 60c; 32 lbs. (bu.) \$1.75



Southland Oats

SEMINOLE (Disease resistant). (110 days.) This new, early maturing oat has been increased on Experiment Station Farms in Florida, and seed is now available. It is resistant to different races of Crown rust and to smut. Due to its earliness it escapes damage from stem rust. A short variety with strong straw and not subject to lodging. The grayish-white grain is larger, plumper and of higher quality than Floriland. Makes a very early, upright growth of medium large leaves. Produces an abundance of early forage and is good for early and mid-winter forage. Also excellent in grain yield. Because of its earliness it is well adapted to central and south Florida.

Lb. 35c; 2 lbs. 50c; 8 lbs. (pk.) 60c; 32 lbs. (bu.) \$1.75

PEANUTS

Plant 1 bushel of seed per acre.

A bushel weighs 25 lbs., except Valencia 22 lbs. per bu.

In addition to the crop of nuts for hogs, peanuts yield a large quantity of nutritious dry forage. Dixie Runner peanuts are by far the best when grown to fatten hogs, as they can be left in the ground until eaten.

It is better to shell the seed that is to be planted. If planted in corn, a bushel will plant 4 acres, but where planted solid it requires a bushel of seed per acre, in 3-foot rows, dropping seeds in hills a foot apart. Plant from March to July. For best results inoculate peanuts with Nitragin. (See page 69.)

Most shelled peanuts are treated so they will keep better and longer, and also to protect them from rot and disease after they are planted and to increase the stand. Do not use Nitragin inoculation with treated shelled peanuts.

For Insect and Disease Control See Page 57.

VALENCIA. (100 days.) This variety is adapted for growing in light sandy soils. It fills the demand for a peanut that is large and contains more nuts than the common varieties. Pods are large, well filled, usually containing 3 or 4 nuts.

Write for prices.

(Continued on next page)

FIELD SEED (Continued)

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.



Field of Dixie Runner Peanuts

DIXIE RUNNER. (120 days.) This variety produces high yields like Spanish peanuts, in four years' tests averaging 53 per cent higher yields than the Florida Runner, and yet remains dormant after maturity like the regular Runners, thus being especially suited to "hogging off." It is a vigorous grower with more prostrate branches, and therefore is more apt to fill the spaces left by skips in the stand. Kernels are much like Florida Runner in size, shape, markings and color. Pods are somewhat lighter and brighter in color, shorter, thicker at the waist and more cylindrical in shape. In addition to high yield and dormancy characters, this variety possesses a valuable quality for shellers, as it shows less than 1 per cent of concealed damage, or decay of the interior of the kernels, which is a serious disadvantage of some varieties going to the shelling plants, and may average as high as 20 per cent concealed damage.

Write for prices.

Shelled and treated Dixie Runners. One bushel of good Dixie Runners should yield about 15 lbs. shelled peanuts. Plant 25 lbs. per acre.

Write for prices.

DWARF ESSEX RAPE

Sow 4 pounds in drills or 8 to 10 pounds broadcast per acre.

For early spring, fall or winter planting to give quick green forage or grazing for hogs and poultry. While mostly planted in fall and early winter, it may be planted in early spring, but it will not stand much warm weather. It makes a quick succulent growth. It can be planted in rows like turnips, or broadcast covering by a light harrowing. It is ready for grazing in eight to ten weeks after planting. After cutting, or grazing, it keeps coming out again. Rape can be sown with fine results any month from September to March. It is a splendid green feed and forage crop relished by all kinds of stock and especially by milk cows.

Pkt. 10c; 1 lb. 55c; 10 lbs. \$2.30; 25 lbs. \$5.25
100 lbs. \$20.00

RICE

Sow $\frac{1}{2}$ bushel (22 lbs.) per acre.

Rice makes fine feed for both stock and poultry. Prepare the ground thoroughly, drill $\frac{1}{2}$ bushel per acre in 2½-foot rows and cultivate a few times. Should be planted in March or April so that it may mature at a time to miss the rice-bird season. If planted in June and July it will also miss the bird season.

BLUE BONNET (Gopher). A popular variety for Florida, generally known as Upland Rice, but will also do well on low land. Grains heavy, broad, and flat.

1 lb. 35c; 2 lbs. 60c; pk. \$1.60; bu. \$6.00

ZENITH (Pearl). Grown extensively in Florida. It produces a heavier, taller, more vigorous stalk with heavier heads than other varieties of this type. Does not shatter so badly as other varieties. Produces long, slender grains. Best adapted for low wet land.

1 lb. 35c; 2 lbs. 60c; pk. \$1.60; bu. \$6.00

RYE

Sow 3 pecks in drills; broadcast 1 to 1¼ bushels per acre. A bushel weighs 56 pounds.

Rye can be planted from September to January but October, November and December are the best months in which to plant for seed. Earlier planting is recommended for winter and early spring grazing. Fertilization is the same as for oats. (See page 44.)

FLORIDA BLACK. A standard, upright, tall growing variety. Northern and western varieties of rye have a tendency to fall over on the ground and are not suitable for grazing or cutting in Florida. Florida Black is upright in growth and is exceptionally well suited for early grazing, making pasturage quickly, and it withstands cold very well. Highly recommended by the Florida Agricultural Experiment Station.

1 lb. 50c; 2 lbs. 85c; 14 lbs. (pk.) \$1.40
56 lbs. (bu.) \$5.00

ABRUZZI. One of the best for Florida, producing a heavier tonnage of green feed and grain than most other varieties. Tall and upright, and a heavy yielder.

1 lb. 40c; 2 lbs. 75c; 14 lbs. (pk.) \$1.15
56 lbs. (bu.) \$4.00



Abruzzi Rye

FIELD SEED (Cont'd)

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.



Sorghum—Texas Seeded Ribbon Cane

SORGHUM

Amount of seed needed per acre depends upon the purpose for which this crop is grown as indicated below.

Sow 8-10 lbs. to 1 bushel (50 lbs.) per acre.

The uses of Sorghum are many; green feed, hay, pasturage, silage, syrup, etc.

It is common practice to sow Sorghum and Cowpeas together for hay and pasturage purposes using about ½ bushel Sorghum and 1 bushel of Cowpeas per acre, sown broadcast. Planted in drills alone in 3- to 3½-foot rows use 1 peck or 12 to 15 lbs. per acre; where planted for making syrup, in 4-foot rows, use 8 to 10 lbs. per acre, thinning plants 8 to 10 inches apart; or where sown broadcast for forage, use 1 bushel of seed per acre. For grain Sorghums apply 300 to 500 lbs. of a complete fertilizer mixture per acre. Plant from March to August.

TEXAS SEEDDED RIBBON CANE. (For syrup, ensilage, forage and green feed.) The plants grow ten to twelve feet tall, and the thick stalks contain an abundance of juice and also produce plenty of leaves for fodder. Not only valuable for syrup-making, but will make a large amount of forage and green feed for livestock. The tallest and heaviest producing Sorghum for both forage and syrup.

1 lb. 55c; 10 lbs. \$2.10; 25 lbs. \$4.75; 100 lbs. \$18.00

HEGARI (Higear). Early Dwarf Type. (For grain and forage.) (100 days.) This crop, which is quite similar to Kaffir corn, has become very popular in the south-east because of its quick growth and heavy yield of grain, sometimes running over a ton to the acre. Unlike Kaffir, the stalks are very sweet and juicy. Grows 2 to 5 feet depending on type of soil, with large heads of white grains which do not shatter. Stands dry weather exceptionally well. Sow ten pounds per acre for grain, one bushel (50 lbs.) per acre broadcast for forage.

1 lb. 40c; 10 lbs. 90c; 100 lbs. \$6.50

HEGARI (Higear). Regular Type. (For forage and cover crop.) (115 days.) This strain is slower to seed but makes more forage quicker than the Early Dwarf Type, and also is better for cover crop. Sow 30 to 40 lbs. per acre broadcast for forage or for a cover crop. On the lower east coast this strain is used extensively as a cover crop and a common practice is to cut the stand just before seed heads form; the plants will then sprout out and make similar height as before, thus giving a double cover crop to turn under.

1 lb. 40c; 10 lbs. 90c; 100 lbs. \$6.50

SHALLU OR EGYPTIAN WHEAT. See "Chicken Corn"—page 43.

SESBANIA

Sow 30 to 40 pounds per acre.

Sesbania is a most valuable summer cover crop to grow on land that is more or less covered with water. Unlike other cover crops, Sesbania will grow in water, and will grow ideally well in the hot weather of mid-summer on low, wet land. It is used extensively as a summer cover crop in low, wet muck lands, as well as in other vegetable and commercial flower-growing lands. It is valuable not only as a green manure cover crop, but also as a means of killing out Bermuda grass, nut grass and other weeds, because it grows so rapidly. In addition, being a leguminous crop, forming nodules on the roots, it fixes atmospheric nitrogen in the soil for seed inoculant. (See page 69.)

We recommend scarified Sesbania seed for quicker and higher germination. Seed is broadcast at the rate of thirty to forty pounds per acre, even a little heavier if weeds are bad. Usually in late May, June and July, dense, tall growth is obtained in about two months, at which time, just before blooming, the crop may be plowed under and the stalks will rot or decay rapidly. In some sections Sesbania is grown for pole bean sticks. For this purpose it should be sown 12 to 15 lbs. per acre.

Write for prices.

SUNFLOWER

Sow 6 pounds per acre.

MEDIUM CALIFORNIA. The Medium California makes three to four times as much seed as ordinary varieties. It is used extensively as a windbreak for beans. Plant from March to September. Plant seed in drills three feet apart, and hills one foot apart in the drill, thinning out to one stalk in the hill when four to six inches high.

1 lb. 55c; 10 lbs. \$2.30; 25 lbs. \$5.25; 100 lbs. \$20.00

VELVET BEANS

Sow 2 pecks (30 lbs.) per acre.

Velvet Beans are good for groves and truck lands in building up the humus; good pasturage for cattle and hogs; and fine land renovators. They are especially valuable because they are resistant or practically immune to nematodes causing root knot disease. Plant from March to August. Use Nitragin. (Page 69.)

90-DAY. (90 to 100 days.) This variety makes a very luxuriant growth and an abundant amount of foliage for pasturage and soil improvement. It matures quicker than other varieties.

1 lb. 45c; 2 lbs. 80c; pk. \$2.00; bu. \$7.50



A typical field of Hegari (Higear) grown for grain

STRAWBERRY PLANTS

All prices quoted are subject to change without notice.

We supply Maryland and Tennessee grown strawberry stock plants of the Missionary and Florida 90 variety from nurseries in those states. These northern grown stock plants are set out in multiplication beds in Florida from January through March. The young plants are usually set in the permanent field from September through November or early December, depending on weather conditions. The northern grown plants are usually available about November 1st and when Florida grown plants are not available are used by commercial growers for November setting and are also used to a great extent for home gardens. Care should be exercised never to allow the roots to dry out.

Strawberry plants may be set in rows three feet apart for single rows, or 4 to 5 feet apart for double rows, allowing 16 inches between double rows on each bed with 12 to 14 inches between plants in the row.

In setting strawberry plants, make holes deep enough to set the roots straight. Plants should be set so that the surface soil comes just to the crown.

Soil should be well firmed around the roots, and during dry weather each plant should be watered daily for several days, preferably in the late afternoon.

If the land is sour or acid, apply 500 lbs. lime or 1200 lbs. hardwood ashes per acre before setting plants. For control of insects and diseases see page 56.

See Fumigants on page 54 for Nematode (root knot) control.

After the plants are well established, 3 or 4 applications of high grade commercial fertilizer, such as 4-7-5 mixture, should be given the plants at about monthly intervals, being sure to make one heavy application of a 4-8-8 mixture when the plants begin to fruit.

Missionary Variety Virus-free	100 plants	250 plants	500 plants	1000 plants
	Postpaid		f.o.b.	Nursery
Maryland plants	\$2.25	\$4.00	\$5.00	\$8.50
Tennessee plants	2.00	4.00	5.50	8.50
Florida 90 Variety				
Maryland plants	2.25	4.00	5.00	8.50
Tennessee plants	2.00	4.00	5.50	8.50

ROSE BUSHES FOR FLORIDA

Planting and Care of Rose Bushes.

Plant bushes 1½ to 2 feet apart each way. Dig holes sufficiently deep and wide to permit roots to be set in natural position. Put two to three pounds of pulverized sheep manure in each hole and mix well with a half pound of Vigoro or Vertagreen fertilizer. (See page 67.) Then pour in a liberal supply of water, set bush, press the soil tightly around the roots, leaving no air spaces. Roses need lots of water, and should be fertilized every six to eight weeks with Vigoro or Vertagreen fertilizer, using about one-half pound per plant, and an occasional application of sheep manure will help. Spray with Nicotine Pyrox or dust thoroughly with Kilgore's Rose Dust (See page 63) at least once a week or as soon as diseases or insects appear.

All orders will be shipped direct from Texas nursery.

Delivery made in 12 to 15 days after receipt of order. Extra strong, No. 1 grade bushes—bloom approximately 2 months after setting out.

Postpaid to any point in Florida

5	10	25	50	100
\$5.00	\$9.50	\$23.00	\$45.00	\$85.00

No orders accepted for less than five bushes. However this does not apply to patented roses.

Plant any time after November 1 to the end of March.

VARIETIES OF BUSH ROSES

RED—Red Radiance, Poinsettia, Etoile De Hollande, Red Talisman.

PINK—Pink Radiance, Editor McFarland, Briarcliff.

WHITE—White American Beauty (Frau Karl Druschki), Caledonia.

YELLOW—Golden Dawn Soeur Therese.

TWO-TONE—Talisman—Orange and copper tones, suffused with pink.

VARIETIES OF CLIMBING ROSES

American Beauty (pink), Paul's Scarlet (scarlet).

Mrs. Pierre S. DuPont (rich golden yellow).

PATENTED AND NOVELTY ROSES

CHARLOTTE ARMSTRONG—Plant Pat. 455. Beautiful cerise pink ... \$2.25

MIRANDY—Plant Pat. 632. Beautiful bright red 2.25

PEACE—Plant Pat. 591. Golden yellow etched with pink 2.50

BRAVO—Plant Pat. 983. Red 2.25

BLANCHE MALLERIN—Plant Pat. 494. White 2.25

FORTY-NINER (HT)—Plant Pat. 792. Large double flower of cherry red 2.50

FLORIBUNDA ROSE

FASHION—Plant Pat. 789. Peach 2.00

See Kilgore's Flower, Vegetable Planting Guide for more complete information on Rose bushes.

The Kilgore Seed Company

General Offices and Mail Order Dept., Plant City, Fla.

FOURTEEN KILGORE SEED STORES
SERVING FLORIDA

A Florida Institution Maintained for Service to
Florida Growers

With over 45 years experience specializing in vegetable, flower and field crop seeds and growers' supplies for Florida, and with fourteen Kilgore seed stores we are best qualified to serve Florida growers.

"Everything for the Florida Grower" is a Kilgore motto. Seeds, insecticides, fungicides, sprays, dusts, spraying and dusting equipment, fertilizers, cans and canning equipment as well as poultry and beekeepers' supplies are always carried in stock in Kilgore's Plant City warehouses and fourteen Florida stores.

High quality with prompt and efficient service at reasonable prices are foundation stones on which this pioneer Florida seed company has been built.



We now have a group of technically trained field men thoroughly familiar with local conditions, who are available to discuss your insect, disease and seed problems with you at any time.

GLADIOLUS BULBS

Rainbow Mixed

Enjoy beautiful cut flowers from your garden. For continuous blooms plant bulbs every 10 days to two weeks. In north and central Florida plant from August to October, and January to May.

In south Florida plant from September to May. Set bulbs 4 inches deep and 6 inches apart in well-prepared and well-drained beds.

Disease-free, well cured, graded extra large bulbs of good color assortment. No finer Glads for Florida can be secured.

Large Exhibition Size Gladiolus Bulbs (2-2½ inches.)

1 dozen bulbs	\$.85 postpaid
25 bulbs	1.50 postpaid
100 bulbs	4.00 postpaid
1000 bulbs	35.00 postpaid





FLOWER SEEDS FOR FLORIDA

We recommend the following kinds and varieties of flowers for Florida planting. Quantities larger than packets are listed in Kilgore's "Flower and Vegetable Planting Guide". If you do not have a copy of this Guide we shall be glad to send one to you.



PRICES—All flower seed packets 15c each, postpaid. All flower seeds are subject to the Florida State Sales Tax. Rate of Sales Tax—11 to 35c—1c tax; 36 to 65c—2c tax; 66c to \$1.00—3c tax.

Ageratum or **Floss Flower**, Blue Perfection, Dwarf Blue Cap. Sept. - Jan.

African Daisy (Arctotis Grandis). Sept. - Jan.

Alyssum—Little Gem, Royal Carpet. Sept. - Jan.

Aster—Crego and Imperial Mixtures, Powderpuffs and Princess Mixtures. Sept. - April.

Baby's Breath—White. Sept. - Jan.

Bachelor's Button—Blue, Dainty Mixture. Sept. - Jan.

Balsam—Double Mixture. Sept. - March.

Beans, Scarlet Runner. Feb. - May.

Blanket Flower or **Gaillardia**—Double Mixture. Any Month.

Blue-Eyed African Daisy or **Arctotis**—Sept. - Jan.

Blue Lace Flower (Didiscus)—Sept. - Jan.

Calendula—Campfire (dark orange), Lemon Queen, Art Shades and Excellent Mixtures. Sept. - Jan.

California Poppy—Gorgeous Single Mixture—Sept. - Jan.

Calliopsis—Choice Mixture—Sept. - Feb.

Candytuft—White Hyacinth—Flowered, Border Group Mixed. Sept. - Feb.

Carnation—(Chabaud's) Charm Mixture. Sept. - Dec.

Celosia or **Cockscomb**—Tall Royal Velvet (Comb Type) Pampas Plume (Plume Type). Jan. - June.

Chinese Forget-Me-Not or **Cynoglossum**—Sky Blue. Sept. - March.

Cleome or **Spider Plant**—Pink Queen. Feb. - Aug.

Cosmos—Fiesta (Orange), Choice Mixture. Mar. - May.

Dahlia—Unwin's Bedding Mixture. Mar. - April.

Delphinium (Perennial Larkspur)—Belladonna Hybrids Mixed. Oct. - Nov.

Floss Flower or **Ageratum**. See Ageratum.

Globe Amaranth or **Gomphrena** (Florida Bachelor's Button)—Mixed Colors. Feb. - July.

Gourds (Ornamental)—Small varieties mixed. Feb. - May.

Hollyhock—Indian Spring (Single), Double Mixed—Rose and Pink Shades. Aug. - Jan.

Larkspur—Regal Finest Mixed. Nov. - Jan.

Linaria (Baby Snapdragon)—Northern Lights Mixed. Sept. - Dec.

Lupine—Tall Assorted Colors. Sept. - Dec.

Marigold—Extra Dwarf Double, Dwarf Double Harmony, Dwarf Gigantea Pot O'Gold, Lemondrop. Tall varieties—Cracker Jack, Crown O'Gold (Collarette), Gigantea Sunset Giants, Glitters (Chrysanthemum), Real Gold and Yellow Supreme (Carnation type). Feb. - May and again Sept.

Morning Glory (Ipomea)—Scarlett O'Hara (Red), Pearly Gates (White), Heavenly Blue, Improved (Blue). Mar. - June.

Mourning Bride or **Scabiosa**—Sunrise Mixture. Sept. - Dec.

Nasturtium—All varieties Semi-Double. Dwarf Gem Mixed, Golden Gleam, Gleam Hybrids, Mahogany Gleam, Scarlet Gleam. Sept. - March.

Pansy—Engelmann's Giant Mixture. Sept. - Dec.

Petunia—Carnival (F-2 Hybrid Mixture), Elks Pride, Flaming Velvet, Fire Chief, Fluffy Ruffles, Heavenly Blue, Howards Star Improved, Rosy Morn, Snow Queen, Superb Mixture. Sept. - June.

Phlox—Dwarf Nana Compacta and Peninsular Mixtures. Sept. - Feb.

Pinks or **Dianthus**—Medal or Imperator and Sweet Wivelsfield Double Mixtures; Single Gaiety Mixture. Sept. - March.

Queen Anne's Lace—White. Sept. - Jan.

Rose Bushes—See page 47.

Salvia—Bonfire. Sept. - June.

Shasta Daisy—Single and Semi-Double Giants. Sept. -

Snapdragon or **Antirrhinum**—Giant Mixture and Giant Ruffled Tetraploid Mixture, both Rust Resistant. Sept. - Dec.

Statice—Straight colors. Blue, Lavender, Rose, White, Yellow. Also Assorted Colors. Sept. - Jan.

Strawflower or **Helichrysum**—Many Colors. Sept. - March.

Sunflower or **Helianthus**—Single Mixed; Sungold (Double Yellow). Feb. - Aug.

Sun Plant or **Portulaca**—Double Mixed. Feb. - Aug.

Sweet Peas (Early Flowering Spencer)—Balls Orange Improved, Hope, Lavanda, Mars, Mrs. Herbert Hoover, Shirley Temple, Rainbow Mixture. Zvolanek's Multiflowered Mixture. Sept. - Dec.

Tithonia or **Florida Sunflower**—Torch. Feb. - June.

Verbena—Master Mixture. Any Month.

Vinca or **Periwinkle**—Pink and White Mixed. Any Month.

Zinnia—Blaze, Crimson Monarch, Polar Bear, Wonder Rose, Crown O'Gold, Fantasy, Giant Hybrids, Giants of California, Peppermint Stick, Persian Carpet, Pompon, Sunburst and Ortho Polka Mixtures. Feb. - Aug.

PESTICIDE SECTION

(Insecticides, Fungicides, Fumigants, Rodenticides, Vermicides, Weed Killers, Spreaders, Stickers, Emulsifiers—Pages 50 to 66.)

The sources of information and recommendations in connection with insect pests and diseases given in this catalog are the State Experiment Stations, The Kilgore Seed Company Laboratory, our technically trained field men, and many members of our store personnel who have had years of close contact with agriculture.

Every attempt has been made to provide such

information in a plain and simple form for the benefit of our growers. As, however, constant changes are taking place in the field of pesticidal problems because of new insect pests and diseases and the introduction of new pesticides, it is difficult to keep an annual catalog strictly up-to-date. We therefore invite you to contact our field and store men. They will be glad to assist you whenever you are facing new problems.

The use of pesticides is now just as necessary as that of fertilizers and various modern cultural practices. Many factors are involved in the use of pesticides in relation to results desired, such as the choice of proper pesticides for the control of different diseases and insects, and the time and method of application. One cannot expect satisfactory control with poor or wrongly timed applications. Too often the grower waits until the disease or insect has too much of a start before spraying or dusting. This usually is more expensive than starting off with a regular spray or dust program and keeping it up to prevent diseases and insects. He not only has to use excessive amounts of sprays or dusts to get control, but he also suffers considerable crop damage.

Now that the Miller Amendment governing the use of pesticides on crops is in effect, and a tolerance on the amount of residue on vegetables at harvest time has been established on most of the pesticides, it is very important that controlling of insects and disease start early, and the crop kept clean, so as to avoid

applying low tolerance pesticides too close to maturity or harvest.

See page 50 for more detailed information on the Miller Amendment.

As a rule, it is much cheaper to use a combination spray or dust when one must contend with both insects and diseases, as one application may do both jobs. Most insecticides and fungicides are compatible and can be used together in sprays or dusts. Although we manufacture numerous combination dusts, only the more important are listed in this catalog. If you do not find the combination listed which you want, write to The Kilgore Seed Company, Plant City, Florida, or call at any of our stores.

With the aid of newly installed mixing and blending equipment in our Plant City factory, you can be assured that you will receive fresh and perfectly blended dusts formulated to meet your particular requirements. Only high-grade technical materials are used by us in producing the various pesticidal compositions that you may desire.

SANITATION—Serious losses may result from neglect of a few simple rules. Diseases will spread readily from sick to healthy plants by various means, depending upon the nature of the disease. Spores of many species of fungi are air- or wind-borne, others may be scattered over fields in rain or irrigation water. Insects, particularly those of the sucking type, such as aphids and thrips disseminate diseases. Certain cultural operations, such as pruning, picking, transplanting, etc., may scatter diseases, especially those caused by bacteria and viruses (mosaic being a specific example of the latter).

Considering methods of infection, it becomes obvious that the source of the disease should be destroyed. Affected plants, as well as decayed fruit, should be removed from the field and burned. Cultivation should be clean and frequent. Mechanical injury to plants by plowing, spraying or picking operations should be avoided as much as possible. Crop rotation is essential, provided that no other crop affected by the responsible disease is grown for several years. This method often proves cheaper than an attempt to control the disease chemically. As weeds serve as host plants to many insects, which may be carriers of fungi, viruses and other undesirable organisms, the necessity for their eradication certainly becomes evident. Whenever practical, soil moisture should be kept fairly constant. Corrective measures should be adopted where soils are either too acid or too alkaline.

MOSAIC—Mosaic disease is caused by virus organisms in the sap of the affected plants. The viruses may be transmitted in various ways from diseased to healthy plants. Insects undoubtedly play an important part in the transmission of the disease, although it may be spread by other means, such as pruning operations, bruising of plants by workers or mechanical equipment, picking fruit, the use of tobacco and by other means.

The symptoms of mosaic vary slightly in different plants, but in general, certain broad characteristic indications can be noted in most all infected plant types. There is usually a stunting or dwarfing with accompanying pattern of light yellowish areas merging with green patches, giving a "mottled" effect. Often there is a roughening of the surface, producing, in some cases, a warty appearance. Edges of leaves may cup downward. The fruit also may be mottled and warted.

Mosaic has become costly and destructive on many crops, especially peppers, tomatoes, cucumbers and squash. This disease apparently does not carry over in the soil, provided cover crops are turned under long enough to be decomposed before planting. It is carried from season to season in wild host plants (weeds) in and around the field, and is transmitted to the plants in the field by insects, particularly aphids and thrips.

Pepper and tomato seedbeds should have clean cultivation two or three weeks before planting the seed. The seedbeds should not be close to ditch banks or fence rows, unless the weeds near the beds are destroyed. Smoking or other use of tobacco should not be permitted while working in the seedbeds or handling plants. Tobacco mosaic has been known to be carried on tobacco. If plants become affected with mosaic in the seedbeds, do not set them in the field. Once the plant gets the disease there is no known cure.

The State Experiment Stations recognize the severity of the problem and they are now very much engaged in the attempt to develop effective control measures by chemical or biological means. A serious effort is also being manifested in the development of resistant varieties of various vegetables commonly susceptible to virus diseases. In the event of success, such new varieties would hardly be available commercially for several years, and we would again like to stress the importance of sanitary practices and effective insect and weed eradication as the only means of crop damage reduction at the present time.

THE MILLER AMENDMENT . . . As it Concerns the Grower

THE MILLER AMENDMENT is a new law which affects all growers who use pesticide sprays and dusts in combatting insects, weeds, rodents and plant diseases.

It sets up new procedures for the Food and Drug Administration in Washington to use in establishing official limits on how much of the spray or dust can remain on a crop.

PESTICIDE SPRAYS AND DUSTS are essential to assure the continued supply of high quality foods. The new law is designed to permit the effective use of these sprays and dusts in the production of food without hazard to the consumer.

THE LAW WILL ENABLE THE GROWER to know with reasonable assurance what sprays and dusts he can use on his crops.

GROWERS WHO FOLLOW THE APPROVED DIRECTIONS for application and good agricultural practices should, under normal conditions, have no problem with pesticide residues.

ALL FRUITS, VEGETABLES AND GRAINS are covered by the law.

BY PESTICIDE RESIDUE is meant the amount of pesticide chemical left in or on the crop when it is harvested.

A TOLERANCE is the official term for the amount of pesticide residue which can legally remain in or on a crop. For example, a tolerance of five parts per million means that no more than five parts per million by weight of that chemical may legally remain in or on any food when it enters interstate commerce.

WHEN A TOLERANCE IS ESTABLISHED by FDA, it means that FDA is satisfied that when properly used, the residue, if any, of the chemical will cause no danger to public health. The tolerance set for a specific chemical does not apply to the use of that chemical on all crops, but only to the specific crops named in the FDA order. Sprays and dusts adhere to some crops more than to others.

A ZERO TOLERANCE on chemicals set by FDA means that such chemicals should be used in such a way that no residue will be left in or on crops. These chemicals

may be sprayed or dusted on crops if no residue remains at harvest time.

WHEN NO TOLERANCE HAS BEEN ESTABLISHED for a spray or dust, it means that if any residue of that chemical remains on a crop, the crop is subject to seizure and condemnation when it is introduced into interstate commerce. Some chemicals leave no residue and may not be given a tolerance for that reason.

A FEW CHEMICALS ARE EXEMPT from the requirement of a tolerance. FDA considers that any residue which results from the normal use of these chemicals on growing crops will be safe if the sprays or dusts are applied according to good agricultural practices. Examples of such chemicals are pyrethrum, rotenone, sabadilla, most pyrethrum synergists and most copper compounds (except arsenical copper compounds). All of the chemicals listed can be used up until harvest. **FOR YOUR AID, ESTABLISHED TOLERANCES TO DATE** are given in this catalog for chemicals offered by us (See below). Information regarding tolerances which may be established after the publication of this catalog will be made available to all of our stores and field men and we urge all growers to approach them at any time for this information.

It is not expected that any special problems or any major changes will be created, provided that the grower carries out instructions for the proper use of pesticides.

USE THE MATERIALS ONLY AS DIRECTED on the CROPS specified, in the AMOUNTS specified, and at the TIMES specified. Read the label before each use. Consult your local agricultural advisor on spraying and dusting schedules.

As to the number of days before harvest, present available information for local uses is scanty, but suggestions made and published by Florida Agricultural Experiment Station and U.S.D.A. on the safe number of days from last application to harvest on many of the pesticides. Much research has already been done on residues, and is still being carried on and we hope to have the answer soon. (See Circular 140, July 22, 1955, entitled "Some Questions and Answers on Vegetable Pesticide Tolerances.")

All of our stores and field men will be kept fully informed whenever additional information is available.

The Following Are Official Tolerances Established Up To Date

Pesticide	Trade Names	Tolerance p.p.m.	Pesticides	Trade Names	Tolerance p.p.m.
Aldrin	—	0.1	Lindane	Isotox, Kilgam	10.0
Aramite	—	1.0	Methoxychlor	Marlate	14.0
Benzene hexachloride	Gamtox, Kilhex	5.0	Malathion	—	8.0
Captan	Orthocide	20.0	Maneb	—	7.0
Chlordane	Ortho-Klor, Kilchlor	0.3	Nicotine	—	2.0
Copper Compounds:			Nabam	—	7.0
Bordeaux Mixture	—		Nicotine Sulphate	—	2.0
Copper oxychloride	Copper-A		Petroleum Oils	—	Exempt
Basic Copper	Kilcop		Parathion	Vapophos, Kilphos	1.0
Sulfate			Pyrethrum	Pyroicide	Exempt
Cuprous oxide	Cuprocide		Rotenone	—	Exempt
Cryolite	Kryocide	7.0	Sabadilla	—	Exempt
DDD	RHothane	7.0	Sulphur	—	Exempt
DDT	Kilcide	7.0	Systox	—	0.75
Dieldrin	—	0.1	TEPP	Vapotone	0
Endrin	—	.0	Toxaphene	Alltox, Kilphene	7.0
EPN	—	3.0	Zineb	Dithane Z-78, Parzate	7.0
Ferbam	Fermate	7.0	Ziram	Zerlate	7.0
Heptachlor	—	0.1			
Lead arsenate	—	7.0 of combined lead			

Safe Combinations of INSECTICIDES AND FUNGICIDES

Insecticides	Trade Names	Can be used with
AGRIMYCIN		Neutral Copper compounds, carbamates and most insecticides in wettable form.
STREPTOMYCIN		
PARATHION	(Vapophos)	DDT, DDD, methoxychlor, chlordane, benzene hexachloride, lindane, toxaphene, aldrin, dieldrin, Dithane, "Parzate," "Zerlate," "Fermate," "Manzate," and sulphur. Most copper compounds can be used with parathion and malathion for spraying mixtures if mixed freshly and used immediately.
MALATHION		
DDT	(Kilcide)	Each other, and also with nicotine, sulphur, copper, Dithane, "Parzate," Zerlate," "Fermate," "Manzate," and "Karathane."
DDD	(RHothane)	
METHOXYCHLOR	("Marlate")	
CHLORDANE	(Kilchlor)	
BENZENE HEXACHLORIDE	(Gamtox)	
LINDANE	(Isotox)	
TOXAPHENE	(Alltox)	
ALDRIN		DDT, DDD, methoxychlor, chlordane, benzene hexachloride, lindane, toxaphene, sulphur.
DIELDRIN		
TEPP	(Vapotone)	
NICOTINE		DDT, toxaphene, sulphur, copper, Dithane, "Parzate."
KRYOCIDE		Sulphur, copper, Dithane, "Parzate."

* Do not use emulsions with sulphur.

DONT'S

A Reference to this table may prevent serious crop losses

DON'T USE:	ON THE FOLLOWING CROPS:
DDT (Kilcide)	Cucurbits (cucumbers, squash, melons, etc.).
DDD (RHothane)	Cucurbits.
TOXAPHENE (Kilphene)	Cucurbits, young tomatoes*, tobacco, lettuce and escarolle**, avocados. Pole beans.
(Alltox)	
B. H. C. (Gamtox)	Cucurbits, corn, young tomatoes, potatoes and other tuber or root crops. Southern peas and beans.
(Kilhex)	
LINDANE (Isotox)	Corn, young cucurbits when heavy dew is present, potatoes and other tuber or root crops.
(Kilgam)	
CHLORDANE (Kilchlor)	Cucurbits, tobacco seedbeds and young tobacco plants in field.
KRYOCIDE	Corn, Asparagus plumosus.
COPPER COMPOUNDS	Corn.
TEPP (Vapotone XX)	Tomatoes.
(Vapotone Dust)	
ALDRIN	Don't use on any crops not approved by USDA. Consult local authorities.
DIELDRIN	

* As toxaphene may prove toxic to young tomato plants in some local areas, local experiment stations should be consulted before treating.

** Do not treat lettuce and escarolle within a period of three weeks from seeding or transplanting.

Avoid combinations of emulsions and wettable powders. Many are not compatible physically.

SEED TREATMENT

(For Disease Control)

The following recommendations are suggested:
Treatment for the prevention of damping-off fungi:

		Oz. Per Bu.	Teasps. Per Lb.
Beans, Lima	Spergon	2	1/3
Beans, Snap and Pole	Spergon	2	1/3
Peas, English	Spergon	2	1/3
Cowpeas	Spergon	2	1/3
Corn	Arasan or Ortho Seedguard 2		1/4
Broccoli	Arasan		1/2
Cantaloupe	Spergon or Semesan		1
Carrot	Spergon		3/4
Celery	Spergon or Semesan		3/4
Cabbage, Cauliflower, etc.	Arasan or Semesan		1/2
Cucumber	Spergon or Semesan		1
Eggplant	Semesan		3/4
Escarolle	Semesan		1
Lettuce	Semesan		1
Pepper	Semesan		1/2
Potato	Orthocide 50-W	4	
Spinach	Spergon or Arasan		1/2
Tomato	Semesan		1
Watermelon	Spergon or Semesan		1

The seed and seed treatment material should be thoroughly mixed together in a rotary drum mixer until all the seed is uniformly coated. In most cases this will take from three to five minutes at a speed of forty revolutions per minute. If Arasan is used on beans and peas, use one-half as much as the Spergon recommendation. Do not use more material than recommended. Screen off excess dust.

Bichloride of Mercury (Corrosive sublimate) treatment affecting organism on surface of seeds:

Cucumbers: Dilute 1:1000, immersing seed for 10 minutes.
Less hard seeds: Dilute 1:1000, immersing seeds for 5 minutes.
For small quantities of seed dissolve 1 tablet in 1 pint of water.
For larger quantities dissolve 1 oz. of the powder form in 7½ gallons of water. It is advisable to dissolve the powder in a small quantity (a pint or quart) of boiling water first.
Place seeds in loosely woven cloth bags (about half full). Submerge bags in solution and stir with a stick to loosen adhering bubbles. After soaking the proper length of time, remove bags and drain thoroughly. Seed should be washed for 15 minutes in several changes of water. Then spread in thin layers to dry. Do not prepare solution of bichloride of mercury in metal containers. Only glass, earthenware or wooden vessels should be used. In working with the compound, the greatest caution should be exercised, as it is extremely poisonous.

HOT WATER TREATMENT FOR CONTROL OF SEED-BORNE DISEASES

BLACK ROT. Cabbage seed which is not known to be free from the black rot organism should be soaked in water held at 122° F. for 25 minutes (cauliflower and broccoli, 18 minutes). Use an accurate thermometer and hold the temperature within 1° F. of that specified to control the disease without injuring the seed. Fill a cheesecloth bag two-thirds full of seed, tie the top, immerse it in the hot water and then stir it enough to remove the air. After treatment, dip the seed in cold water, drain and spread in a thin layer to dry. Use new land for seedbeds or old land in which cabbage or other related crops have not been grown for two years.
Do not use plants from a plant bed in which black rot has been found.

DILUTION TABLE FOR INSECTICIDES AND FUNGICIDES

Material	Average amounts needed to make spray		
	50 Gals.	4 Gals.	1 Gal.
Agri-mycin 100 at 200 p.p.m. level	9.34 oz.	4 tbsps.	1 tbsps.
Aldrin 2 Spray	1/2 pt.		
Alltox 40-W	2 lbs.	8 tbs.	2 tbs.
Alltox Spray No. 4.8	1 1/2 pts.	3 3/4 tbs.	4 tsp.
Aramite	3/4 lbs.	4 tbs.	1 tbs.
Arsenate of Lead	1 1/2 lbs.	10 tbs.	2 1/2 tbs.
Black Leaf 40	1 pt.	8 tsp.	2 tsp.
Ceresan	1 lb.	3 tbs.	2 1/4 tsp.
C-M-Z Bean Spray	5 lbs.	20 tbsps.	5 tbsps.
C-M-Z Spray Improved	3 lbs.	14 tbs.	3 1/2 tsp.
C-M-Z Garden Spray	2 1/2 lbs.	8 tbs.	2 tbs.
Copper-A Compound	2 lbs.	8 tbs.	2 tbs.
Cuprocide—Yellow	1 lb.	2 tbs.	1 1/2 tsp.
DDT 25% Emulsion	1 pt.	5 tbs.	4 tsp.
DDT 50% Wettable	1 1/4 lbs.	8 tbs.	2 tbs.
Dieldrin Emulsion (1.5)	Consult label		
Dithane D-14 (Nabam)	1 qt.	5 tbs.	4 tsp.
(Zinc Sulphate)	6 oz.	3 tsp.	3/4 tsp.
Dithane M-22 (Maneb)	1 lb.	5 tbs.	4 tsp.
Dithane Z-78 (Zineb)	1 lb.	5 tbs.	4 tsp.
Endrin	Consult label		
Fermate	1 lb.	9 1/3 tbs.	7 tsp.
Gamtox 10 Wettable	1 1/4 lbs.	8 tbs.	2 tbs.
Heptachlor—2 Spray	Consult label		
Improved Isotox Garden Spray M	1 pt.	8 tsp.	2 tsp.
Isotox Spray No. 200	1/2 pt.	4 tsp.	1 tsp.
Isotox Wettable No. 25	1/2 lb.	4 tbs.	1 tbs.
Karathane WD	Consult label		
Kelthane	Consult label		
Kilchlor 40-W (Chlordane)	1 1/4 lbs.	6 3/4 tbs.	4 tsp.
Kilcop "45"	Consult label		
Kilcop "53"	2 lbs.	10 tbs.	2 1/2 tbs.
Kryocide	3 lbs.	13 tbs.	3 1/3 tbs.
Malathion "50" Spray	1 qt.	5 tbs.	4 tsp.
Malathion 25% Wet.	1 lb.	7 tbs.	5 tsp.
Marlate 50 Methoxychlor Spray	1 1/2 lbs.	11 tbs.	8 tsp.
Manganese Sulphate	1 1/2 lbs.	6 2/3 tbs.	5 tsp.
Manzate (Maneb)	3/4 lb.	3 1/4 tbs.	4 tsp.
Oil Emulsion (Fico)	1 gal.	10 1/2 oz.	5 1/2 tbs.
Orthotran 50W	1/2 lb.	2 1/2 tbsps.	2 tsp.
Ortho-Klor 40-W	1 1/4 lbs.	8 tbs.	2 tsp.
Ortho Streptomycin (200 PPM)	8 oz.	4 tbs.	1 tbs.
Orthocide 50-W	2 lbs.	12 tbs.	3 tbs.
Orthol Garden Spray	1 1/2 qts.	7 1/2 tbs.	5 1/2 tsp.
Ortho-Klor 44 Spray (Chlordane)	1 1/2 qts.	8 tbs.	2 tbs.
Ortho-Klor 8 Spray (Chlordane)	Consult label		
Orthophos 4 Spray (Parathion)	1/3 pt.		
Parzate (Zineb)	1 lb.	5 tbs.	4 tsp.
Phygon-XL	1/2 lb.	3 tbs.	2 1/4 tsp.
Red Arrow Insect Spray	1 pt.	8 tsp.	2 tsp.
RHothane WP-50	1 1/4 lbs.	8 tbs.	2 tbs.
RHothane 25% Emulsion	1 pt.	8 tsp.	2 tsp.
Spergon Wettable	1 1/2 lbs.	11 tbs.	8 tsp.
Sulfocide	1 qt.	5 tbs.	4 tsp.
Systox	3/4 pt.	2 tbs.	1 1/2 tsp.
Triton B-1956	3 tbs.	3/4 tsp.	20 drops
Vapophos 15 (Parathion)	1/2 lb.		
Vapotone-XX Spray (TEPP)	4 oz.		
Volck Isotox Spray	1 gal.	20 tbs.	5 tbs.
Volck Oil Spray	Consult label		
Zerlate	1 lb.	5 tbs.	4 tsp.
Zinc Sulphate	1 lb.	3 tbs.	2 tsp.

BUREAU OF STANDARDS HOUSEHOLD MEASUREMENTS

Tsp.—level teaspoon	Gal.—gallon	1 Cup—8 ounces
Tbs.—level tablespoon	3 Tsp.—1 tbs.	2 Cups—1 pint
Pt.—pint	2 Tbs.—1 ounce	2 Pts.—1 quart
Qt.—quart	16 Tbs.—1 cup	4 Qts.—1 gal.
Oz.—ounce		

Sprays or dusts required per acre depend chiefly upon stage of growth, variety of crop and severity of insect infestation or disease infection. Spray for small plants require from 75 to 100 gals. per acre, for larger plants and heavy growth 100 to 150 gals. per acre. Dusts for small plants 25 to 30 lbs. per acre, for large plants and heavy growth 35 to 50 lbs. per acre.

INSECT AND DISEASE RECOMMENDATIONS

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 50.

BEANS—

INSECTS

		Coined Name	Common Name	
Leaf miners	Kilphos Dust No. 2	Parathion	20-30 lbs., acre	
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.	
Bean leaf hoppers	Kilphos Dust No. 2	Parathion	25 lbs., acre	
	Kilcide 5-Sulphur Dust	DDT	20-35 lbs., acre	
	Pyrocide Dust 7½	Pyrethrum	20-30 lbs., acre	
	DDT Wettable—50%		2 lbs., 100 gals.	
Thrips	Kilchlor-Kilphene Poison	Chlordane	15-30 lbs., acre	
	Bait No. 11	Toxaphene		
Army worms	Kilphene Poison Bait No. 9	Toxaphene	15-30 lbs., acre	
	Kilchlor Poison Bait No. 8	Chlordane	15-30 lbs., acre	
Cutworms				
Mexican bean beetles	Kilphos Dust No. 2	Parathion	20-30 lbs., acre	
	Rotenone Dust No. 1		20-30 lbs., acre	



Adult
BEAN LEAF HOPPER


DISEASES

Rust	Sulphur Dust	20-40 lbs., acre
Powdery mildew	Sulphur—Wettable	10-16 lbs., 100 gals.

BROCCOLI, CABBAGE, CAULIFLOWER—

INSECTS

Aphids	Kilphos Dust No. 4	Parathion	20-30 lbs., acre
	Vapotone XX Spray	TEPP	1 pt., 100 gals.
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
	Orthophos 4-Spray	Parathion	1/3 pt., 100 gals.
Cabbage worms	Kilchlor 5% Dust	Chlordane	25-35 lbs., acre
	Kilphene 10 Dust	Toxaphene	25-35 lbs., acre
	Kilgam Dust No. 2	Lindane	20-30 lbs., acre
	Kryocide Dust 30-70		25-35 lbs., acre
	Marlate 50	Methoxychlor	3 lbs., 100 gals.
	Marlate Dust No. 1	Methoxychlor	25-35 lbs., acre
	Rotenone Dust No. 1		25-35 lbs., acre
	25% DDT Emulsion		1 qt., 100 gals.
Leaf miners	Kilchlor 40-W	Chlordane	2½ lbs., 100 gals.
	Kilphos Dust No. 2	Parathion	25-35 lbs., acre
	Kilchlor-Sulphur Dust No. 3	Chlordane	25-35 lbs., acre
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
Cutworms			
Army worms	Kilchlor-Kilphene Poison	Chlordane	15-30 lbs., acre
	Bait No. 11	Toxaphene	
Harlequin cabbage bug	Kilphene Poison Bait No. 10	Toxaphene	15-30 lbs., acre
Grasshoppers	Kilchlor Poison Bait No. 8	Chlordane	15-30 lbs., acre
Mole crickets			



CABBAGE LOOPER



Adult
HARLEQUIN
CABBAGE BUG

DISEASES

For Seedbeds		Consult Local Experiment Station for rate of application.
Downy mildew	Spergon Dust No. 3	2 lbs., 100 gals.
	Spergon Wettable	
For Crops in Field		2 qts. plus ¾ lb., 100 gals.
Alternaria leaf spot	Nabam plus zinc sulphate	2 lbs., 100 gals.
Downy mildew	Zineb	


CELERY—

INSECTS


Cutworms	Alltox 40 W	Toxaphene	3 lbs., 100 gals.
	Kilphene Poison Bait No. 9	Toxaphene	15-30 lbs., acre
Army worms			
Celery leaf tiers	Kilphene-10 Dust	Toxaphene	20-30 lbs., acre
	25% DDT Emulsion		1 qt., 100 gals.
Celery loopers	Marlate 50	Methoxychlor	3 lbs., 100 gals.
	Marlate Dust No. 1	Methoxychlor	25-35 lbs., acre
Aphids	Kilphos Dust No. 1	Parathion	20-30 lbs., acre
	25% DDT Emulsion		1 qt., 100 gals.
Thrips	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.

DISEASES

Early blight	Nabam plus zinc sulphate	Nabam	2 qts. plus ¾ lb., 100 gals.
	Kilcop "53"	Copper	4 lbs., 100 gals.
	Copper A Compound		6 lbs., 100 gals.
	Maneb		1½ lbs., 100 gals.
Late blight			
Bacterial leaf spot in seedbeds	Agrimycin 200 ppm plus 4 lbs. neutral copper, 1 lb. 3 oz., 100 gals.		
	Streptomycin 200 ppm plus 4 lbs. neutral copper, 1 lb. 100 gals.		



IMPORTED CABBAGE WORM







CELERY LEAF-TIER

INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 50.

CORN (SWEET) —

INSECTS

		Coined Name	Common Name	
Corn earworms Corn bud worms Fall army worms	Kilphos Dust No. 4	Parathion	30-35 lbs., acre	
	Kilphene-10 Dust	Toxaphene	30-35 lbs., acre	
	Kilcide 10 Dust	DDT	30-35 lbs., acre	
	Vapophos Wettable 15%	Parathion	2 lbs., 100 gals.	
	Alltox 40 W	Toxaphene	2½ lbs., 100 gals.	
	DDT Wettable—50%		2 lbs., 100 gals.	
Corn lantern flies Corn silk flies	25% DDT Emulsion		1 qt., 100 gals.	
	Kilphos Dust No. 4	Parathion	30-35 lbs., acre	
Cutworms Wire worms	Vapophos Wettable 15%	Parathion	2 lbs., 100 gals.	
	Alltox 40 W	Toxaphene	2½ lbs., 100 gals.	
Corn leaf blight	Kilchlor 40 W	Chlordane	4 lbs., 100 gals.	
	Aldrin Emulsion		Broadcast 2 gals., acre in 100 gals. water	
	Zineb Dust No. 2		25-30 lbs., acre	
	Zineb		2 lbs., 100 gals.	
	Nabam plus zinc sulphate		2 qts. plus ¾ lb., 100 gals.	

CORN EARWORM
(TOMATO FRUITWORM)

WIRE WORM

COWPEAS (SOUTHERN PEAS) —

INSECTS






Pod weevils	Kilphene 5—Sulphur Dust	Toxaphene	25-35 lbs., acre
	Kilcide 5—Sulphur Dust	DDT	20-35 lbs., acre

DISEASES

Scab and leaf blight	Kilcop Dust No. 1	Copper	25-35 lbs., acre
	Nabam plus zinc sulphate		2 qts. plus ¾ lb., 100 gals.

CUCUMBER, SQUASH, CANTALOUPE —

INSECTS

Aphids	Kilphos Dust No. 1	Parathion	25-35 lbs., acre	
	Kilgam Dust No. 2	Lindane	25-35 lbs., acre	
	Pyrocide Dust 7½		20-30 lbs., acre	
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.	
	Vapotone XX	TEPP	1 pt., 100 gals.	
	Isotox Wettable No. 25	Lindane	1 lb., 100 gals.	
Leaf miners	Kilphos Dust No. 2	Parathion	30-35 lbs., acre	
	Vapophos 15%	Parathion	1 lb., 100 gals.	
Cucumber beetles	Kilgam Dust No. 2	Lindane	25-35 lbs., acre	
	Kryocide Dust 30-70		25-35 lbs., acre	
Pickle worms	Kilphos Dust No. 1	Parathion	25-35 lbs., acre	
Melon worms	Marlate 50	Methoxychlor	3 lbs., 100 gals.	
Squash bugs	Marlate Dust No. 1	Methoxychlor	25-35 lbs., acre	
	Isotox Wettable No. 25	Lindane	1 lb., 100 gals.	
	Kryocide Wettable		6 lbs., 100 gals.	
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.	

Adult
STRIPED CUCUMBER
BEETLE

SQUASH BUG

DISEASES

Downy mildew	Zineb Dust No. 1		25-35 lbs., acre
	Powdery mildew	Nabam plus zinc sulphate	2 qts. plus ¾ lb., 100 gals.
Powdery mildew only	Karathane	Capryl	6 to 8 oz., 100 gals.
Angular leaf spot	Karathane Dust 1%	Capryl	20-30 lbs., acre
	Anthracnose	Orthocide 50-W	Captan.....4 lbs., 100 gals.

EGGPLANT —

INSECTS

Aphids Thrips	Kilphos Dust No. 2	Parathion	20-60 lbs., acre
	Kilhex-Sulphur Dust—1½%	BHC	20-60 lbs., acre
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
	Vapotone XX	TEPP	1 pt., 100 gals.
	Gamtox Wettable No. 10	BHC	2½ lbs., 100 gals.
	Dusting Sulphur		20-60 lbs., acre
Red spiders	Vapotone XX	TEPP	1 pt., 100 gals.
	Kilphos Dust No. 1	Parathion	20-60 lbs., acre
Cutworms Mole crickets	Kilchlor-Kilphene	Chlordane	15-30 lbs., acre
	Poison Bait No. 11	Toxaphene	

LETTUCE, ESCAROLLE —

INSECTS

Cutworms Mole crickets Aphids and loopers	Kilchlor-Kilphene	Chlordane	15-30 lbs., acre
	Poison Bait No. 11	Toxaphene	
	Pyrocide Dust 7½		20-30 lbs., acre



SQUASH VINE BORER

INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 50.

OKRA—

INSECTS

		Coined Common Name	
Aphids	Vaportone XX	TEPP	1 pt., 100 gals.
	Kilphos Dust No. 1	Parathion	25-35 lbs., acre
Leaf miners	Kilphene 10 Dust	Toxaphene	25-35 lbs., acre
Pumpkin bugs	Alltox Wettable No. 40	Toxaphene	2½ lbs., 100 gals.
Okra caterpillars			



GREEN PUMPKIN BUG
(SOUTHERN STINK BUG)

ONION—

INSECTS

Thrips	Gamtox Wettable No. 10	BHC	2½ lbs., 100 gals.
	DDT 25% Emulsion		1 qt., 100 gals.

PEPPER—

INSECTS

Aphids	Kilphos Dust No. 1	Parathion	20-30 lbs., acre
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
	Black Leaf 40		1 qt., 100 gals.
Pepper weevils	Kilhex Dust No. 7	BHC	25-30 lbs., acre
Fall army worms	DDT 25% Emulsion		1 qt., 100 gals.
Blister beetles	Gamtox Wettable No. 10	BHC	2½ lbs., 100 gals.
	Marlate 50	Methoxychlor	3 lbs., 100 gals.
	Marlate Dust No. 1	Methoxychlor	25-35 lbs., acre



PEPPER WEEVIL

DISEASES

Cercospora leaf spot	Kilcop Dust No. 6	Copper	30-35 lbs., acre
	Kilcop "53"	Copper	5 lbs., 100 gals.
Bacterial leaf spot	Kilcop Dust No. 6	Copper	30-35 lbs., acre
Phytophthora blight	Nabam plus zinc sulphate	Nabam	2 qts. plus ¾ lb., 100 gals.
	Zineb	Zineb	2 lbs., 100 gals.
Bacterial leaf spot only	Agrimycin (200 PPM)		1 lb. 3 oz., 100 gals.
	Streptomycin (200 PPM)		1 lb., 100 gals.
Damp-off	Orthocide 50-W	Captan	4 lbs., 100 gals.
	Yellow Cuprocid	Copper	2 lbs., 100 gals.

POTATOES—

INSECTS

Aphids	Kilphos Dust No. 1	Parathion	20-40 lbs., acre
Army worms	Kilphene-5 Dust	Toxaphene	20-40 lbs., acre
Cutworms	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
Colorado potato beetles	Alltox Wettable No. 40	Toxaphene	2½ lbs., 100 gals.
	DDT Emulsion—25%		1 qt., 100 gals.
Leaf miners	Same as above except omit DDT Emulsion 25%		
Pumpkin bugs			
Leaf-footed plant bugs			

DISEASES

Late blight	Zineb Dust No. 2		25-35 lbs., acre
	Kilcop Dust No. 6	Copper	25-35 lbs., acre
Early blight	Nabam plus zinc sulphate		2 qts. plus ¾ lb., 100 gals.
	Zineb		2 lbs., 100 gals.
	Kilcop "53"	Copper	4 lbs., 100 gals.



Adult
COLORADO POTATO BEETLE

POTATOES (SWEET)—

INSECTS

Caterpillars	Kilphene-10 Dust	Toxaphene	25-35 lbs., acre
Gold bugs	Alltox Wettable No. 40	Toxaphene	2½ lbs., 100 gals.
Grasshoppers			

RADISH, TURNIP, MUSTARD —

INSECTS

Aphids	Pyrocide Dust—7½	Pyrethrum	15-35 lbs., acre
	Vaportone XX	TEPP	1 pt., 100 gals.



Wingless Adult
APHID

INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 50.

STRAWBERRIES—


INSECTS

		Coined Common Name	
Thrips	Malathion 25-W		2 lbs., 100 gals.
	Kilphos Dust No. 1		20-25 lbs., acre
Pameras			
Ants			
Flea beetles	Pyrocide Dust 7½	Pyrethrum	20-25 lbs., acre
Cucumber beetles	Kilchlor-Sulphur Dust No. 3	Chlordane	20-25 lbs., acre
Leaf rollers			
Grasshoppers			
Mole crickets	Kilchlor-Sulphur Dust No. 3	Chlordane	20-25 lbs., acre
Corn borers	Kilchlor Poison Bait No. 8	Chlordane	15-25 lbs., acre
Cutworms	Kilchlor-Kilphene Poison	Chlordane	
Field crickets	Bait No. 11	Toxaphene	20-25 lbs., acre
	Dusting Sulphur		20-30 lbs., acre
Red spiders	Malathion 25-W		2 lbs., 100 gals.
	Kilphos Dust No. 1		20-25 lbs., acre


Note: Use Chlordane before fruiting only.

DISEASES

Anthracnose	Kilcop Dust No. 2	Copper	20-30 lbs., acre
Leaf spots	Kilcop "53"	Copper	4 lbs., 100 gals.
	Zineb Dust No. 1		20-30 lbs., acre
	Maneb		1½ lbs., 100 gals.



RED SPIDER



CUTWORM


TOMATOES—

INSECTS

Aphids	Kilphos Dust No. 1	Parathion	25-35 lbs., acre
Thrips	Kilhex Dust 1½%	BHC	25-35 lbs., acre
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
	Gamtox Wettable No. 10	BHC	2 lbs., 100 gals.
Army worms	Kilphos Dust No. 1	Parathion	25-35 lbs., acre
Cutworms	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
Tomato fruit worms	DDT 25% Emulsion		1 qt., 100 gals.
Pin worms	DDT 50% Wettable		2 lbs., 100 gals.
	RHothane WP-50	TDE—DDD	2 lbs., 100 gals.
	Kilchlor-Kilphene Poison	Chlordane	20-25 lbs., acre
	Bait No. 11	Toxaphene	
Leaf miners			
Pumpkin bugs	Kilphos Dust No. 1	Parathion	25-35 lbs., acre
Leaf-footed plant bugs	Vapophos Wettable 15%	Parathion	1 lbs., 100 gals.
Horn worms	Kryocide-Sulphur Dust 30-70		35-45 lbs., acre
	RHothane 25% Emulsion	TDE—DDD	1 qt., 100 gals.
	Marlate 50	Methoxychlor	3 lbs., 100 gals.
	Marlate Dust No. 1	Methoxychlor	25-35 lbs., acre



TOMATO PINWORM



Nymph THRIPS

DISEASES

	Maneb		1½ lbs., 100 gals.
Late blight	Zineb-Sulphur Dust No. 4	Zineb	30-40 lbs., acre
Early blight	Kilcop-Sulphur Dust No. 3	Copper	30-40 lbs., acre
Gray leaf-spot (Red rust)	Nabam plus zinc sulphate		2 qts. plus ¾ lb., 100 gals.
	Kilcop "53"		4 lbs., 100 gals.
	Zineb		2 lbs., 100 gals.
	Phygon XL		¾ lb., 100 gals.
Bacterial leaf spot	Agri-mycin (200 PPM)		1 lb. 3 oz., 100 gals.
	Streptomycin (200 PPM)		1 lb., 100 gals.


WATERMELON—

INSECTS

Pumpkin bug			
Leaf-footed plant bug			
Squash bug	Kilphos Dust No. 1	Parathion	20-35 lbs., acre
Other insect same as Cucumbers, etc.	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
Page 54			

DISEASES

Downy mildew	Zineb Dust No. 1	Zineb	30-40 lbs., acre
Anthracnose	Kilcop Dust No. 5	Copper	30-40 lbs., acre
Gummy stem blight	Zineb	Zineb	2 lbs., 100 gals.
	Kilcop "53"	Copper	4 lbs., 100 gals.
	Orthocide 50-W	Captan	4 lbs., 100 gals.



LEAF-FOOTED PLANT BUG

INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 50.

TOBACCO—

INSECTS

		Coined Name	Common Name	
Aphids	Kilphos Dust No. 1	Parathion	10-35 lbs., acre	
	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.	
	Vapotone XX	TEPP	1 pt., 100 gals.	
Bud worms	RHothane 5-Tobacco Dust	TDE-DDD	15-35 lbs., acre	
	DDT 50% Wettable		2 lbs., 100 gals.	
	RHothane WP-50	TDE-DDD	2 lbs., 100 gals.	
Horn worms	RHothane 5-Tobacco Dust	TDE-DDD	25-30 lbs., acre	
	Kryocide Dust 30-70		15-35 lbs., acre	
	RHothane WP-50	TDE-DDD	2 lbs., 100 gals.	
Mole crickets	Kilchlor Poison Bait No. 8*	Chlordane	15-25 lbs., acre	

* CAUTION: Do not use directly on plants.
Scatter in paths or walk ways.

DISEASES—PLANT BEDS

Blue mold	Zineb Dust No. 2	Zineb	1½ to 4 lbs. to 100 sq. yds.
	Zineb Wettable	Zineb	4 lbs., 100 gals.
	Fermate Wettable	Ferbam	3 lbs., 100 gals.



Adult
MOLE CRICKET

PEANUTS—

INSECTS

Velvet bean caterpillars	Kilcide 5—Sulphur Dust	DDT	25-35 lbs., acre
	Kryocide—Sulphur Dust 30-70		25-35 lbs., acre

DISEASES

Leaf spot	Dusting Sulphur	15-25 lbs., acre
-----------	-----------------	------------------



Adult
THRIPS

GLADIOLUS—


INSECTS

Thrips	Kilcide-Kilhex Dust No. 3*	DDT-BHC	25-35 lbs., acre
	DDT 50% W in combination with Gamtox W No. 10*	BHC	2 lbs. plus 1¼ lbs. Gamtox, 100 gals.
	DDT 50% Wettable		4 lbs., 100 gals.
Army worms	Dieldrin Emulsion (1.5 lbs. technical per gal.)		3 pts., 100 gals.
	Kilphos Dust No. 2	Parathion	25-35 lbs., acre
Red spider	Vapophos Wettable 15%	Parathion	1 lb., 100 gals.
	Wettable Sulphur		12 lbs., 100 gals.

DISEASES

Curvularia	Orthocide 50-W	Captan	4 lbs., 100 gals.
Botrytis	Nabam plus zinc sulphate	Nabam	2 qts. plus ¾ lb., 100 gals.
Stemphylium			

* CAUTION: Kilhex and Gamtox Wettable may give spikes a musty odor if sprayed just prior to shipping.



Adult
CHINCH BUG

LAWNS—

INSECTS

Chinch bugs	Kilgore's 25% Emulsion Concentrate	1 qt. to 1000 sq. ft.
	Kilgore's Kil-Chinch	Treat infested areas with 2 lbs., 100 sq. ft.
Army worms	Saba Chinch	Treat infested areas with 4 lbs., 1000 sq. ft.
	Kilgore's 25% Emulsion Concentrate	1 qt. to 1000 sq. ft.
Nematodes	Kilgore's Nema-Kil	1 qt. to 600-800 sq. ft.

NOTE: See our 1956-57 Annual Flower Guide for Florida, pages 28, 29, 30, 31.

PASTURES—

INSECTS

Aphids	Vapotone XX	TEPP	1 pt., 100 gals.
	Vapophos 15%	Parathion	1 lb., 100 gals.
Sod webworms	RHothane 5-Sulphur Dust*	TDE-DDD	20-30 lbs., acre
Army worms	Kilphene 10-Dust*	Toxaphene	20-30 lbs., acre
Grasshoppers			

* CAUTION: Keep cattle off pastures for 2 weeks after applying these insecticides. Do not use in pastures for dairy cows.



Nymph
CHINCH BUG


The Kilgore Seed Company processes many combinations of insecticides and fungicides to suit local conditions which are too numerous to recommend in our General Catalog. If you do not find the combination you want, write our General Office in Plant City or call at your nearest Kilgore Store; it is likely we will have a dust in stock to fill your requirements.

INSECT AND DISEASE RECOMMENDATIONS (Continued)

For detailed information on the Miller Amendment governing the use of pesticides on crops, see page 50.

SOIL BORNE PESTS—

Nematodes	DD Soil Fumigant, see page 64
As preplanting treatment	Methyl Bromide, see page 64
	Soilfume (EDB), see page 64
	Vapam
Nematodes	
On established plants	Kilgore's Nema-Kil, see page 64
Wire worms and other soil insects	Aldrin 2 Spray 2 gals., acre in 100 gals. water
	Ortho-Klor 8 2 gals., acre in 100 gals. water
	Heptachlor 2 Spray 2 gals., acre in 100 gals. water



WIRE WORM

Termite Control

		Coined Common Name	
Termites	Ortho-Klor 8 Spray	Chlordane	5 qts., 100 gals.
	Ortho-Klor 44 Spray	Chlordane	2 gals., 100 gals.



SUBTERRANEAN TERMITE
WINGED ADULT

Apply to infected soil areas with low pressure spraying equipment. Thoroughly soak area. Where outside trenching operations next to foundation walls are desired, dig trench 1 to 2 ft. deep and 8" to 12" wide. Apply at the rate of one gallon per linear foot of trench. Also treat soil as trench is being refilled. For treating fill dirt under concrete, apply 3 gals. diluted material per sq. foot—Make diluted spray with 1/3 pint 75% emulsion to 25 gals. water. If above recommendations are followed, present results show that the protection period should be at least five years.

Livestock Pests


DAIRY CATTLE—

		Coined Common Name	
Flies	"Marlate" 50-Insecticide	Methoxychlor	8¾ lbs., 100 gals.
Lice	"Marlate" 50-Insecticide	Methoxychlor	25 lbs., 100 gals.

DAIRY BARNs AND MILK SHEDS—

Flies	Isotox Dairy Spray	Lindane	10 lbs., 100 gals.
	"Marlate" 50-Insecticide	Methoxychlor	25 lbs., 100 gals.
	Ortho Fly Killer		Follow label directions

CAUTION: All feed troughs, feed and utensils should be covered before spraying.



ADULT CATTLE
FEVER TICK

RANGE CATTLE—

SPRAYS

(Use one of following)

Horn flies	DDT 50% Wettable		8 lbs., 100 gals.
Lice	Gamtox Wettable No. 10	BHC	3¾ lbs., 100 gals.
Mosquitoes	Isotox Dairy Spray	Lindane	1½ lbs., 100 gals.
	"DuPont Livestock Spray and Dip No. 30"	BHC & DDT	11 lbs., 100 gals.

(Use one of following)

Mange mites	Kilchlor 40-W		5 lbs., 100 gals.
	Gamtox Wettable No. 10	BHC	2½ lbs., 100 gals.

DIPS*

(Use one of following)

General purpose dip	DDT 50% Wettable		8½ lbs., 100 gals.
	Gamtox Wettable No. 10	BHC	2½ lbs., 100 gals.

Combination of:	Ortho Healthy Herd Wettable Powder	BHC & DDT	10 lbs., 100 gals.
	"DuPont Livestock Spray and Dip No. 30"	BHC & DDT	11 lbs., 100 gals.
	Ortho Kleen Stock Spray or Dip	Lindane and Toxaphene	See label for directions



ADULT
HORN FLY

***CAUTION:** In using dipping vats, all traces of arsenicals, oils or sludge from such materials must be thoroughly removed before charging with Gamtox, Isotox, "Marlate" or DDT. A mixture of arsenicals or oil solutions with Gamtox, Isotox, "Marlate," or DDT is highly toxic to animals and may cause death. Directions on all containers should be carefully followed.

Screw worms	"Smearex," Screw Worm Preparation		
	"Ortho 1038 Screw Worm Control"		Follow Label Directions
	Ortho EQ 335 Screw Worm Remedy		

PESTICIDES

For household and garden insecticides, see Page 63.

When ordering pesticides allow enough for postage because all prices quoted are f.o.b. Plant City or any Kilgore store unless otherwise specified. Inquire of your postmaster for proper parcel post rates.

All pesticides not used for commercial agricultural purposes are subject to 3% State Sales Tax.

All prices quoted are subject to change without notice.

INSECTICIDES

ALDRIN 2 SPRAY—A liquid formulation containing 2 lbs. aldrin per gallon. 1 gal. \$4.65; 5-gal. can \$19.25. 30 gals. \$111.00; 53 gals. \$190.80.

ARAMITE—15-W. An effective miticide. Used on citrus, ornamentals and vegetables. Controls red spider mites. Use 1½ lbs. per 100 gals. Price 4-lb. bag \$2.88; 50 lbs. \$34.50.

ARSENATE OF LEAD POWDER. Use 6 lbs. to 100 gals. water. Pound canister 60c; 4 lbs. \$1.45

BENZENE HEXACHLORIDE PREPARATIONS.

GAMTOX WETTABLE—No. 10 (Contains 10% gamma isomer.) Use 2½ lbs. per 100 gals. water. 5-lb. bag \$1.60; 50-lb. bag \$14.50.

KILHEX DUST—No. 7 (Code No. G-3)—(Gamtox) Contains 1½% gamma isomer. Lb. 10c; 25 lbs. \$1.90; 50 lbs. \$2.75; 100 lbs. \$5.50.

KILHEX-SULPHUR DUST—(Code No. G-4)—(Gamtox) Contains 1½% gamma isomer. Lb. 10c; 25 lbs. \$2.20; 50 lbs. \$3.40; 100 lbs. \$6.80.

BLACK LEAF 40. For the control of aphids, thrips. Use from 1 to 2 pts. per 100 gallons water. Is effective as a spray against poultry lice. 1 oz. 49c; 5 oz. \$1.39; 1-lb. \$2.98; 2 lbs. \$5.29; 5 lbs. \$7.60; 10 lbs. \$14.70. Prices quoted are prepaid.

CALCIUM ARSENATE POWDER. 1-lb. can 45c; 4-lb. bag 65c.


CHLORDANE PREPARATIONS

KILCHLOR 40-W. (Contains 40% technical chlordane.) Use 2½ lbs. per 100 gals. of water. 1-lb. can 90c; 5-lb. bag \$3.35; 10-lb. bag \$6.50; 25 lbs. \$13.25; 50 lbs. \$25.50.

ORTHO-KLOR "44"—(Containing 44% liquid chlordane). Readily emulsifiable in water. For control of ants, mole crickets, cutworms, army worms, grasshoppers, roaches and many other insects. 4 oz. 69c; 8 oz. \$1.19; pt. \$1.98; qt. \$2.98.

ORTHO-KLOR-8 SPRAY. (Contains 72% chlordane 1 gal. \$11.20; 5 gals. \$49.75. Use 2 qts. per 100 gallons for soil applications.

KILCHLOR 5% DUST (Code No. H-1)—(Contains 5% chlordane.) Lb. 15c; 25 lbs. \$2.50; 50 lbs. \$3.95; 100 lbs. \$7.90.



KILCHLOR-SULPHUR DUST NO. 3 (Code No. H-3)—(Contains 5% chlordane). Lb. 15c; 25 lbs. \$2.85; 50 lbs. \$4.65; 100 lbs. \$9.30.

DDT PREPARATIONS

DDT 50% WETTABLE POWDER. Use 2 lbs. to 100 gals. water. 1-lb. canister 70c; 3-lb. bag 99c; 4 lbs. \$1.32; 6-lb. bag \$1.98; 50-lb. bag \$15.00.

DDT 25% EMULSION. Use 2 qts. per 100 gals. water. Qt. \$1.35; gal \$3.25; 5 gals. \$12.50; 30-gal. drum \$66.00; 50 gals. \$100.00.

KILCIDE 5 DUST—(Code No. D-3)—(Contains 5% DDT). Lb. 10c; 25 lbs. \$1.65; 50 lbs. \$2.35; 100 lbs. \$4.70.

KILCIDE 5-SULPHUR DUST—(Code No. D-10)—(Contains 5% DDT). Lb. 10c; 25 lbs. \$2.00; 50 lbs. \$3.00; 100 lbs. \$6.00.

KILCIDE 10 DUST—(Code No. D-4)—(Contains 10% DDT). Lb. 10c; 25 lbs. \$2.20; 50 lbs. \$3.45; 100 lbs. \$6.90.

KILCIDE 5-MANGANESE-SULPHUR DUST — (Code No. DX-2)—(Contains 5% DDT). Lb. 10c; 25 lbs. \$2.15; 50 lbs. \$3.25; 100 lbs. \$6.50.

DIELDRIN EMULSION. (Contains 1.5 lbs. technical dieldrin per gal.) 1 gal. \$8.00; 5 gals. \$38.75; 50 gals. \$375.00.

DIELDRIN—10% Granular. Used for control of flies, mosquitoes, beetles, etc. Consult label for complete information. 5-lb. bag \$2.75; 50-lb. bag \$24.00.

DIELDRIN—50% wettable. For the control of many insects. Consult label for complete information. 5 gal. \$10.20.

ENDRIN—1.6 spray. Use strictly in accordance with label cautions, warnings and directions. 1 gal \$13.20; 5 gal. \$64.50.

HEPTACHLOR—2 Spray. For control of wire worms and many soil borne pests. Consult label. 5 gals. \$19.75; 30 gal. \$114.00; 55 gal. \$203.50.

KELTHANE FC EMULSION. New effective miticide for use on flowers and ornamentals. Use 1 pint to 1 qt. in 100 gals. water, or 1 to 2 teaspoons per gallon. 1 gal. \$15.00; 5 gal. \$72.50.

KELTHANE-W—Wettable powder. Use 1 to 2 lbs. per 100 gal. water, 4 to 8 level tablespoons in 3 gallons water. 3 lbs. \$1.60; 50 lbs. \$72.50.

KILGORE'S SLUG DUST. For controlling slugs in nurseries, greenhouses etc. Use 2 lbs. per 1000 sq. ft. 1-lb. canister \$1.00; 6-lb. bag \$3.30; 50-lb. drum \$25.00.

KRYOCIDE (Natural Cryolite). A fluorine spray for chewing insects. Use 3 pounds Kryocide to fifty gallons of water. 1-lb. can 45c; 4-lb. bag \$1.10; 50-lb. bag \$10.65.

KRYOCIDE DUST 30-70 (Code No. K-6)—(Contains 30% Kryocide). A fluorine dust especially recommended for worms. Lb. 15c; 25 lbs. \$2.60; 50 lbs. \$4.20; 100 lbs. \$8.40.

KRYOCIDE-SULPHUR DUST 30-70 (Code No. K-4)—(Contains 30% Kryocide.) Lb. 15c; 25 lbs. \$2.95; 50 lbs. \$4.85; 100 lbs. \$9.70.

LINDANE PREPARATIONS

ISOTOX SPRAY—No. 200 (Contains 20% gamma isomer). Use 1 pt. per 100 gals. water. 4-oz. bottle 95c; 16-oz. bottle \$2.25; 1 gal. \$14.59.

ISOTOX—No. 25-W (Contains 25% gamma isomer). Use 1 lb. per 100 gals. water. For cucumbers, squash, melons, etc. 1-lb. can \$2.10; 4-lb. can \$8.20; 20 lbs. \$40.00.

INSECTICIDES (Continued)

KILGAM DUST—No. 1 (Code No. Q-4)—(Isotox) Contains 1% gamma isomer of Lindane. Lb. 15c; 25 lbs. \$2.60; 50 lbs. \$4.15; 100 lbs. \$8.30.

KILGAM DUST—No. 2. (Code No. Q-5)—(Isotox) Contains 1½% gamma isomer of Lindane. Lb. 15c; 25 lbs. \$3.25; 50 lbs. \$5.55; 100 lbs. \$11.10

KILGAM—SULPHUR DUST No. 6 (Code No. Q-6) (Isotox) Contains 1% gamma isomer of Lindane. Lb. 15c; 25 lbs. \$2.90; 50 lbs. \$4.75; 100 lbs. \$9.50.

MALATHION EMULSION. (Contains 5 lbs. technical malathion per gal.) Use 1 qt. per 100 gals. water. 1 oz. 49c; 5 oz. \$1.39; 1 pt. \$2.69; 1 qt. \$4.29; ½ gal. \$7.95.

MALATHION. 25% Wettable Powders. Use 2 lbs. per 100 gals. water 1 lb. 75c; 5 lbs. \$2.80; 50 lbs. \$24.30.

"MARLATE" 50—A residual insecticide with exceptionally low toxicity to animals and humans. There is only a minimum hazard from the spray residue. Especially recommended for worm control on cucumbers, squash, and melons—will not harm plants. 4-lb. bag \$3.40; 50-lb. bag \$40.00.

"MARLATE" 2-MR—(Emulsifiable)—A residual insecticide for control of leafhoppers and other pasture insects. Also used for chinch bugs, fleas, soil insects in lawns, and insect and fly control of dairy and beef cattle as spray or in "back-rubber". 1 gal. \$4.75; 5 gal. \$22.20; 30 gal. \$124.50.

MARLATE DUST NO. 1 (Code No. W-1) (Contains 3% Methoxychlor). 1 lb. 15c; 25 lb. \$3.00; 50 lbs. \$5.00; 100 lbs. \$10.00.

OIL EMULSION 60. For the control of scales, mealy bugs, red spiders, etc. Dilute with water 1 to 60. Gal. 80c; gals. \$3.25; 55 gals. \$22.00. Deposit on containers.

PARATHION PREPARATIONS. Parathion products are highly toxic to man and warm blooded animals. It is urged that all precautions and recommendations stated on labels be carefully followed.

VAPOPHOS WETTABLE 15%. A wettable powder containing 15% parathion. Use at the rate of 1 lb. to 100 gals. of water. 1 lb. 56c; 3-lb. bag \$1.20; 4-lb. bag \$1.60; 5-lb. bag \$2.00; 100-lb. drum \$40.00.

VAPOPHOS 4 FLOWABLE. 5 gals. \$58.50; 30 gals. \$346.50.

ORTHOPHOS 4 SPRAY. (Contains 42% parathion per gal). Use 1/3 pt. per 100 gallons of water. 1 gal. \$22.00; 5 gals. \$108.75.

KILPHOS DUST NO. 1. (Code No. E-1)—Contains 1% parathion. 50 lbs. \$2.75; 100 lbs. \$5.50.

KILPHOS DUST NO. 2 (Code No. E-2)—Contains 1% parathion and wettable sulphur. 50 lbs. \$3.50; 100 lbs. \$7.00.

KILPHOS DUST NO. 4 (Code No. E-4)—Contains 2% parathion. 50 lbs. \$3.95; 100 lbs. \$7.90.

KILPHOS-KILPHENE-SULPHUR DUST No. 2 (Code No. EI-2)—Containing ¾% parathion and 5% toxaphene. 50 lbs. \$4.30; 100 lbs. \$8.60.

PARIS GREEN. ¼ lb. 30c; 1 lb. 80c; 5 lbs. \$3.45.

POISON BAITS. Ready for use to control crickets, cutworms, grasshoppers and field crickets. Also ideal for airplane application. Do not use poison baits No. 10 and 11 on cucumbers, squash and melons.

POISON BAIT NO. 8 (Code No. B-8)—(Contains chlordane). 1 lb. 10c; 25 lbs. \$1.95; 50 lbs. \$3.45; 100 lbs. \$6.90.

POISON BAIT NO. 10 (Code No. B-10)—(Contains toxaphene). 1 lb. 10c; 25 lbs. \$1.90; 50 lbs. \$3.40; 100 lbs. \$6.80.

POISON BAIT NO. 11 (Code No. B-11)—(Contains chlordane and toxaphene). 1 lb. 15c; 25 lbs. \$2.30; 50 lbs. \$4.15; 100 lbs. \$8.30.

PYROCID DUST NO. 7½ (Code No. P-7)—Recommended for control of worms, flea hoppers, aphids, thrips and etc. on vegetable crops. Can be used with safety up to harvest. 1 lb. 25c; 25 lbs. \$4.80; 50 lbs. \$8.60; 100 lbs. \$17.20.

RED ARROW INSECT SPRAY. Kills garden aphids, melon aphids, worms, leaf hoppers, white flies, red spiders, caterpillars, thrips, celery leaf tiers. Average dilution 1 to 400 parts water. Safe to use on vegetables up to harvest. 1 oz. 35c; 4 oz. \$1.25; pint \$3.30; quart \$6.30; gal. \$20.00.

RHOTHANE. A compound closely related to DDT. Its toxicity to most insects is equal to that of DDT and less toxic to humans and warm blooded animals.

RHOTHANE WP-50. A wettable form of powder, containing 50% of the technical material. Use at the rate of 2 lbs. to 100 gals water. 4-lb. bag \$2.04; 50-lb. bag \$23.50.

RHOTHANE EMULSION CONCENTRATE. (Contains 25% of the technical material.) 1 gal. \$3.00; 5 gals. \$13.50; 30-gal. drum \$75.00; 50-gal. drum \$120.00.

RHOTHANE 5-TOBACCO DUST. (Code No. J-2)—(Contains 5% DDD.) 1 lb. 15c; 25 lbs. \$2.35; 50 lbs. \$3.65; 100 lbs. \$7.30.

ROTENONE DUST, No. 1. (Code No. R-6)—Safe to use up to harvest on vegetable crops for control of worms, aphids, thrips, etc. 1 lb. 15c; 25 lbs. \$3.30; 50 lbs. \$5.55; 100 lbs. \$11.10.

SCALECIDE. Excellent for citrus. Combined with Sulfocide at the rate of two gallons Scalecide and one gallon Sulfocide to two hundred gallons of water. It takes care of many citrus troubles. Pt. 85c; qt. \$1.25; gal. \$3.00; 5 gals. \$11.50.

SYSTOX—A systemic insecticide (which acts from within the plant) effective against aphids and red spider mites. Systox is harmless to beneficial insects. Permitted for use on ornamentals, broccoli, Brussels sprouts, cabbage, cauliflower, muskmelons, potatoes, strawberries and oranges. 1-gal. can \$32.00.

TOXAPHENE PREPARATIONS

ALLTOX 40W. (Contains 40% toxaphene.) Use from 3 to 4 lbs. per 100 gals. of water. 6-lb. bag \$1.80; 50-lb. bag \$13.50.

ALLTOX SPRAY NO. 4.8. (Contains 50% toxaphene.) Use from 1 to 1½ qts. per 100 gals. of water. Gal. \$4.20.

KILPHENE 5-DUST (Code No. 1-3)—(Contains 5% toxaphene.) 1 lb. 10c; 25 lbs. \$1.75; 50 lbs. \$2.45; 100 lbs. \$4.90.

KILPHENE 10-DUST. (Code No. 1-2)—(Contains 10% toxaphene.) Lb. 10c; 25 lbs. \$2.30; 50 lbs. \$3.55; 100 lbs. \$7.10.

KILPHENE 5-SULPHUR DUST. (Code No. I-1)—(Contains 5% toxaphene.) Lb. 10c; 25 lbs. \$2.05; 50 lbs. \$3.10; 100 lbs. \$6.20.

TEPP PREPARATIONS

VAPOTONE XX SPRAY (20%) for control of aphids, thrips, red spiders. Use from ½ to 1 pt. per 100 gals. water. 1 gal. \$12.00; 5 gal. \$58.50.


ORTHO TEPP SPRAY (40%). Use ½ to 2/3 pt. per 100 gals. water per acre. For control aphids, thrips, red spiders and leaf hoppers. 1 gal. \$12.00; 5 gal. \$58.75.

VOLCK OIL SPRAY. A liquid oil spray and spray carrier, furnished in the following sizes only, 4 oz. 35c; 16 oz. 75c; 32 oz. \$1.25; 1 gal. \$3.00.



FUNGICIDES

ANTIBIOTIC SPRAYS



Agri-Mycin and Ortho Streptomycin, two of the Antibiotic materials, are now being used successfully for the control of bacterial spot on tomatoes and peppers, blackleg and soft rot of potatoes, wildfire and blue mold of tobacco. Experimental work is being continued on many other plants. Work on compatibility with insecticides and other fungicides is being conducted. Antibiotics are now being recommended in combination with neutral copper for diseases on celery and pepper. General recommendation is 200 P.P.M. (Two hundred parts per million). Two 9.34 oz. bottles of Agri-Mycin 100

or two 8 oz. bags of 17% Ortho Streptomycin in 100 gallons of water makes a 200 P.P.M. spray. 100 P.P.M. in combination with 4 lbs. of neutral copper is recommended.

Agri-Mycin—9.34 oz. bottle \$6.60; 25-lb. drum (equivalent to 42½ bottles) \$219.00.

Ortho Streptomycin—8 oz. bag of 17% \$6.60; 1-lb. bag \$13.20; 25-lb. drum \$219.00 f.o.b. Plant City or Kilgore stores.

PHYTOMYCIN. A 20% Liquid Streptomycin. 6¼ oz. bottle in 100 gals. of water makes 100 P.P.M. spray. 6¼ oz. bottle \$6.50.

ARASAN. (50% Thiram). An effective treatment for vegetable seed, reduces loss from seed decay and damping-off. 8 oz. \$1.00; 2 lbs. \$3.20; 10-lb. bag \$10.35.

ARASAN-75. (75% Thiram). An effective treatment for vegetable seed. Reduces loss from seed decay and damping-off. 1 oz. pkg. 50c; 8-oz. can \$1.50; 25-lb. drum \$40.50; 100-lb. drum \$155.00.

CERESAN (New Improved). Destroys seed-borne diseases by direct contact with the spores or penetrating vapor fumes. Ceresan has been used successfully in treating Gladiolus bulbs for control of Fusarium Rot before planting. Use ½ lb. Ceresan, 5 tablespoons of DuPont Spreader-Sticker to 25 gallons water. Dip bulbs in solution, let remain 15 minutes. Bulbs can be planted immediately after treatment. 1 lb. \$1.75; 100 lbs. \$110.00.

COPPER COMPOUNDS

BLUESTONE (copper sulphate) Granular or Snow-form. 1 to 10 lbs. 25c per lb; 25 lbs. \$5.45; 50 lbs. \$10.40; 100 lbs. \$21.25.

COPPER-A COMPOUND PREPARATIONS

COPPER-A COMPOUND—An excellent Copper fungicide for spraying purposes containing 45% metallic copper. Formulated especially for greater safety on copper sensitive plants. Use 4 lbs. per 100 gals. of water. 6-lb. bag \$3.12; 50-lb. bag \$24.95.

KILCOP DUST NO. 1 (Code No. A-1)—(5.4% metallic copper.) Lb. 15c; 25 lbs. \$2.85; 50 lbs. \$4.70; 100 lbs. \$9.40.

KILCOP DUST NO. 2 (Code No. A-2)—(6.75% metallic copper.) Lb. 15c; 25 lbs. \$3.15; 50 lbs. \$5.35; 100 lbs. \$10.70.

KILCOP-SULPHUR DUST NO. 3 (Code No. A-3)—(6.75% metallic copper.) Lb. 15c; 25 lbs. \$3.55; 50 lbs. \$6.10; 100 lbs. \$12.20.

CUPROCID-YELLOW. (Contains 80% metallic copper.) A cuprous oxide for the control of various truck crop diseases. Use 2 lbs. per 100 gals. of water. 3-lb. bag \$3.30; 100-lb. drum \$105.00.

KILCOP "53" PREPARATIONS

KILCOP "53." (Contains 53% metallic copper.) Kilcop is extremely fine, which is essential for good coverage and effective control of fungus. Use 4 lbs. per 100 gals. water. 1-lb. can 95c; 2-lb. bag \$1.00; 6-lb. bag \$2.70; 50-lb. bag \$21.50.

KILCOP DUST NO. 5 (Code No. O-2)—(7.4% metallic copper.) Lb. 15c; 25 lbs. \$2.70; 50 lbs. \$4.40; 100 lbs. \$8.80.

KILCOP DUST NO. 6 (Code No. O-3)—(9% metallic copper.) Lb. 15c; 25 lbs. \$2.95; 50 lbs. \$4.90; 100 lbs. \$9.80.



DITHANE PREPARATIONS

DITHANE D-14 (Nabam). This fungicide has given excellent control on a variety of crop diseases such as late and early blight of potatoes and tomatoes, leaf blights of peppers, curvularia leaf spot of glads, anthracnose of beans and downy mildew of cucumbers, squash and melons and helminthosporium leaf blight of corn. Use 2 qts. Dithane D-14, ¾ lb. zinc sulphate to 100 gals. water. 1 gal. \$2.10; 5 gals. \$8.75; 30 gals. \$49.50.

DITHANE Z-78 (Zineb) (Wettable). Effective in the control of downy mildew and anthracnose of cucumbers, squash, melons and blue mold on tobacco. Helminthosporium leaf blight of corn). Use 2 lbs. to 100 gals. 1-lb. canister \$1.50; 3-lb. bag \$2.55; 50-lb. bag \$41.00.

DITHANE M-22 (Maneb). Controls a number of diseases on tomatoes, celery, peppers, eggplant and potatoes. Particularly effective where manganese deficiency exists. Use 1½ lbs. per 100 gals. water. 3-lb. bag \$3.36; 4½-lb. bag \$5.04; 25-lb. drum \$27.50.

ZINEB DUST NO. 1 (Code No. M-1)—(Contains 4% zineb). Lb. 15c; 25 lbs. \$2.45; 50 lbs. \$3.90; 100 lbs. \$7.80.

ZINEB DUST NO. 2 (Code No. M-2)—(Contains 6% Zineb). Lb. 15c; 25 lbs. \$3.05; 50 lbs. \$5.05; 100 lbs. \$10.10.

ZINEB-SULPHUR DUST NO. 4 (Code No. M-5)—(Contains 4% zineb). Lb. 15c; 25 lbs. \$2.85; 50 lbs. \$4.70; 100 lbs. \$9.40.

"FERMATE" (R) (Ferbam)—Fungicide for control of tobacco blue mold, early and late blight of celery; downy mildew of cabbage, damping-off in celery and lettuce seedbeds. Use 2 lbs. per 100 gals. water. 3-lb. bag \$1.89; 50-lb. bag \$30.00.

KARATHANE: A specific fungicide for the control of powdery mildew on cantaloupes, cucumbers, squash and ornamentals.

Use ½ lb. to ¾ lb. per 100 gals. water.
Price—3-lb. bag \$5.10.

KARATHANE DUST. 1% (Code No. Z-1). 1 lb. 15c; 25 lbs. \$3.00; 50 lbs. \$4.30; 100 lbs. \$8.60.

"MANZATE" (R)—(Maneb)—Fungicide—Specific for control of tomato diseases such as late blight and gray leaf spot. Provides protection without yellowing or stunting of plants.

Also controls major foliage diseases of gladioli, cucumbers, cantaloupes, potatoes, celery, and other crops. Use 1½ lbs. per 100 gallons of water. 3-lb. bag \$3.36; 4½-lb. bag \$5.04; 25-lb. drum \$27.50.

ORTHOCIDE 50-W (Captan). For control of damping-off in pepper seedbeds and other fungus diseases. For Irish potato seed treatment see page 52. Use 4 lbs. per 100 gals. water. 4-lb. bag \$3.00; 50-lb. bag \$34.00.

ORTHOCIDE 75 SEED PROTECTANT. Contains 75% captan. A seed treatment to prevent seed rot and damp-off diseases. 4½ oz. shaker can \$1.19; 1-lb. can \$2.60; 4-lb. can \$10.20; 20-lb. drum \$33.00.

FUNGICIDES (Continued)

"PARZATE" (R)—Zineb—Fungicide (Wettable)—Effective in the control of downy mildew, and anthracnose of cucumbers, cantaloupes, squash, and watermelons.

For control of helminthosporium leaf blight of corn, early and late blight of potatoes, blights and gray leaf spot of tomatoes, blue mold of tobacco, and leaf spots of many other crops. Use 2 lbs. per 100 gals. of water. 3-lb. bag \$2.55; 50-lb. bag \$41.00.

"PARZATE" (R)—Nabam—Fungicide—Used as an effective fungicide wherever nabam or zineb is recommended. Especially useful for potato blights and helminthosporium leaf blight of corn. Use 2 qts. "Parzate" and $\frac{3}{4}$ lb. zinc sulphate per 100 gals. 5-gal. can \$8.75; 30-gal. drum \$49.50.

PHYGON-XL (Wettable). A fungicide for the control of leaf blights of tomatoes, beans and various other plant diseases. Use $\frac{1}{2}$ to 1 lb. per 100 gals. of water. 2 oz. \$1.00; 1 lb. \$3.00; 4 lbs. \$6.60; 100 lbs. \$158.00.

SEMESAN. An effective treatment for vegetable and flower seeds to control damping-off fungi in the soil. Also repels ants. Directions on package. 4-oz. pkg. \$1.80; 12 ozs. \$4.95; 25 lbs. \$105.00.

SEMESAN BEL. For treatment of potatoes, controlling certain diseases of this crop. Directions on package. 3 oz. \$1.05; 1 lb. \$3.30; 4 lbs. \$10.65; 40 lbs. \$99.80.

SPERGON. Very valuable for treating seeds, especially beans and peas, to control soil diseases for better stands and increased yields. Use 2 ounces per bushel of seed with beans and peas. 1 oz. 50c; 5 ozs. \$1.75; 1 lb. \$2.50; 10 lbs. \$20.00; 50 lbs. \$95.00.

SPERGON (Wettable). Valuable for spraying to control diseases—especially downy mildew in cabbage seedbeds. Use at the rate of 2 lbs. to 100 gals. of water 1 lb. \$2.25; 10 lbs. \$11.40; 50 lbs. \$48.50.

SPERGON DUST NO. 3. (Code No. Y-3)—Contains 5% Spergon—Wettable. For the control of downy mildew on cabbage. Lb. 15c; 25 lbs. \$3.15; 50 lbs. \$5.25; 100 lbs. \$10.50.

SULFOCID. A vegetable spray which is also valuable for citrus fruits. Checks mildew, rust and anthracnose on beans. Also used in washing water before shipping beans, tomatoes, eggplants and peppers to check rot and other diseases while in transit. Use 2 qts. to 100 gals. of water. $\frac{1}{2}$ pt. 65c; pt. \$1.00; qt. \$1.50; gal. \$3.00; 5 gals. \$11.50.


SULPHUR (Dusting). Lb. 5c; 25 lbs. \$1.00; 50 lbs. \$1.58; 100 lbs. \$3.15; ton \$62.00.

SULPHUR (Wettable). 50 lbs. \$1.70; 100 lbs. \$3.40; ton \$67.00.

TERSAN (R) 75—A lawn and turf fungicide for control of "brown-patch" and "dollar spot." Use two tablespoons full to $1\frac{1}{2}$ gal. water to 60 sq. ft. 8-oz. cans \$1.65; 3-lb. bag \$5.50.

ZERLATE (R) Ziram—Fungicide—Specific for control of "frog-eye" spot on peppers and leaf blights on celery. Use two pounds per 100 gals. 3-lb. bag \$1.95; 50-lb. bag \$31.00.

Insecticide and Fungicide Combinations



Prices quoted are f.o.b. Plant City
or Kilgore stores

Prices subject to change without notice

The Kilgore Seed Company processes many combinations of insecticides and fungicides which are too numerous to list in our General Catalog. If you do not find the combination you want, write our General Office in Plant City or call at your nearest Kilgore Store and it is likely we will have a dust in stock which will fill your requirements.

KILCIDE-KILCOP DUST NO. 1 (Code No. DA-1)—(Contains 3% DDT plus 5.4% metallic copper.) Use where DDT is needed with a neutral copper. Lb. 15c; 25 lbs. \$3.20; 50 lbs. \$5.40; 100 lbs. \$10.80.

KILCIDE-KILCOP - MANGANESE - SULPHUR DUST NO. 1 (Code DAX-1)—(Containing 5% DDT, 5.4% metallic copper, manganese sulphate and sulphur) a combination dust for control of insects, disease and nutritional deficiencies. 1 lb. 20c; 25 lbs. \$3.80; 50 lbs. \$6.60; 100 lbs. \$13.20.

KILCIDE-ZINEB DUST NO. 1 (Code No. DM-1)—(Contains 5% DDT and 4% Zineb.) For use where DDT is desirable for general insect control with Zineb. 1 lb. 15c; 25 lbs. \$3.05; 50 lbs. \$5.05; 100 lbs. \$10.10.

KILGAM-KILCOP DUST NO. 1 (Code No. QA-3)—(Contains 1% lindane [essentially pure gamma isomers of benzene hexachloride] and 5.4% metallic copper.) An excellent combination designed for use on cucumbers, squash, and melons which cannot tolerate ordinary BHC compounds (Kilhex), as well as for cabbage, celery, tomatoes, peppers and eggplants, at periods when dusts containing ordinary BHC (Kilhex) cannot be

used because of the possible contamination of edible portions due to odor or flavor. 1 lb. 20c; 25 lbs. \$4.30; 50 lbs. \$7.65; 100 lbs. \$15.30.

KILHEX—KILCOP DUST NO. 1 (Code No. GA-1)—(Contains 1% gamma isomer and 6.75% metallic copper.) A practical insect and disease control combination at a minimum cost for cabbage, celery, tomatoes, peppers and eggplants for application before fruits or edible portions are well developed. 1 lb. 20c; 25 lbs. \$3.75; 50 lbs. \$6.45; 100 lbs. \$12.90.

KILPHENE-KILCOP DUST NO. 2. (Code No. IA-2)—A convenient insect and disease control combination containing 5% toxaphene and 6.75% metallic copper. (Under no circumstances use this material on cucumbers, squash, or melons.) 1 lb. 20c; 25 lbs. \$3.80; 50 lbs. \$6.60; 100 lbs. \$13.20.

KILPHOS-ZINEB DUST NO. 1. (Code No. EM-1)—(Contains 1% parathion and 6% Zineb.) Can be used on most crops for most insect and disease control. 50 lbs. \$6.60; 100 lbs. \$13.20.

KRYOCIDE-KILCOP DUST NO. 3. (Code No. KA-1)—(Contains 30% Kryocide plus 5.4% metallic copper.) A good insecticide and fungicide combination for most vegetable crops. Lb. 20c; 25 lbs. \$4.40; 50 lbs. \$7.75; 100 lbs. \$15.50.

ZINEB KILGAM DUST NO. 1 (Code No. MQ-1)—(Contains 1% lindane and 4% Zineb.) This combination is recommended for purposes where Zineb has been proved more effective than copper. Refer to Kilgarn-Kilcop Dust No. 1 for further important details. 1 lb. 20c; 25 lbs. \$3.95; 50 lbs. \$6.90; 100 lbs. \$13.80.

ZINEB-KRYOCIDE DUST. (Code No. MK-1)—(Contains 4% Zineb and 30% Kryocide.) This combination is recommended for purposes where Zineb has been proved more effective than copper. Lb. 20c; 25 lbs. \$4.00; 50 lbs. \$6.95; 100 lbs. \$13.90.

HOUSEHOLD AND GARDEN PESTICIDES

KILGORE'S NEMA-KIL. (Contains Nemagon). A new nematocide, which kills nematodes, including rootknot, sting nematodes, burrowing nematodes, stubby-root nematodes, etc. Can be used safely on many established plants, such as hibiscus, roses, gardenias, azaleas, ixoras, as well as lawn grasses and most flowers without harm to plants or grasses. NEMA-KIL is emulsified in water and easy to apply. It is the answer to a long-felt problem as to how to get rid of nematodes in your yard without digging up lawns, shrubbery, etc. One quart of NEMA-KIL will treat from 600 to 800 square feet of surface.



Price: pt. \$1.95; qt. \$3.30; 1 gal. \$10.00; 5 gal. \$47.50; 30-gal. drum \$282.00

IMPROVED ISOTOX GARDEN SPRAY-M—An effective spray containing 5% lindane, for control of insects on roses, camellias, azaleas, ixoras, hibiscus, bougainvillea, and other ornamentals, house and garden plants. This spray is marked by the absence of odor and visible residue. Use four teaspoons per gallon of water. Price: 4-oz. bottle \$1.00; 8-oz. \$1.85; 16-oz. \$2.95.

KILCHLOR 40-W. A chlordane concentrate for the control of chinch bugs, army worms, sod webworms and ants by spraying. 1-lb. can 90c; 5-lb. bag \$3.35; 10-lb. bag \$6.50; 25-lb. bag \$13.25; 50-lb. bag \$25.50.

KILCOP "45"—(Contains 45% metallic copper.) Kilcop is extremely fine which is essential for good coverage and effective control of fungi. Consult label for dosages. 1-lb. can 90c.

KILGORE'S C-M-Z GARDEN SPRAY.—A combined nutritional and fungicidal spray for shrubs, fruit trees and vegetables. Use two tablespoons per 1 gallon of water on Avocados, Mangos and shrubs, 1¼ tablespoons on citrus and 2½ tablespoons on vegetables. No wetting agent required. 1 lb. can 75c.

KILGORE'S GENERAL PURPOSE DUST. (Code No. P-6.) Contains pyrethrins. A non-poisonous dust, which can be used in or around the house to kill ants, roaches, flies and many garden insects. The ideal dust for use on garden vegetables up to picking time, Dust Gun is refillable and can be used over provided the gun is kept in a dry place. Dust gun complete with dust 75c; refills for dust gun (dust only), 1 lb. 30c.



DRY FLY BAITS.—For use against house flies and their maggots in out-buildings, kennels, around garbage cans, etc. Easy to apply—scatter by hand. No spraying equipment needed. Kills within a few minutes. Lasts for several days when applied to dry ground. Do not use inside of dwellings.

FLY FLAKES—Apply 4 oz. per 100 sq. ft. 1-lb. shaker can 98c; 2-lb. bag 89c; 5-lb. bag \$1.79; 10 lbs. \$3.25; 25-lb. drum \$6.00.

ORTHO FLY KILLER (Dry Bait)—Apply 4 oz. to 500 sq. ft. 1 lb. can 98c; 5-lb. bag \$1.98.

KILGORE'S KIL-CHINCH. Granulated. (Code No. V-2.) A recently developed material for satisfactory chinch bug control in St. Augustine type of grasses. Furnished in a granular form, similar to coarse sea sand. The toxic ingredient is malathion. Apply at the rate of two pounds per 100 sq. ft. 5-lb. bag \$1.50; 10-lb. bag \$2.50; 50-lb. bag \$10.00; 100 lbs. \$20.00.

KILGORE'S FLEA-KILL (Code No. G-10.) Apply dust under houses and other buildings, around kennels and on bare sandy areas at the rate of 2 lbs. per 100 sq. ft. Dust lawns at the rate of 2 lbs. per 250 sq. ft. Be sure grass is dry before applying dust. 5 lbs. 90c; 10 lbs. \$1.50; 25 lbs. \$2.75; 50 lbs. \$4.50; 100 lbs. \$9.00.

KILGORE'S SLUG DUST—A 15% Methaldehyde dust providing effective control of slugs in nurseries, green-houses, etc. Use at the rate of 2 lbs. per 1000 square feet. Usually three applications are required.

1 lb. Canister \$1.00; 6-lb. bag \$3.30; 25-lb. Drum \$15.00
50-lb. Drum \$25.00

KILGORE'S KILCHLOR. A residual household spray. Contains 2% chlordane plus isobornyl thiocyanate, for control of roaches, ants, and most other household insects. Qt. 65c; gal. \$2.15.

KILGORE'S KILSPRAY. An excellent indoor space spray for flies, mosquitoes, roaches, ants, fleas and other household insects. Not a residual spray. It should be used when insects are present. Qt. 65c; gal. \$2.15.

KILSPRAY INSECT BOMBS—Kills flies, mosquitoes, gnats, flying moths, roaches, ants and Silverfish.

Kills quickly. Easy to use.

Price 12 oz. Bomb. 97c prepaid

KILSPRAY—Air Freshener, Bombs—Stops all offensive odors. Quick and easy to use. Non-toxic, non-inflammable. Comes in two odors—Bouquet and Pine needle.

Price 10 oz. Bomb. 97c each



MALATHION Emulsion 50%. For control of scale and many other lawn and garden insects. Use 2 tablespoons per gal. of water. 1 oz. 49c; 5 oz. \$1.39; pt. \$2.98; qt. \$5.29.

KILGORE'S 25% DDT EMULSION CONCENTRATE. For the control of insects on truck crops, shrubs, etc. Qt. \$1.35; gal. \$3.25

KILCHLOR-KILPHENE POISON BAIT NO. 11. (Code No. B-11)—(Contains chlordane and toxaphene). For control of mole crickets, cutworms, and many soil insects. 1 lb. 15c; 25 lbs. \$2.30; 50 lbs. \$4.15; 100 lbs. \$8.30.

ORTHO-KLOR "44" SPRAY.—For control of ants, mole crickets, cutworms, army worms, roaches and many other pests. For Termite Control see page 58. 4 oz. 70c; 8 oz. \$1.20; 16 oz. \$1.98; 32 oz. \$2.98.

ORTHOL GARDEN SPRAY. An oil emulsion containing nicotine and DDT for the combined control of aphids, red spiders, mealy bugs, thrips and many other insects. Use at the rate of 6 tablespoonfuls to 1 gal. of water. 4 oz. 35c; 16 oz. 75c; 32 oz. \$1.25; 1 gal. \$2.85.

RED ARROW INSECT SPRAY. Kills aphids, worms, leaf-hoppers, white flies, red spiders, caterpillars, thrips, celery leaf tier. Average dilution 1 to 400 parts water. 4 oz. \$1.25; pint \$3.30; quart \$6.30; gal. \$20.00.

KILGORE'S ROSE DUST. (Code No. PO-4.) A combined insecticide and fungicide, to keep rose bushes free of insects and diseases, by applying at intervals of seven days. An excellent material as a general insecticide and fungicide for garden vegetables. Dust Gun, complete with dust 75c. Refills for dust gun (dust only) 1 lb. 35c.

SABA CHINCH. (Code No. S-2.) For control of chinch bugs, army worms, sod webworms, and ants. Use 4 lbs. per 100 sq. ft. Lb. 15c; 25 lbs. \$3.10; 50 lbs. \$5.15; 100 lbs. \$10.30.

SEEDTONE. Seed Protectant. (Containing 50% captan)—A guard against seed decay, damping-off and seedling blight. ¼ oz. pkt. 25c.

VOLCK OIL SPRAY. 4 oz. 35c; 16 oz. 75c; 32 oz. \$1.25; 1 gal. \$3.00.


VOLCK ISOTOX SPRAY. For use on ornamental plants against red spiders, certain scale insects, mealy bugs, white flies and certain other garden pests. Use 5 tablespoonfuls per 1 gal. water. Don't use more than three applications per season. 4 oz. 40c; 8 oz. 60c; 16 oz. 95c; 32 oz. \$1.75.



FUMIGANTS

CARBON DI-SULPHIDE (Hi-life.) Recommended for ridding grain of weevils, and killing ants, etc., and for destroying vermin. Sold in gallon can only. Per gallon \$2.25; 5 gal. \$10.00. (Not mailable.)

PYRENONE, GRAIN PROTECTANT. A contact material for the protection of corn and other grains stored in open cribs or bins. Offers freedom from toxic hazards to man and animals. Effective against weevils and other storage insects. Prices: 50-lb. bag \$12.50, f.o.b. Plant City.



CYANO GAS ANT-KILLER. Cyanogas Ant-Killer kills ants with the least trouble. It is a gas-producing powder—flows freely through the spout on the can. Not a bait. It's the gas that kills 'em. 4-oz. can 50c. (Not mailable.)

CYANO GAS A-DUST. For killing the following pests: ants (Garden, Mound Building; Leaf Cutting). Rats (in burrows; along ditch banks; in city dumps; under pavements and tight floors; in lumber piles; on poultry farms; in grain elevators and stockyards). Mice (field mice and pine mice), moles, gophers. 1-lb. tin \$1.25; 5-lb. can \$4.25; 25-lb. can \$13.75. (Not mailable.)

CYANO GAS G-FUMIGANT. Comparable to sea sand in size of particles. Used for fumigating greenhouses, warehouses, buildings and for grain fumigation. 4-oz. can 50c; 5-lb. can \$4.25; 25-lb. can \$13.75; 100-lb. drum \$43.35. (Not mailable.)

KILGORE'S NEMA-KIL (Contains Nemegon). A new nematocide which kills nematodes in the soil. Can be used around many established plants. See page 63 for more detailed information. Pt. \$1.95; qt. \$3.30; gal. \$10.00; 5 gal. \$47.50; 50 gal. \$282.00.

V-C 13 NEMACIDE. For protecting growing trees, shrubs and lawn turfs from nematodes. Use 1 pt. in 50 gal. water to 180 sq. ft for turfs and lawns. Do not use on anything grown for human or animal consumption. 8 oz. bottle \$2.50; 1 qt. \$8.50; 1 gal. \$28.50; 50 gal. \$115.00.

VAPAM-SOIL FUMIGANT. For control of certain weeds, soil-barn fungus diseases, nematodes and garden centipedes, etc. Consult label for detailed information. 1 gal. \$5.60; 5 gal. drum \$24.25; 30 gal. drum \$120.00.

METHYL BROMIDE (with 2% chloropicrin). For treatment of nursery seedbeds and greenhouse soil. 1-lb. can 85c.

JIFFY APPLICATORS. For 1-lb. methyl bromide cans \$3.35.

Note—For plastic covers used in treating seedbeds with methyl bromide. Contact Kilgore Seed Co., Plant City or your nearest Kilgore store.

SOIL FUMIGATION FOR CONTROL OF SOIL-BORNE PESTS. Nematodes are present in most of our soils, and seriously affect most of our vegetable crops and reduce yields. Nematodes can be controlled by the proper use of soil fumigants, either solid treatment, which is applied with power applicator about 12 inches apart, or row treatment, which is only applied in the row that is to be planted. This method has become very popular, especially with small growers, as cheap handmade applicators with gravity flow, can be used attached on a tractor or a horse drawn plow. The cost of this method of treatment is only about one-third the cost of solid treatment, and has proved very satisfactory. "D-D" and "Soilfume" are the two materials most extensively used in commercial treatment at this time. Information on soil fumigation can be obtained from any of the Kilgore stores or write The Kilgore Seed Company, Plant City, Florida.

D-D Soil Fumigant. 1 gal. will treat approximately 1500 square feet of soil area applied in 15 inch squares, 10 inches deep, with hand applicator. 22 gals. will treat approximately one acre applied with a machine in 12 inch rows. Applied only in the row that is to be planted. 1 pt. will treat 260 feet of row, or approximately 5½ gals. per acre in 4 foot rows. Gallon \$2.00; 30-gal. drum \$46.50; 55-gal. drum \$82.50.

SOILFUME 80-20 (ethylene dibromide). 1 gal. will treat approximately 1400 square feet of soil area applied in 12-inch squares, 10 inches deep with hand applicator. 30 gals. in 12-inch rows applied with a machine will cover one acre. Applied only in the row that is to be planted, 1 pt. will treat approximately 140 ft. of row or 10 gals. per acre in four foot rows. 1 gal. \$1.65; 30-gal. drums \$46.50; 50-gal. drums \$75.00.

RODENTICIDES AND VERMICIDES

CARBOLINEUM. A wood preservative and vermicide. Prevents damage by termites. Kills and protects against chicken mites and fowl ticks. ½-gal. can \$1.65; 1 gal. \$2.50; 5 gals. \$12.25; 35 gals. \$64.75; 55 gals. \$96.25, f.o.b. Plant City.

NO NIB'L RABBIT REPELLENT is recommended as a rabbit repellent for flowers, shrubs and other ornamentals. When used as directed, will not harm plants or soil and is non-injurious to birds or animals. Easy to use and is effective either as a dust or as a spray. 6-oz. shaker-top can 98c.

DIXIE KILLER. Exterminates rats, mice, roaches, waterbugs, land crabs, cutworms and other destroyers of food and property. 2-oz. jar 35c; 1 lb. \$1.10; 2 lbs. \$2.00. Prepaid.

MAGIKIL JELLY ANT BAIT. Kills ants and is harmless to plants. Ant trap 25c; Garden size 40c; Estate tube \$1.00; 7½ oz. Jumbo size \$1.95; 10 lbs. \$23.00. Prepaid.

TERRO ANT KILLER. Will rid your place of ants quickly. 1¼-oz. bottle 35c; 3-oz. bottle 50c. Prepaid.

HARRIS ROACH TABLETS. 2-oz. pkg. 30c; 4-oz. pkg. 60c.

GATOR ROACH HIVES are sanitary, open-end fibre tubes, containing a specially made gum that roaches, silverfish, waterbugs and crickets eat with relish but results in their certain death. Small size (3 baits) 20c; Large size (3 baits) 40c. Prepaid.

COLONIAL FLEA and TICK KILLER. For cats and dogs. For dogs 4½-oz. shaker-top can 89c; 1-lb. can \$1.50; for dip 8-oz. bottle \$1.50. For cats 4½-oz. can 89c.

COLONIAL "333" RAT AND MOUSE BAIT (Pivalyn Water Bait) 1½ oz. pkg. Makes 1 qt. water bait. 1½ oz. pkg. 45c.

COLONIAL-42.—An effective bait containing Warfarin for the control of all species of rats and mice. 4-oz. pkg. 35c; 10 oz. 60c; 1 lb. 90c; 2 lbs. \$1.65.

KILGORE'S RAT AND MOUSE BAIT. A Warfarin bait especially designed for the control of rats and mice in fields and groves. 10 lbs. \$3.50; 25 lbs. \$8.00; 50 lbs. \$15.00.

QUICK DEATH RAT KILLER. A fortified red squill formula. Safe to use in homes. Price, 25c per package.

Spreaders, Stickers, Emulsifiers, Etc.

DU PONT SPREADER-STICKER—A liquid wetting, spreading, sticking, deposit building agent that increases the effectiveness of sprays on crops, livestock and weeds. Use with soluble fertilizers and nutrient sprays to insure uniform deposit and decrease chances of spot burn. Spreader-Sticker is compatible with hard waters. Use at 1 to 4 oz. per 100 gallons. Oz. 49c; pt. \$1.45; gal. \$8.45; 5 gals. \$41.25.

TRITON B-1956. An efficient emulsifier and spreader for insecticides and fungicides. A combined spreader,

sticker and deposit builder. Soluble in water and oil, it is widely used as an emulsifier in fruit, truck crop and ornamental sprays for maximum spreading and sticking qualities. Use at the rate of 2 to 4 ounces per 100 gallons. Gal. \$7.70; 5 gals. \$37.75; 50 gals. \$370.00.

FISH OIL SOAP. Increases the efficiency of Black Leaf 40 and many other spray materials. Use 1½ to 2 lbs. to 50 gallons as a spreader and sticker. 1 lb. 40c; 1-gal. can \$2.35; 5-gal. can \$8.35.

WEED KILLERS

Note: Read labels carefully—they are put on the package for your protection. Use Weed Killers wisely and you will be using them safely. Be sure to clean spray equipment thoroughly according to approved methods after using weed control materials of any kind. f.o.b. Plant City.

"AMMATE" (R)-WEED AND BRUSH KILLERS—For complete kill of undesirable brush, trees, poison oak, poison ivy, and many other types of weeds. "Ammate" provides the safest way to control brush, trees and weeds that may adjoin cropland. There are no vapors to drift onto sensitive crops or foliage. "Ammate" is non-volatile, non-inflammable and can be used without hazard to humans or animals. Use 60-75 lbs. per 100 gals. plus Spreader-Sticker at 4 oz. per 100 Gal.



2 oz. bottle \$1.15; 6 oz. bottle \$2.95; 50-lb. drum \$15.25; 75-lb. bag \$19.95.

"KARMEX" (R) W—WEED KILLER (Formerly DuPont CMU)—Kills weeds and grasses and prevents regrowth for a prolonged period of time.

Use at ½ lbs./acre for control of annual weed seedlings in sugar cane, pineapple, gladioli and other bulb ornamentals.

Use at 1 lb./1000 sq. ft. wherever it is desired to kill all vegetation for a prolonged period of time (9-12 mos.)—ditches, around storage yards, barns, packing houses, fence rows, etc.

"Karmex" W is non-volatile, non-toxic, non-straining, non-flammable, and non-corrosive.

Do not apply on or near valuable trees or other plants or on areas into which their roots may extend. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Do not apply to locations from which "Karmex" W may be washed or moved into contact with the roots of valuable plants.

2-lb. can \$9.10; 10-lb. drum \$40.50; 50-lb. drum \$187.50.

POLYBOR-CHLORATE. A non-selective herbicide for the prolonged control of weeds about brick walks and patios, fence lines, driveways, or tennis courts, around stores, industrial properties, etc. It is non-selective in its action and will kill any plant with which it comes in contact, consequently cautions, as per label instructions should be observed in order to protect valuable shrubs and trees adjacent to the areas to be treated. Dissolved at the rate of four pounds in one or two gallons of water to be applied to a 100 sq. ft. area. Price 5-lb. pkg. \$2.25; 50-lb. bag \$10.25.

WEED-B-GON—64X. An amine salt of 2, 4-D. Excellent for killing water hyacinth and most broad leaf plants. Apply as per directions on container. Caution—do not use on or let material drift on plants you do not want to kill. Price, 1 gal. \$4.43; 5 gals. \$21.00; 30 gals. \$119.70; 50 gals. \$195.50.

PENITE—6. A highly concentrated, black opaque viscous Sodium Arsenite solution. A non-selective weed killer used for killing weeds on ditch banks, fence rows, etc. Is especially recommended for killing off potato tops before harvest. Permits earlier harvest, toughens the skin and decreases mechanical injury. Use 1 gallon of Penite 6 to 100 gallons of water, applied at the rate of 100 to 150 gallons per acre. Will give excellent potato vine kill. Comply with cautions on label. 5 gal. \$13.75; 30-gal. drum \$63.00.

WEED KILLER (90% sodium TCA). A grass killer for non-selective control in areas not planted to crops. For maximum control of Para grass along ditch banks it is suggested to first use a "pre-weakening" method, such as treating with a relatively cheap contact herbicide, or by flame singeing. TCA can then be applied at fairly low dosages. As an alternate, several applications of TCA in small amounts will produce better results than one application at a higher dosage. West Palm Beach, Fla. 10 lbs. \$5.20; 50 lbs. \$22.50; 100 lbs. \$44.50.

TRIOX. An arsenical weed killer. Toxic to the soil and kills vegetation. Prevents plant growth for one to two years. A liquid, easy to apply with watering can. Do not use on lawns. Consult label. Qt. \$1.00; gal. \$3.50; 5 gals. \$13.50; 30-gal. drum \$72.00.

WEEDONE BRUSH KILLER—32. Contains 2,4,5-T, plus 2,4-D. Kills poison ivy, poison oak stumps and other woody plants. Use 1 to 1½ gals. to 100 gals. water or ½ to 1 pt. to 5 gals. water. Caution—do not use this material on plants you do not want to kill. Price qt. \$3.75; gal. \$7.25; 5 gals. \$34.50.



NEW IMPROVED WEEDONE. A selective weed killer containing 2,4-D plus 2,4,5-T. For complete kill, all the foliage must be sprayed as there is little cross-movement within the stems. Weedone is effective only when the leaves are green and actively growing, for the chemical does not move through dormant or dead tissue. Do not use Weedone on St. Augustine type grasses. 8-oz. bottle \$1.00; 1-qt. bottle \$2.75; 1-gal. bottle \$6.75; 5 gals. \$24.75.

A number of pre-emergence and post-emergence weed control materials are now being offered to truck crop growers. As most of these materials have not been tried sufficiently under Florida conditions we hesitate to make specific recommendations in this catalog. We suggest that growers interested should consult local authorities.

WE WILL MAKE EVERY EFFORT TO SUPPLY WEED CONTROL MATERIALS OF THIS NATURE RECOMMENDED BY THE FLORIDA EXPERIMENT STATIONS FOR EACH AREA.

LIVESTOCK PREPARATIONS

Prices are subject to change without notice and are f.o.b. Plant City unless otherwise specified.

DDT 50% Wettable. For spray on range cattle. To control horn flies, lice and mosquitoes use 20 lbs. per 100 gals. water. Do not use on dairy cattle. 1-lb. canister 70c; 3-lb. bag 99c; 50-lb. bag \$15.00.

Gamtox Wettable No. 10 (10% Gamma isomer of BHC). For spray on range cattle to control horn flies, lice and mosquitoes use 3¾ lbs. per 100 gal. water. Do not use on dairy cattle. 5-lb. bag \$1.60; 50 lbs. \$14.50.

Marlate Methoxychlor (50% Wettable). For spray on dairy and range cattle to control flies and lice. For lice use 25 lbs. per 100 gal. water. For flies use 8¾ lbs. per 100 gal. water. 4-lb. bag \$3.40; 50-lb. bag \$40.00.



DuPont Dairy Cattle Spray. (Methoxychlor, 50% wettable). For control of flies on dairy cattle use 1 lb. in 12 gals. water. 1-lb. can \$1.25; 1-lb. bag \$1.15.

DuPont Livestock Spray and Dip No. 30. For control of horn flies, lice and mosquitoes on range cattle. Do not use on dairy cattle. 2-lb. bag \$1.35; 11-lb. bag \$5.75; 55-lb. bag \$25.50.

Isotox Dairy Spray. (Contains lindane). Spray for killing flies in dairy barns and milk sheds. Do not spray on dairy cows. 1 lb. \$2.19; 4 lbs. \$8.60.



ORTHO Healthy Herd. A wettable powder. (Contains 45% DDT and 2.5% gamma isomer of BHC and 12.5% other isomers of BHC.) Use as spray or dip on cattle, horses, hogs, sheep, and goats to control lice, ticks, horn flies and mosquitoes. Use 10 lbs. per 100 gals. water. (Not for use on dairy cattle or in dairy barns.) 5-lb. bag \$3.25; 50-lb. bag \$25.50.

ORTHO KLEEN Stock. An emulsifiable liquid containing toxaphene and lindane for the control of ticks, lice, flies, etc. on beef cattle and hogs. One pt. makes 4 to 12½ gals. finished spray depending on use. Directions on container. (Not for use in dairy barns or on dairy cattle.) Price 16-oz. bottle \$1.40; 32-oz. bottle \$2.40; 1 gal. \$6.95; 5 gals. \$27.50.

DRY FLY BAITs

For use against flies and their maggots in dairy and livestock barns, stables, poultry houses, feed rooms, kennels, manure piles, etc. Do not apply where poultry can feed upon the poison.


Fly Flakes. For adult flies apply 4 oz. per 100 sq. ft. For maggots, use 1½ lbs. per 100 sq. ft. 1-lb. shaker-top can 98c; 2-lb. 89c; 5-lb. bag \$1.79; 10-lb. bag \$3.25; 25-lb. bag \$6.00.

Ortho Fly Killer Dry Bait. For adult flies use 4 oz. to 500 sq. ft. 1-lb. canister 98c; 5-lb. bag \$1.98.

DIAZINON 25E (25% Emulsifiable Solution.) A 25% emulsifiable solution for use in overall residual fly sprays, "spot" sprays and liquid baits for application in dairy barns and other farm buildings except poultry houses, to corrals, porches, garbage dumps, manure piles and other areas where flies congregate. It possesses excellent stability when diluted with water.

Residual sprays give control of flies, including resistant strains, for periods up to 8 weeks. It is equally effective against fly maggots.

Use 1 Pint to 6 gallons of water. Consult label for more detailed information. Price—1 Pint Bottle—\$2.85.



Smearex — (Screw Worm Preparation). ½ pt. 50c; pt. 90c; qt. \$1.75; gal. \$6.00.

Ortho 1038 Screw Worm Control. 4-oz. bottle 90c; 16 oz. \$2.75; 32 oz. \$5.00; 1 gal. \$18.00.

Note: Follow label directions on all screw worm preparations.



BIG STINKY OUTDOOR FLY TRAPS

The action of Big Stinky's Control Fluid with water on the starting bait in producing a potent scent which lures flies in the vicinity into the trap. Big Stinky actually lures flies away from houses, barns, stock pens and other buildings where you don't want them—it's irresistible—to filthy flies.

1-gal. size with 8 oz. control fluid	\$4.95
"Little Stinky" with 3 oz. control powder	2.95
"Wee Stinky" with 1½ oz. control powder	1.69
8-oz. bottle control fluid	1.50
3-oz. tube control powder89

POULTRY PREPARATIONS

Walko Tablets. Use in drinking water for baby chicks. Aids in treating cold in head, simple catarrh, sneezing, wheezing, and roup. Package 60c.

Tilghman Condition Powder. Aromatic stimulant and carminative for cattle, horses, sheep, hogs, and chickens. Package 25c. Postpaid.

Germozone. For use in drinking water. Valuable as preventive of simple bowel trouble, also ocular roup. 4 fluid oz. 50; 12 fluid oz. 90c.

Kilcide 10 Powder. (Containing 10% DDT.) For treating chickens against lice. Apply by dusting it well into the feathers in order to get the dust down to, or near, the skin surface. 1 lb. 15c; 25 lbs. \$3.45.

FERTILIZERS

KILGORE'S STORES CARRY A COMPLETE LINE OF FERTILIZERS AND FERTILIZER MATERIAL FOR TRUCK CROPS, AND GROVES, THAT IS NOT LISTED IN THIS CATALOG OR SOLD THROUGH MAIL ORDER.

All prices quoted are f.o.b. and subject to change without notice.

CAMELLIA AND AZALEA PLANT FOOD

A special prepared fertilizer for camellias and azaleas. Containing the three primary plant foods, plus secondary elements. Price 5 lbs. 65c; 10 lbs. \$1.05; 25 lbs. \$1.80; 100 lbs. \$5.15, f.o.b. Plant City.



VIGORO—Complete balanced plant food supplies the nutrient elements plants must get from the soil for best growth and development. Vigoro is odorless, sanitary and easy to apply. Made for Florida soils and conditions. Use it on everything you grow. 1 lb. 15c; 5 lbs. 55c; 10 lbs. 95c; 25 lbs. \$1.70; 50 lbs. \$2.55; 100 lbs. \$4.85 f.o.b. Plant City or any Kilgore store. Write for prices on larger quantities.

Granular Aero Cyanamid. A fertilizer carrying both nitrogen 22% and lime 70%. It is also valuable in helping to control soil disease organisms, especially nematodes. Packed in convenient, even weight, paper lined, 100-lb. sacks. 100 lbs. \$4.50 f.o.b. Plant City; ton \$85.00.

Cyanamid—Lawn and Garden

10-lb. bag \$1.85; 25-lb. bag \$3.75; 50-lb. bag \$6.50

SOLUBLE PLANT SEEDS

HY-GRO. A soluble plant food containing the three major plant food elements, nitrogen, phosphoric acid and potash, plus minor elements.

An excellent starter solution for potted plants, seedbeds and transplanting. Use at the rate of two tablespoonfuls dissolved in one gallon water or 1 teaspoonful to quart. Lawns can be easily watered and fertilized at the same time by attaching a Hy-Gro Gat sprayer (see page 71) to your garden hose.

Hy-Gro can be used on growing and fruiting plants as a plant stimulant spray, at the rate of 4 lbs. to 100 gals. of water.

3 oz. 35c; 1 lb. \$1.00; 2 lbs. \$1.45; 5-lb. bag \$2.15; 10-lb. bag \$4.00; 100 lbs. \$25.00, f.o.b. Plant City.

ORCHID HY-GRO. A special prepared formula for feeding orchids. Use 1 teaspoon per gal. of water for spray or watering. 14 oz. jar \$1.25; 25-lb. drum \$13.85.

GRO-CROP. A liquid plant food containing nitrogen, phosphorous and potash, plus minor elements needed to produce healthy plant growth. Used as ground application, starter solution and foliage spray.

Compatible with most insecticides and fungicides spray. Excellent for airplane spraying. It is a liquid and won't clog nozzles.

Use 2 tablespoonfuls per gallon water, or 2 qts. per 100 gallons.


8 oz. 59c; 16 oz. 98c; 32 oz. \$1.79; 1 gal. \$4.25; 30-gal. drums \$97.20.

NUTRI-LEAF 60. A soluble fertilizer containing 20% nitrogen, 20% phosphoric acid and 20% potash plus minor elements. Compatible with most insecticides and fungicides as a foliage spray for plant stimulant. Increases growth and size of fruit and prolongs the life of the plant and picking period. Use 5 lbs. per 100 gal. water. 5-lb. bag \$1.45; 30-lb. bag \$8.25.



VERTAGREEN is a high quality, scientifically blended plant food, for lawns, flowers, shrubs, trees, vegetables. Contains nitrogen, phosphorous and potash, besides several other important plant food elements. Clean, odorless, easy to apply. Used with outstanding success.

Price per lb. 15c; 5 lbs. 55c; 10 lbs. 95c; 25 lbs. \$1.70; 50 lbs. \$2.55; 100 lbs. \$4.85 f.o.b. Plant City, or Kilgore Stores.



PULVERIZED SHEEP MANURE

Sheep manure is highly concentrated, with all weed seeds killed. It supplies the humus and natural plant-food so greatly needed by Florida soils. For lawns, flowers and shrubs use 10 to 12 lbs. per 100 sq. ft. For commercial vegetable growing use 1,000 to 1,500 lbs. per acre drilled in the row. 25 lbs. \$1.00; 50 lbs. \$1.75; 100 lbs. \$3.50; ton \$67.50.


DAIRY ORGANIC COMPOST

High in bacteria. Supplies humus and organic plant food so greatly needed in most Florida soils. Excellent for starting lawns, flowers, shrubs and garden vegetables. Odorless, and will not burn. Weed seed is destroyed in processing.

Use 20 to 25 lbs. per 100 square ft. 5-lb. bag 65c; 50-lb. bag \$2.50; ton \$80.00.

VHPF "VIP"—A soluble fertilizer—containing the three major plant foods, nitrogen, phosphoric acid and potash plus essential minor elements. Used as a starter solution, a nutritional spray, proved very successful in combination with fungicide and insecticide sprays on growing and fruiting plants as a stimulant. Use 1 lb. to 20 gals. or 6 lbs. to 100 gals. water.

Price: 1-lb. can 50c; 3-lb. bag 75c; 6-lb. bag \$1.47; 30-lb. bag \$7.20, f.o.b. Plant City or Kilgore stores.



HYPONEX—Valuable in seedbeds and to start plants after transplanting. Comes in a powder which makes liquid plant food containing all plant food elements plus vitamin B₁. Dilute 1 oz. in 6 gals. water. 1 oz. 10c; 3 oz. 25c; 7 oz. 50c; 1 lb. \$1.00; 10 lbs. \$8.00, f.o.b. Plant City.

NUGREEN. A soluble plant food containing 45% nitrogen in the form of urea. Gives excellent results when quick acting nitrogen is needed. Can be applied as a foliage spray in combination with insecticides and fungicides at the rate of 4 to 5 lbs. per 100 gals. Can also be applied through irrigation or as top dresser. 80-lb. bag \$5.80, ton \$137.00.

ORTHO-GRO. A liquid plant food for potted plants, flowers, lawns and gardens. 4-oz. bottle 35c; 16 oz. 90c; 32 oz. \$1.50; 1 gal. \$4.95.

PLANTABBS FOR HOUSE PLANTS. Plantabbs are a complete, balanced plant food of the high analysis 11-15-20. Use Plantabbs for all indoor potted plants and flowers, window and balcony boxes. Also for garden flowers and vegetables. Plantabbs with vitamin B₁ make vastly stronger root growth.

10 Tablets 10c; 30, 25c; 75, 50c; 200, \$1.00; 1000, \$3.50.



NUTRIENTS



ACTIVO. A compost activator. Converts compost quickly and easily into available plant food.

No. 2 Size for 350 lbs. of Compost\$1.00
No. 7 Size for 1200 lbs. of Compost 2.95

ADCO.—A clean odorless powder for converting leaves, grass clippings etc. into compost. 5 lbs. makes 200 lbs. organic compost. 5-lb. bag \$1.35; 20-lb. bag \$3.45.

ALUMINUM SULPHATE. For increasing acidity of soils where aluminum toxicity is no factor. 10c per lb.; 100 lbs. \$4.00.

CALCIUM CHLORIDE - FLAKE. Recommended for control of blossom-end rot on peppers and tomatoes if caused by calcium deficiency. Use 4 lbs. per 100 gal. water per acre. 4 to 5 applications may be necessary. Sold in 100 lb. bags only. 100-lb. bag \$5.00

ES-MIN-EL. For soil application. Contains many essential mineral elements in proper proportions. Add to ordinary fertilizers at rate of 1 lb. or 2 lbs. to 10 lbs. regular fertilizer or 100 to 200 lbs. per ton of regular fertilizer. 5-lb. bag \$1.05; 100-lb. bag \$10.50; f.o.b. Plant City.

ES-MIN-EL. For spray or dust. For spray use 4 lbs. to 100 gals. water. For dust use 16 lbs. Es-Min-El mixed in 84 lbs. of talc, clay or other inert ingredients. Price: 4-lb. bag \$1.60, 50-lb. bag \$14.00, f.o.b. Plant City

KILGORE IMPROVED C-M-Z Spray. A combined nutritional and fungicidal spray for tomatoes, potatoes, peppers and other vegetables. Do not use on sweet corn. Apply at the rate of 6 lbs. per 100 gals. of water. 6-lb. bag \$2.50.

KILGORE'S C-M-Z BEAN SPRAY—For use on beans for the correction of copper, manganese and zinc deficiencies and for the control of rust and powdery mildew.

Use 10 pounds per 100 gallons of water at intervals of 10 days. Where rust is a serious problem apply every 5 days. The addition of spreaders is not necessary. Price: 10-lb. bag \$1.70; 50-lb. bag \$7.15.

HORMONES

HORMODIN POWDERS. Hormodin is a hormone-like plant growth substance. Its principal use is to stimulate rapid growth on cuttings. It has wide-spread use in propagating cuttings. Many cuttings formerly impossible to propagate may now be rooted with the aid of Hormodin. Easy to use, consisting of simply dropping moistened stems in the powder and then planting. The cost of treatment is only a fraction of a cent per cutting.

No. 1—For Chrysanthemums, Poinsettias, Roses, etc. ¼ oz. 25c; 1¾ oz. 50c; 1 lb. \$3.00.

No. 2—For semi-woody plants and most common shrubs.

1¾ oz. 75c; 1 lb. \$4.50.


No. 3—For resistant species, such as Evergreens and dormant leafless cuttings.

1 oz. \$1.00; ½ lb. \$4.50.

Combination Package, powders Nos. 1, 2 and 3, in carton, 75c.

FRUITONE is a wettable powder containing several hormones plus an effective wetting agent for securing

KILGORE'S IRON-AID. (Code No. X-5). Contains Versen-ol, a newly developed chelated iron compound, which is completely available to plants under most soil conditions—Designed for the correction of iron chlorosis wherever symptoms are indicated in flowers or shrubs. These symptoms indicate a tendency of paling the areas between leaf veins, while the veins themselves still retain their original green color. We suggest the use of it in all cases where iron chlorosis is indicated as well as in questionable borderline cases. Consult label for specific directions before applying.



1 lb. 60c; 5 lbs. \$1.25; 10 lbs. \$2.00; 25 lbs. \$3.75;
50 lbs. \$5.90; 100 lbs. \$11.80 f.o.b.

LIME, HYDRATED. Land-sweetener. Use a half ton to a ton per acre, broadcast, before breaking the ground. If soil is new and sour, the application should be heavier. Also used for the neutralization of manganese, zinc and copper sulphates. 10-lb. bag 30c; 50-lb. bag 90c; ton \$31.00; ½ ton at ton rate, f.o.b. Plant City.

MANGANESE SULPHATE (70% spray grade). A quick acting material for applying manganese to plants in the form of a spray when plants are turning yellow on very alkaline soils. Apply at the rate of 2 lbs. to 50 gallons of water. 1 lb. 10c; 50-lb. bag \$2.60, f.o.b. Plant City.

MAGNESIUM SULPHATE (Epson Salts)—For magnesium deficiency. 100 lbs. \$6.50; less than 100 lbs. 10c per lb.

SULPHUR-MANGANESE DUST—10%. (Code No. X-2)—For manganese deficiencies in beans, peppers, etc. Lb. 10c; 25 lbs. \$1.60; 50 lbs. \$2.20; 100 lbs. \$4.35.

VERSEN-OL.—A chelated iron compound for soil application—soluble in water. Not recommended for foliar applications.

1 lb. can \$2.98, f.o.b. Plant City.

ZINC SULPHATE. For zinc deficiencies in beans, tomatoes, etc. 100 lbs. \$10.95; less than 100 lbs. 13c per lb.

a thorough coverage. Fruitone when sprayed on plant blossoms and young set fruit will help to prevent premature dropping. Cut flowers and twigs dipped in Fruitone will hold up longer.

Use 1 level teaspoonful to 1 gal.; 2/5-oz. to 5 gals.; 2 oz. to 25 gals.; 8 oz. to 100 gals of water.

Price: ¾-oz. pkg. 25c; 2-oz. can \$1.00; 12-oz. can \$5.00.

ROOTONE. A hormone powder to hasten the rooting of cuttings. It is easy, economical, safe and sure treatment for rooting cuttings. Low cost—1/30 to 1/50 cent per cutting.

¼ oz. 25c; 2 oz. \$1.00; 1 lb. \$5.00; prepaid.

Rootone No. 10 recommended especially for rooting azaleas and camellias.

2 oz. jar \$2.00; 1 lb. jar \$10.00.

2 oz. pkg. will treat up to 3,000 cuttings.

TRANSPLANTONE supplies the plant with chemicals which stimulate root formation and growth after transplanting time. ½ oz. makes five gallons solution.

½ oz. 25c; 3-oz. can \$1.00; 1-lb. can \$4.00.

SEED INOCULANTS

All prices quoted are postpaid.

When planting a legume feed or cover crop, such as clovers, cowpeas, beggarweed, crotalaria, soy beans and velvet beans, first inoculate the seed. This helps the crop to add the maximum amount of valuable nitrogen plant food material from the air to your soil. A good leguminous cover crop can take 50 to 150 pounds of nitrogen per acre from the air—if inoculated. For your soil's sake, inoculate all legume seed.



Group A—Alfalfa and Sweet Clover	1 bu. size 60c 100 lb. size 80c
Group B—Clovers White Dutch Special	1 bu. size 60c 100 lb. size 80c
Group C—Vetch, Garden and Sweet Peas	100 lb. size 55c
Group D—Beans—all varieties	1 bu. size 60c
Group E—Peanuts, Cowpeas, Velvet Beans, Crotalaria, Beggarweed and Alyce Clover	100 lb. size 55c
Group L—Lespedeza	100 lb. size 60c
Lupines—all varieties	100 lb. size 55c
Group S—Soy Beans	2 bu. size 45c 5 bu. size 65c
Special Cultures: Lima Bean Trefoil, Dalea or Woods Clover, Crown Vetch, Chick Peas, Hairy Indigo and Sesbania	1/8 bu. size 25c 1 bu. size 70c
Garden Size, for Garden Peas, Sweet Peas, All Beans, Soy Beans and Lupines.	Each package will inoculate 12 lbs. seed—25c

INOCULANT MADE IN FLORIDA ESPECIALLY FOR FLORIDA SOILS

Dixie Inoculant for	Quantity of seed will inoculate (size)	Price
Soy Beans	2 bushels 5 bushels	\$.30 .60
Lespedeza, Alyce Clover Hairy Indigo	100 pounds50
Cowpeas, Crotalaria, Lima Beans, Peanuts and Velvet Beans	100 pounds50
All Vetches, Austrian Peas, Wild Winter Peas, Garden and Sweet Peas	100 pounds50
Lupines, Special for Sweet, Yellow and Blue	100 pounds50
Clovers, Red Crimson, Ladino, White Dutch and Louisiana White	100 pounds70
Alfalfa, Hubam, Floranna, Sweet and Bur Clovers	100 pounds70
Garden Size	8 pounds15



Culture Group 1 Alfalfa, Bur, Button, Hubam Clovers	60 lb. size 60c 100 lb. size 80c
Culture Group 2 Clovers—Alsike, Crimson, Red, White, Ladino	60 lb. size 60c 100 lb. size 80c
Culture Group 3 Vetches, English Peas	100 lb. size 55c
Culture Group 4 Beans—Snap, Pole, Navy, Pinto, Kidney	60 lb. size 60c
Culture Group 5 Lupines	100 lb. size 55c
Culture Group 6 Soy Beans	2 bu. size 45c 5 bu. size 65c
Culture Group 7 Peanuts, Cowpeas, Velvet Beans, Crotalaria, Beggar Weed and Alyce Clover	100 lb. size 55c 5 bu. size 60c

Garden size, inoculates up to 6 lbs. seed. Peas, Beans, Sweet Peas, Lupines, Soy BeansPrice 15c

POULTRY SUPPLIES

All prices f.o.b. Plant City or Kilgore Stores and subject to change without notice.

More complete line of poultry supplies carried in Kilgore stores.

HUDSON HANDY FILL CHICK FOUNTAIN



No. A-4—Capacity one gallon	\$1.45
No. A-8—Capacity two gallons ..	1.70
C-3—Capacity three gallons	2.85
C-5—Capacity five gallons	3.65

MASON JAR FOUNTAIN AND FEEDER

C-25 Plastic Pan with three point clip to fit Mason jar	\$.20
---	--------

REEL TOP FEEDERS

No. 118 Metal Feeder 18-inch length	\$.50
No. 224 Metal Feeder, 24-inch length75
No. 336 Metal Feeder, 36-inch length	1.00

HUDSON "KLEEN-EZY" REEL TOP FEEDERS

No. 6360A—36 in. length—Reel Top	\$1.75
No. 5480A—Feeder, 48-inch length Reel Top	2.40

GRILL TOP FEEDER

No. 7480A Feeder, 48-in. length Grill Top	\$3.65
---	--------

Key Ring Leg Bands. Colors, red, yellow, green, white and blue. Sizes for baby chick, pigeon, bantam, medium, intermediate and large. Specify size and color. 1c each; 75c per 100.

	Each	Doz.
Nest Eggs—china	\$.05	\$.50

THERMOMETERS

Black oxidized brass scales, white filled figures and graduations, mercury tubes.
Weather thermometer, No. 5151, 8-in. wood\$.75
Weather thermometer, No. 5140, 7 7/8-in. wood case 1.00
Weather thermometer, No. 5132, 7-in. Metal Utility 1.50

KNAPSACK AND HAND DUSTERS

THE CHAMPION NO. 100 DOUBLE ACTION DUSTER is carried like a knapsack. It fits comfortably on the back and leaves both hands free for efficient operation. The easily operated bellows throws a screen of dust exactly where wanted within a range of 30 feet. The complete duster weighs only 14 pounds empty and will hold as much as 17 pounds of fluffy powder. Can be converted to single action by removing upper intake valve.


Price \$29.35, Prepaid



Champion No. 100
Double Action Duster


ROOT HAND GUN MODEL C-3A

Root Hand Duster C-3A
with "Y" Attachment
\$34.20, Postpaid



Root Hand Duster C-3B
(Less "Y" Attachment)
Price \$31.25, Postpaid

CHAMPION HAND DUSTER — MODEL 103



Champion Hand
Duster Model 103

Simplicity of design and ease of operation make the Champion Hand duster ideal equipment for flowers, vegetables, shrubs and small trees. It is used extensively in home gardens and small estates as well as in experimental and show gardens; very efficient and economical for spot dusting small plants in the field such as watermelons, etc. It is of highest quality construction. All parts


are enclosed and the steel tank is lead coated for long life. It is extremely light in weight and will carry up to eight pounds of powder, although it operates effectively with only a fraction of an ounce of powder in the tank.

Price \$9.85, Prepaid

LITTLE CHAMP NO. 902 Rotary Hand Duster. Made especially for home garden. Sturdy built, 8" tall. Wt. 2¾ lbs. Capacity 2 lbs. Easy to turn. Puts out a strong blast of dust. \$9.85.

HAND DUSTERS

Made of aluminum sheet and castings for lightest weight. The Hudson Roto-Power "806" is designed for growers who want the easiest turning and lightest carrying duster, one that produces the high velocity and volume of air necessary to break up dust and blast it right into the plant. Holds 14 lbs. average density dust. Applies from 5 to 45 lbs. of dust per acre. High velocity: 1 crank turn produces 60½ fan revolutions.



Hudson No. 806B Roto-Power Duster (Aluminum)
Price \$39.80, Postpaid
f.o.b. Plant City or Kilgore stores

SMALL HAND DUSTERS

f.o.b. Plant City



Brown Crystal Duster, Glass Jar, capacity 1 quart	\$2.40
Hudson No. 612-A Patrol Duster	2.10
Hudson No. 673 Cadet	.75
Hudson Admiral No. 766-C	2.90
Hudson Adjutant No. 674	1.45
Root P-1	2.45
Root P-2	3.00
Dobbins No. 132, 1-lb. capacity, price	3.00



HUDSON ADMIRAL No. 766-C

An ideal duster for the average home garden, for use on vegetables, flowers, trees, and shrubs.

RESPIRATORS —

Approved for parathion dusting	each \$5.95
Replacement cartridges, No. 11-A (4 to box)	Box 2.40
Replacement filters, No. R-490 (10 to box)	Box 1.60
Monogoggles No. 93A Clear Lens	2.75
Monogoggles No. 93A Green Lens	2.90

Postpaid

FLAME SPRAYERS

Flame Sprayers have many uses. Develops 2000° F. of easy portable heat in a flame that can be directed at the will of the operator. Can also be converted to a liquid sprayer by using spray attachment No. 225.

Hudson No. 211 Saf-T-Lok, Galv. Tank, Brass Pump. Capacity 2¼ gal., price	\$20.50
Dobbins No. 410, 4-gal. capacity Flame Sprayer, price	\$35.85
No. 225 Liquid Spray Attachment, price	7.15

POWER AND HAND SPRAYERS

Prices quoted are f.o.b. Plant City unless otherwise indicated

HUDSON POWER SPRAYER

High pressure sprayer recommended for estates, small groves, dairy barn and cattle spraying, and landscape gardens. Handles all insecticides, fungicides, weed control chemicals, and whitewash.



No. 31030 Matador Cushion Tires, 30-gal. tank ..\$269.75
No. 31015, Matador Cushion Tires, 15-gal. tank .. 247.50

COMPRESSED AIR SPRAYERS

Prepaid

Compressed air sprayers are strong, efficient and easy to use. Ideal for dozens of spraying jobs in homes, truck gardens, dairy barns, poultry houses, etc.

Kilgore No. 178,
1½ gal. capacity\$ 5.95
Kilgore No. 179,
3½ gal. capacity 6.95



Brown Compressed Air,
Galv. Open-Head
No. 64-D (4-gal.)\$15.65
Hudson Compressed Air
Perfection Sprayer
(4-gal.) No. 310-B 19.10
Hudson Bugwiser No. 220-B
(2½-gal.) 10.00
Hudson Leader No. 293-B
(3½-gal.) 12.85
Hudson Climax No. 335-A
(3-gal.) 16.75
Dobbins 44-A (4 gal.) 19.80
Dobbins No. 21-A
(3½-gal. capacity) 13.50
Dobbins No. 24-A Budget
(3½-gal. capacity) 10.95



Leader No. 293-B

SPRAYER HOSE

¾" Hose. Price per ft.25


CHAMPION SPRAYER MODEL-1



by the operating lever. This lever also operates the brass agitator which keeps 4½ gallons of any liquid well mixed.

Price \$33.75, Postpaid

This all-purpose portable sprayer is top in its field. The Champion combines in one unit all the desirable qualities of unusual power, simple operation, effective control, continuous agitation, and long life. Uniform pressure of 200 lbs. in the sturdy pressure cylinder is pumped as needed



Hudson Mercury No. 333

Household sprayers

f.o.b. Plant City or Kilgore stores

No. 208, Hudson Nebu-lizer, Super Cloud
8 oz. capacity\$.55
No. 333, Hudson Nebu-lizer, Super Fog
14 oz. capacity80
No. 444, Hudson Nebu-lizer, Super Misty
27 oz. capacity95
No. 409, Hudson, Nebu-lizer, Super Rapid
2 qt. capacity 1.80


Household continuous sprayers

f.o.b. Plant City or Kilgore stores

Hudson Cardinal No. 436, 3-quart capacity 3.10
Hudson Mercury No. 433, 1-quart capacity 2.10
Hudson Comet No. 431A, 1-quart capacity 1.55
Ken Sprayall (Trombone Type) No. 202 9.95
No. 214 Redisprayer w/Extension 7.95
Ken-Sprayall Little Giant No. 224 4.95

GAT SPRAYER. Attaches to garden hose for dissolving, mixing and spraying Hy-Gro and other soluble plant foods.

Empty 2-lb. Hy-Gro jar with
Gat Sprayer\$2.50



Insect-O-Gun

BRADSON SPRAY GUNS

For use with standard garden hose in obtaining pressure. Insect-O-Guns, Gard-N-Gun and Bug-Gun for spraying insecticides and fungicides.




Bug-Gun

Gro-Gun for application of soluble fertilizers and plant foods.


Insect-O-Guns—Model-3\$5.95
Gard-N-Gun—Model-2 3.95
Gro-Gun—Model-20 2.95
Bug-Gun—Model-4 2.65

LAWN SPRINKLERS RAIN KING

Model-K Adjustable from 5 to 50 feet\$ 9.75
Model-K-GSS., Rust resistant for coastal areas .. 10.25
Model-D Double nozzle adjustable sprinkler 5.95
Model N1 Garden hose nozzle 1.39
Park Sprinkler Model 911 2.95



Model K



Model D

Miscellaneous Farm and Garden Tools

All prices quoted are f.o.b. Plant City or Kilgore stores



Planet Jr.
No. 4 Hill
and Drill
Seeder

No. 4-D Seeder—Shipping weight 41 lbs. Price ..\$32.30
No. 4 Seeder—Shipping weight 50 lbs. Price 40.45
No. 300-A Seeder—Shipping weight 55 lbs. Price 41.00




No. 17 Single
Wheel Hoe.
Shipping wt.
22 lbs.\$16.75




No. 19 Wheel Hoe.
Shipping weight 18
lbs. Price\$11.35

Planet Jr. No. 219
Fertilizer Distributor



No. 219 Fertilizer Dis-
tributor. Shipping
weight 33 lbs. Price..\$24.75
No. 215 Fertilizer At-
tachment for No. 219
Distributor. Ship-
ping weight 17 lbs.
Price 12.50
No. 8556X, Stand for
No. 215. Shipping
weight 2 lbs. Price.. 1.20
No. 8558X, Covers for
No. 215. Shipping
weight 2 lbs. Price.. 1.55

No. 119 High Wheel Plow. Shipping wt. 18 lbs....\$11.75



BPI (2 HP) Planet
Jr. Tractor with
rubber tires
Price \$199.75


B - 12 cultivator
attachment as il-
lustrated. Ship-
ping wt. 76 lbs.
Price\$45.40

B-8 (2½ h.p.) Tractor with 600 x 12 Rubber Tires.
Shipping weight 197 lbs.\$261.00

Tuffy (1 h.p.) Shipping weight 90 lbs. 102.00

Super Tuffy (1½ h.p.) Shipping weight 99 lbs... 150.00

Write for Panet Jr. Tractor catalog and prices on
tools and attachments for tractor.




Planet Jr.
No. 220
Fertilizer
Distributor

Shipping
Weight
115 lbs.

No. 220 Fertilizer distributor\$66.50

Planet Jr.
No. 13 Double
and Single
Wheel Hoe



No. 13 Double and Single Wheel Hoe—Shipping
weight 27 lbs. Price\$19.70

Planet Jr. Grass Edger No. 2. Shipping wt. 3¼
lbs. Price 3.25

CYCLONE SEEDER

The Cyclone will sow many acres per
hour, adjustable to any amount per
acre.

In ordering seeder to be used with
power take-off specify make and
model of tractor to be used on. Elec-
tric motor model operates from 6 volt
battery from jeep or truck.

Hand Seeder\$4.40

Power Model M-1 Electric Motor\$40.50

Power Model S-1 for Power Take-Off 50.70

Extension Hoppers for M-1 and S-1—
1½ bu. capacity 6.30




INSCO Soil Fumigant Applicator

A 4-gal. tank gravity flow applicator for single row
application. Can be attached to horse drawn plow or
small tractors.

Price f.o.b. Plant City or Kilgore stores\$15.00

MISCELLANEOUS FARM and GARDEN TOOLS

Prepaid unless specified. Prices subject to change without notice.



ECLIPSE builds precision power mowers with extra features for unmatched performance in low-cost maintenance of beautiful lawns. Balanced design, durable construction, dependable power, self-propelled, precise cutting action and effortless handling make lawn mowing easier, faster and better.

Eclipse Lark 18-in.	\$114.50
Eclipse Rocket 20-in. (Recoil starter)	139.50
Eclipse Parkhound 21-in.	164.50


Prices f.o.b. Plant City or Kilgore stores
Sharpeners for reel type mowers available

Eclipse—16-inch Hand Mower	\$34.50
Zephyr 16-inch	30.00
Vogue 16-inch	27.00

Prices f.o.b. Plant City or Kilgore stores

NO. 811—3 PC. GARDEN SET

Sturdy, attractive hand made tools finished in baked green enamel. Set includes trowel, fork and cultivator.




No. 800 Trowel	25c each
No. 801 Fork	25c each
No. 802 Cultivator	25c each
No. 804 Transplanter	25c each
No. 805 Weed Cutter	25c each
No. 811 3-piece Set	80c per set

WISS GARDEN TOOLS

Postpaid

FLOWER SHEARS

FH-4 Cutter Holder, Chrome Plated 6½"	\$2.50
---	--------



No. 808 Draw Cut Pruning Shears

No. 808 Draw Cut	\$3.95
No. 607-R Pruning Shears	2.95
No. C-1 Orange Clippers	2.75



No. 8½-E Hedge Shears

No. 9-A Hedge Shears	\$4.75
No. 8½-E Hedge Shears	5.50
No. 9½-E Hedge Shears	5.95



No. 224-T Lopping Shears

No. 20-V 20 inch	\$4.95
No. 321-T 21 inch	6.75
No. 224-T 24 inch	6.25



No. 5600 Forged Grass Shears

No. 5600 Forged Grass Shears	\$2.95
No. 500 Grass Shears	1.39
No. 900 Grass Shears—Long Handle	4.75
No. 801 "Quick Trim"	1.95
No. 700 Grass Shears	2.25

LAWN FERTILIZER DISTRIBUTORS

f.o.b. Plant City or Kilgore stores




MASTERS MODEL 30

50-lb. capacity. All steel construction, tubular steel handle with two "bicycle type" rubber grips. Rubber tires. Finest home lawn spreader ever developed. Price \$13.95.

MASTERS MODEL 25

20-lb. capacity all steel construction with tubular steel handle, rubber handle grip. Steel wheels with rubber tires. For the customers who want "something better" in a small spreader. Price \$10.45.



MASTERS HANDI-CART Model YM-66

The Masters Handi-Cart has an outstanding reputation among users as a sturdy, dependable tool for hundreds of moving jobs around the home and garden. Capacity 3 cu. feet. Price \$8.95.

Miscellaneous Farm and Garden Supplies

Prices subject to change without notice



BIRD NEPONSET PAPER POTS

Nutrient Treated

Made of tough waterproof paper, adapted for growing plants from seed to transplanting age. Can be sold in pots. Plants get no setback when transplanted from pots. Performs the major function of clay pots at a fraction of the cost.

NEPONSET RED PAPER POTS

Neponset Red Paper pots are designed for shipping plants which have been grown in clay pots or in other containers. Shipping costs are lower by using Neponset Red

Pots because there is considerable less weight and less breakage, and the clay pots can be saved for further use.

By mail postpaid			f.o.b. Plant City		
100 pots			1000 pots		
	Vita	Red	Vita	Red	
2 inch	\$2.55	\$1.50	\$16.05	\$ 9.45	10 lbs.
2½ inch	2.65	1.60	17.05	9.55	16 lbs.
3 inch	2.95	1.70	19.15	10.05	22 lbs.
3½ inch	3.85	2.15	25.00	13.40	32 lbs.
4 inch	4.55	2.65	28.30	16.20	21 lbs.
5 inch	5.07	3.25	36.75	19.95	33 lbs.
6 inch	7.40	3.95	48.50	24.50	47 lbs.

PLANT BANDS



PICK UP LIKE THIS



SET IN PLACE



Postpaid per 100 f.o.b. Plant City per 1000

(Contains sufficient fertilizer for 10 weeks)

Vita-Bands 10

1¼ x 1¼ x 2½ inches	\$.90	\$4.30
2 x 2 x 2½ inches	1.05	5.30
2 x 2 x 3 inches	1.15	5.85
2½ x 2½ x 3 inches	1.30	6.60
3 x 3 x 3 inches	1.45	7.60
4 x 4 x 3 inches	1.80	9.90

Vita Band-D

A nutrient treated band which disintegrates in the soil.

	Postpaid per 100	f.o.b. Plant City per 1000
1½ x 1½ x 2½	\$.85	\$4.45
1¾ x 1¾ x 2½	.95	4.80
2 x 2 x 2½	1.10	5.40
2½ x 2½ x 3	1.30	6.65
4 x 4 x 3	1.95	10.20

TOM THUMB FLATS

	Per 100	f.o.b. Per 1000
No. 1 (for bands 1¼ x 1¼ x 2½)	\$5.40	\$42.20
No. 2 (for bands 2 x 2 x 2½)	6.10	54.60

Terra-Lite Vermiculite



TERRA-LITE (Vermiculite). An excellent light weight medium for starting cuttings, seed in flats, paper pots or plant bands. Can be mixed with soil in seedbeds. Increases germination by reducing damping-off. Holds moisture for several days without rewatering.

4-qt. bag 39c; ½-bu. bag 89c;
4-cu. ft. bag \$2.35,

f.o.b. Plant City

Write for complete descriptive pamphlet on Terra-Lite.

PEAT MOSS

Good to add humus and holds moisture in seedbeds. Also an excellent poultry litter.

Georgia Peat Humus, 1½ cu. ft. bag \$1.20 each
5 cu. ft. bags \$3.50 each

European Peat. Compressed in burlap

Wrapped bales 20" x 20" x 36" each \$4.95
Bulk 10c per lb. f.o.b. Plant City

SPHAGNUM MOSS

Burlap Wrapped Bales 14" x 16" x 45" each \$4.25
Bulk 50c per lb. f.o.b. Plant City

OSMUNDIA FIBRE—Used for growing Orchids.

Per Bale\$8.75
Bulk - per lb.35



RUBBER STRIPS—For Budding and Grafting

Size	Length	Width	Approx. pcs. to the lb.	Price per lb.	Price per 5 lb. pkg.
No. 520	5 in.	3-16 in.	1400	\$2.50	\$11.25
No. 620	6 in.	3-16 in.	1080	2.50	11.25
No. 838	8 in.	3-8 in.	450	2.50	11.25

Less than 1 lb. 25c per oz.

AIRWRAP

Coated plastic adapted to grove and plant propagation. Coated with nutrient and rooting hormones.

9" width—8 yd. Rolls\$1.95
9" x 8" Price per Sheet06
9" width—Less than 8 yds. per yd.25

Write for quotations on larger quantities.



AIRLAYERING ROOTING KIT

For plant propagation. Kit contains sphagnum moss, plastic wraps and tying strips. Complete instructions in box.

Kit complete (Prepaid)\$1.95

GLOVES

GREEN THUMB—(Ladies' size) Plastic

Small, medium and large98 pr.


GIRL FRIDAY

Plastic coated—knit-lined\$1.40 pr.

EEZY WEAR (Kidskin)

Ladies' size, small, medium and large\$1.60 pr.

Men's size—medium and large\$1.70 pr.



Girl Friday


Miscellaneous Farm and Garden Supplies

Prices subject to change without notice

✂️TWIST-EMS'

Postpaid

Help you support plants quickly, neatly, permanently.



Package of 35—8-inch size	10c
Package of 50—8-inch size	15c
Box of 100—8-inch size	25c
Box of 200—4-inch size	25c
Box of 100—16-inch size	50c

UTILITY TWIST-EMS

1000 1/4" x 8"	\$1.15 prepaid
1000 1/4" x 16"	3.90 prepaid




TRAIN-ETTS

New Weatherized trellis netting, ideal support for sweet peas and any vining flower or vegetable plants. Comes in three sizes. No. 1, 60 x 72 in. 75c; No. 2, 60 x 96 in. 95c; No. 3, 60 x 180 in. \$1.35, postpaid.

BAMBOO CANE STAKES

Dyed green pencil-thickness

3 ft. Stakes, per 1000 f.o.b. Plant City	\$12.50
3 ft. Stakes, per 100 postpaid	1.50
Less than 100 2 1/2c each postpaid.	
4 ft. Stakes, per 1000 f.o.b. Plant City	16.00
4 ft. Stakes, per 100 postpaid	2.00
Less than 100 3c each postpaid.	



LABELS—Plant and garden (postpaid)

	Per 100	Per M
4" Pot Labels—Painted—(1M per ctn.)	\$.45	\$3.45
5" Pot Labels—Painted—(1M per ctn.)50	3.80
6" Pot Labels—Painted—(1M per ctn.)55	4.20
8" Garden Labels—Painted—(250 per ctn.)	1.00	8.65
10" Garden Labels—Painted—(250 per ctn.)	1.10	9.75
12" Garden Labels—Painted—(100 per ctn.)	1.80	16.50
3 1/2" Copper Wired—Painted—(1M per ctn.)55	4.35

GRAFTING WAX—Postpaid

A prepared wax recommended for all hand and brush grafting operations. 1/4 lb. 40c; 1/2 lb. 55c; 1 lb. 85c.

BUDDING AND GRAFTING TAPE

Grafting Tape 1/4-in. wide, 60-yd. Roll	\$1.00
Grafting Tape 1/2-in. wide, 60-yd. Roll	1.30
Grafting Tape 1-in. wide, 60-yd. Roll	2.00

BUDDING KNIVES—Postpaid

No. 11 Budding Knife, brass lining, 4 1/2 inch, Price	\$3.50
No. 21SH Florist Knife, 2 blades, Price	3.00
No. 8 1/2C Grafting Blade, Price	1.60
No. 9C Budding Blade, Price	1.60

HOTKAPS

Hotkaps are "individual hothouses" for every plant—protecting them against frost, rain, wind, insects and crusting. They force plant growth, making hardier, healthier plants and producing crops two or three weeks earlier. They may be used with practically any crop that needs protection, but are especially suited for use with vine crops such as tomatoes, cucumbers, cantaloupes and watermelons.



25 pkg. Complete with Paper Setter, each	\$.95
100 with Fibreboard Garden Setter	3.45
250 with Fibreboard Garden Setter	6.50
1000 without Setter	19.95
5000 without Setter, per 1000	19.85
10000 without Setter, per 1000	19.75
25000 without Setter, per 1000	19.65
Fibreboard Garden Setter, alone20
Steel Setter, each	1.95

Prices quoted are f.o.b. Plant City and all Kilgore stores

LABEL PASTE

Ours is a genuine wheat paste, especially made for pasting labels on vegetables and fruit boxes. Mixes easily in cold or warm water. Will not discolor labels. Unaffected by water or atmospheric changes after drying.

Prices quoted are f.o.b. Plant City and all Kilgore stores
25c per lb.; 100 lbs. \$22.75

TWINE (Jute)

Soft vegetable 2 ply (app. 1520 ft. to lb.) tying twine, excellent quality. Used for tying tomato plants and bunching vegetables.

Per lb. 47c; 50 to 60-lb. reels, per lb. 45c.
1/2-lb. balls—30c each.

COTTON TWINE (Variegated)

Used for tying pole beans. 2 1/2-lb. cones. Price per cone \$3.00.

f.o.b. Plant City

RAFFIA

The plain or natural color comes in uneven weight hanks. 1 to 3 lbs. \$1.00 lb.; 10 lbs. \$9.00. Prepaid.

TARPAULINS

Write for prices stating size desired.

HAND TALLY REGISTERS—Pocket Size

A fast and accurate way to count live stock, poultry, watermelons, plants, packed crates, etc. No. O counts to 999, \$3.80. No. 1 counts to 9999, \$4.90.


SUDBURY SOIL

TEST KIT

Home Garden Model

Complete equipment with instructions and solutions for 20 tests, for Nitrogen, Phosphorous, Potash and Acidity.

\$2.98 Postpaid



JR. PROFESSIONAL MODEL

Contains complete equipment and instructions—solutions for 50 tests for nitrogen, phosphorous, potash and acidity (pH). Transparent plastic case. Price \$5.89 Postpaid.


Two Minute Acidity Kit

Write for prices on larger models.

CANNING DEPARTMENT

Prices subject to change without notice

Florida offers the greatest opportunity for home canning of any state in the Union, due to the fact that every month in the year produces some fruit or vegetable, and yet there is probably less canning done here than in almost any other state; but we are becoming more thrifty each year in this respect.



BURPEE PRESSURE COOKER

Pressure cooker or canner made by forming heavy plate aluminum. Very strong — half the weight of "cast" aluminum. Guaranteed not to crack, break or split. It will not stain because there are no "pores" in this metal to fill up with food and create a bad odor.

MODEL AR25


Holds 10 qt. or 17 pt. cans or 7 qt. or 14 pt. jars. Equipped with canning basket, recipe book, and cooking pans.

Postpaid

Burpee AR25 Heavy Type Aluminum\$29.95

BURPEE PRESSURE CANNER

Pressure canner made by forming heavy plate aluminum. Very strong — half the weight of "cast" aluminum. Guaranteed not to crack, break or split. It will not stain because there are no "pores" in this metal to fill up with food and create a bad odor.



MODEL AR1525

Holds 15 qt. or 25 pt. cans, or 14 qt. or 21 pt. jars. Equipped with 2 canning baskets, recipe book, and pans.

Postpaid

Burpee AR1525 Aluminum\$34.95


Home canning is fine training for the children, both girls and boys, as it teaches thrift. In some states the women and children add several hundred dollars to the family income during the year from home canning, thus securing money for many other things. It matters not whether products are used at home or sold, it's money saved just the same.

Hardly is there a day in Florida when there is not some luscious vegetable or fruit that can easily be conserved for later use. Meats, poultry and fish can also be conserved by modern safe and scientific methods—same methods as used by commercial canners. A pantry full of wholesome home-canned products means a better balanced diet for the entire family the year round. It is fascinating, too, avoids waste and builds a bank account.

HOT-LIFT CAN TONGS




Hot-Lift Can Tongs, price Postpaid\$.60



BURPEE SIMPLEX SEALER WITH CAN SAVER ATTACHMENT

The Leader in Can Sealers

Seals cans quickly and easily. Set the can in position, spin the crank with the right hand and move a lever with the left hand. Made of hardened steel for long life. Rustproof. Equipped with attachment which simplifies changing from one size can to another. Clamps on a table. Seals, opens and reflanges two sizes of cans, the No. 2 and No. 3. Tested canning recipes and illustrated instructions with every sealer. Net weight 11 lbs. Price \$17.95, prepaid.



*Simple, accurate
and foolproof.
Insert can, turn
crank, and can
is automatically
sealed.*

*So simple!
Complete canning
recipe book with
each machine.*

DIXIE SIMPLIFIED AUTOMATIC HOME CAN SEALER

Will seal No. 1, No. 2, No. 2½ and No. 3 Sanitary cans.

Prepaid

With No. 2 or No. 3 chuck and base\$35.00

Extra chuck and base 8.50

CANNING DEPARTMENT

SANITARY CANS



Our sanitary open top cans are ideal for home canning. The finest fruits or vegetables and all your hard work can be wasted if the cans are not right.

Our cans for home canning are uniform, accurate, dependable, and strong. For years they have been recognized as cans of highest quality, preferred by home canners.

CANS

f. o. b. Plant City

	No. in Case	Case	1,000
No. 2 Plain Sanitary cans, wt. 26 lbs.	100	\$4.95	\$48.55
No. 2 Plain Sanitary cans, wt. 6 lbs.	24	1.30	51.85
No. 3 Plain Sanitary cans, wt. 36 lbs.	100	7.35	72.65
No. 3 Plain Sanitary cans, wt. 9 lbs.	24	1.90	77.35
No. 2 Sanitary lacquered cans, wt. 26 lbs.	100	4.85	47.45
No. 2 Lacquered cans, wt. 6 lbs.	24	1.45	58.40
No. 3 Lacquered cans, wt. 36 lbs.	100	7.65	75.65
No. 3 Lacquered cans, wt. 9 lbs.	24	1.95	80.40

These cans are of the highest quality, uniform, accurate, dependable, and strong.

Friction top or syrup cans are used for packaging syrup, honey and for many other purposes.



10-LB. SYRUP CAN

	No. in Case	Case	1,000
1½ Friction-top cans, wt. 8 lbs.	36	\$2.15	\$ 59.50
No. 2½ Friction-top (approx. ¼ gal.) cans, wt. 11 lbs.	24	2.15	88.20
No. 5 Friction-top, holds 4½ lbs. of Honey, wt. 24 lbs.	50	6.65	131.55
No. 5 Friction-top cans, wt. 6 lbs.	12	1.70	139.30
No. 10 Friction-top (approx. 1 gal.) cans, wt. 39 lbs.	50	9.75	193.95
5 gal. Square Honey Cans—each 85c; case (16)			\$11.25.
50-lb. Lard Can—each 90c; case of 12—			\$10.35.
Corrugated Shipping Cartons for 5-gal. Honey Cans, each 25c; 10 for \$2.30; 25 \$5.50; <i>f.o.b. Plant City.</i>			

CANNING SUPPLIES

<i>f.o.b. Plant City</i>	Per 100
Extra Covers for No. 2 Plain Sanitary cans	\$1.25
Extra Covers for No. 3 Plain Sanitary cans	1.50

CANNING LABELS

<i>f.o.b. Plant City</i>		
Beans, Corn, Blackeye Peas, Tomato and Syrup.		
Price, No. 1½ and 2½, Syrup ..	.25	\$.50
Price, No. 2 and 3, Vegetables ..	.30	1.00
Price, No. 5 and No. 10, Syrup ..	.25	.60
Rubber Stamp, two-line \$1.75; three-line		\$2.00

ATLAS FRUIT JARS

f.o.b. Plant City

Atlas Jars are made of clear crystal glass of highest quality giving a natural color to packed fruits and vegetables. Best preserving jar on the market, guaranteed to give entire satisfaction when properly used.



Atlas Square Mason Jars

This style jar has been on the market for over 50 years but has recently been improved by the strong shoulder feature.

FRUIT JARS

f.o.b. Plant City

	Per doz.
Atlas Square Mason Jars, ½-pint, wt. 7 lbs.	\$1.00
Atlas Square Mason Jars, pint, wt. 9 lbs.	1.05
Atlas Square Mason Jars, quart, wt. 12 lbs.	1.20
Atlas Square Mason, ½-gal., wt. 22 lbs.	1.50
Atlas Wide Mouth, pint, wt. 10 lbs.	1.20
Atlas Wide Mouth, quart, wt. 16 lbs.	1.35

Caps for all types jars

	Per doz.
Atlas Seal All Standard Caps, 2-piece, wt. 9 oz.	\$.33
Atlas Seal All Standard Lids only, wt. 4 oz.	.15
63 MM 2-piece Caps for Coffee and Relish Jars, wt. 8 oz.	.33
Rubber Rings for Old Style Zinc Caps, wt. 4 oz.	.10
Wide Mouth Caps (Rings and Lids), wt. 10 oz.	.48
Wide Mouth (Lids Only), wt. 5 oz.	.25

JELLY GLASS

Wt. per Doz., 7 lbs.	Per doz.
No. 100 1/3 pt. size, Tumbler Style	\$.75
No. 25 1/3 pt. size, Ribbed Style	.80
No. 92 ½ pt. size, Tumbler style	.80
No. 3 ½ pt. size, Ribbed Style	.85



HONEY JARS

<i>f.o.b. Plant City</i>	Utility Round per doz.	Queenline per doz.
½-lb., 2 doz. to case, wt. 8 lbs.	\$.55	\$.60
1-lb., 2 doz. to case, wt. 12 lbs.	.65	.80
2-lb., 1 doz. to case, wt. 9 lbs.	.85	1.00
No. 5 Glass Honey Pail, wt. 10 lbs.	1.75	
No. 10 Glass Honey Pail, wt. 14 lbs.	2.00	

Jars for Comb and Liquid Honey

	wt. per case	per doz.
1 lb. Round Wide Mouth 24 to case	14 lbs.	.80
2½ lb. Square Wide Mouth 12 to case	12 lbs.	1.40
5 lb. Square Wide Mouth 6 to case	10 lbs.	2.40



QUEENLINE



Bee-Keepers' Supplies, Bee-Hives and Equipment

Prices subject to change without notice and are f.o.b. Plant City or any Kilgore store.

If you wish your shipment to be made by mail, please add postage when ordering.

Most all Bee Suppliers are exempt from State Tax except Hive Bodies and Covers.

The following is only a partial list of our large and complete line of bee supplies carried in stock in all Kilgore's Florida stores and in Kilgore's mail order department at Plant City, Florida. We can always get whatever a customer wants from the factory if we happen to be out of stock, in which case the price will be f.o.b. factory, Medina, Ohio. 5% discount will be allowed on orders of \$100.00 or more. Shipped direct from factory (average time of delivery two to three weeks).

We buy Bees Wax. Write for price.

Write for a complete catalog of Bee-keepers' supplies.

Brood Foundation—(3-Ply)

8x16¾ in.—7 sheets to pound

		Weight	Price
H-1	10-sheet Carton	3 lbs.	\$ 2.30
H-3	50-sheet Carton	9 lbs.	10.70

Medium Brood

For standard frames

8 1/16 x 16¾ in.—8 sheets to pound

		Weight	Price
H-20	10-sheet Carton	2 lbs.	\$ 1.98
H-22	50-sheet Carton	8 lbs.	9.28

Special Brood Foundation

8 1/16 x 16¾ in.—10 sheets to pound

		Weight	Price
H-32	10-sheet Carton	2 lbs.	\$ 1.50
H-34	50-sheet Carton	7 lbs.	7.25
H-36	200-sheet Carton	33 lbs.	28.70

Medium Brood

For shallow frames

4½ x 16¾ in.—16 sheets to pound

		Weight	Price
H-26	10-sheet Carton	2 lbs.	\$ 1.20
H-28	50-sheet Carton	5 lbs.	5.40

Medium Brood, Crimp-Wired

8½ x 16¾ in.—7 sheets to pound

		Weight	Price
D-350	10-sheet Carton	3 lbs.	\$ 2.35
D-350	50-sheet Carton	10 lbs.	10.75

Medium Brood, Crimp-Wired

4¾ x 16¾ in.—12 sheets to pound

		Weight	Price
D-352	10-sheet Carton	2 lbs.	\$ 1.50
D-352	50-sheet Carton	6 lbs.	6.50

Thin Super

J-1.	1-lb. Carton, 3⅞ x 15¾ in., 28 sheets to lb.	2 lbs.	\$1.70
J-3.	5-lb. Carton, 3⅞ x 15¾ in., 28 sheets to lb.	7 lbs.	7.95
J-22.	1-lb. Carton, 4½ x 16½ in., 22 sheets to lb.	2 lbs.	1.70
J-24.	5-lb. Carton, 4½ x 16½ in., 22 sheets to lb.	7 lbs.	7.95
J-27.	1-lb. Carton, 5 x 16½ in., 22 sheets to lb.	2 lbs.	1.70
J-29.	5-lb. Carton, 5 x 16½ in., 22 sheets to lb.	1 lb.	7.95

Standard 10-Frame Hives

One-story standard hive with aluminum cover

16¾ x 20 in.

Hives in the flat are furnished with nails, tin rabbet punched for nails, reversible bottom boards, metal cover, inner cover. Hoffman frames, and complete directions for nailing.

A-1.	Knocked down without foundation.	
	Wt. 31 lbs.	\$9.95

One-story standard hive, with excelsior (wood) cover

A-6. Knocked down without foundation.

Wt. 27 lbs.	\$8.20
-------------	--------

Shallow 10-Frame Extracting Supers

C-12.	1 KD Empty only, no fittings. Wt. 5 lbs.	\$1.70
C-13.	5 KD Empty only, no fittings. Wt. 28 lbs.	7.75

Beeway 10-Frame Section Supers

D-5.	1 KD Without sections or foundation.	
	Wt. 7 lbs.	\$ 2.90
D-6.	5 KD Without sections or foundation.	
	Wt. 36 lbs.	13.30

List of Inside Fixtures for Supers

E-1.	7 Beeway Section Holders (4¼x4¼x1⅞ in.) KD	Wt. 3 lbs.	\$ 1.00
E-2.	50 Beeway Section Holders (4¼x4¼x1⅞ in.) KD	Wt. 15 lbs.	5.55
E-3.	100 Beeway Section Holders (4¼x4¼x1⅞ in.) KD	Wt. 23 lbs.	10.95
E-15.	10 Beeway Wood Separators, 1/10 in. thick	Wt. 2 lbs.	.50
E-35.	100 Super Springs (3 used to a super)	Wt. 1½ lbs.	1.65

Covers, Metal, 10-Frame

B-1.	1 KD Metal Cover with inner cover	Wt. 11 lbs.	\$ 3.95
B-2.	5 KD Metal Cover with inner cover	Wt. 51 lbs.	17.40

The Excelsior Wood Cover, 10-Frame

B-4.	5 KD Excelsior All-Wood Cover	Wt. 31 lbs.	9.25
------	-------------------------------	-------------	------

Inner Covers, 10-Frame


B-8.	1 KD Inner Cover, without bee escape	Wt. 3 lbs.	1.10
B-9.	5 KD Inner Cover, without bee escape	Wt. 12 lbs.	4.75

Bottom Boards, 10-Frame

B-12.	1 KD Reversible Bottom Board	Wt. 7 lbs.	2.20
B-13.	5 KD Reversible Bottom Board	Wt. 37 lbs.	9.25

Comb Honey Sections

	Mill Run	
F-1	100 Wt. 7 lbs.	\$ 3.75
F-3	500 Wt. 32 lbs.	16.75



Full depth 10-Frame Extracting Supers or Hive Bodies

BEE-KEEPERS' SUPPLIES (Continued)

For information on Beekeeping in Florida, write the Florida Agricultural Experiment Station, Mailing Department, Gainesville, Florida, for free copy of Bulletin No. 151, entitled Beginning Beekeeping.

This publication is illustrated and contains much valuable information for beginners.

Frames


Furnished in the flat with nails. All frames are furnished with solid bottoms unless otherwise specified.

Standard frames. Heavy top bar.	10	50	100
and solid bottom bar, 9½x17½ in	G-1	G-2	G-6
Wt. 46 lbs. per 100	\$2.10	\$8.65	\$16.65

Standard frames with heavy top bars			
2 place bottom bar. Wt. 45 lbs.	G-14	G-15	G-16
per 100	\$2.10	\$8.65	\$16.65

Shallow Extracting Frames, with regular Hoffman top bar, with solid bottom bar. Wt. 38 lbs., per 100	G-25	G-26	
	\$7.75	\$15.25	

Shallow frames, L. T. thick top 2 piece bottom bars	G-24	G-27	
	\$1.95	\$7.75	




L 15

Honey Extractors

L-15	1 2-frame reversing (9½x16 in. pocket)	100	\$ 69.00
L-20	1 20-frame Simplicity	300	210.00
L-25	1 4-frame reversing (9½x16 in. pocket)	215	149.00
L-35	1 3-frame Novice, hand power	52	36.95
L-45	1 45-frame Simplicity	470	335.00

Large Electric Extractor can be supplied upon order



Bee Escape

O-5	Root Double Escape without Board.		
	Wt. 1 oz.		\$.35

Queen Excluder, 10-Frame

Bees pass through rapidly, without hesitation.

10-frame Standard 3 wire and wood,	Each	10	
16¼x20 in. Wt. 4 lbs.	B-30	B-31	
	\$1.70	\$16.25	
10-frame Special 7-wire and wood.	B-35	B-36	
Wt. 4 lbs.	\$1.75	\$16.95	
	B-40	B-41	
Unbound Zinc Excluders	\$1.40	\$13.15	

Wire for Frames

		Wt.	
G-61.	¼-lb. Spool Tinned Wire, 350 ft.	8 oz.	\$.37
G-62.	½-lb. Spool Tinned Wire, 700 ft.	1 lb.	.58
G-63.	1-lb. Spool Tinned Wire, 1400 ft.	2 lb.	.90
G-64.	5-lb. Spool Tinned Wire, 7000 ft.	7 lb.	4.00
N-65.	Spur Wire Imbedder	8 oz.	.45



Bristle Bee Brush

N-85	Bristle Bee Brush. Wt. 4 oz.	\$.65
------	------------------------------	--------

Steel Hive Tool

N-81.	Hive Tool. 10-inch. Wt. 6 oz.	\$1.15
-------	-------------------------------	--------

Queen and Drone Trap (10-frame)

O-8.	Wire Trap, Wt. 2 lbs.	\$1.40
------	-----------------------	--------


Feeders

O-2.	Boardman Feeder, with cap. Wt. 12 oz.	\$.35
------	---------------------------------------	--------

Standard Smoker


N-3	Standard Tin Smoker, 3¼x7 in. Wt. 2 lbs.	\$2.50
-----	--	--------

N-7.	Big Jumbo Tin Smoker. Wt. 3 lbs.	3.95
------	----------------------------------	------



Leather and Drill Bee Gloves

Leather (medium or large)	Wt. 2 lbs.	\$3.85
Drill (small, medium or large)	Wt. 1 lb.	1.55



Veils

N-10.	Root Indestructible Bee Veil. Wt. 1 lb.	\$1.95
N-12.	Delphos Steel Bee Veil. Wt. 1 lb.	2.45
N-15.	Cotton Tulle Veil. Wt. 3 lbs.	1.45

INDEX

	Page		Page		Page
SEED AND PLANTING INFORMATION		Sesbania	46	INSECTICIDE and FUNGICIDE	
Home Gardens for Better Health	2	Shallu	43	DUST COMBINATIONS	
Important Directions to Customers	1	Sorghum	46	HOUSEHOLD and GARDEN PESTICIDES	
Kilgore Stores, Locations	Front Cover	Spinach	29	FUMIGANTS	
Location and Plan of the Vegetable		Squash	30-31	Carbon Di Sulphide	64
Garden	3	Strawberry Plants	47	Pyrenone Grain Protectant	64
Making a Seedbed	3	Sunflower	46	Cyanogas	64
Ordering and Mailing Information	1	Swiss Chard	9	D-D Soil Fumigant	64
Preparation of the Soil for the		Tomato	32-33	Methyl Bromide	64
Vegetable Garden	3	Turnip	35	Nema-Kil	64
Reference Table for Florida		Watermelon	36-37	Soilfume 80-20	64
Vegetable Garden	4	Wheat, Egyptian	43	Vapam	64
Sales Tax Information	2	INSECTICIDE AND		VC. 13 Nemacide	64
Storing and Handling Seed	34	FUNGICIDE INFORMATION		SPREADERS, STICKERS and EMULSIFIERS	
What and When to Sow Corn, Grass,		The Miller Amendment on Pesticide		RODENTICIDES and VERMICIDES	
Clover and Field Crop Seeds in Florida	5	Tolerances	50	Ant Killers	64
What and When to Sow Vegetable		INTRODUCTION	49	Carbolineum	64
Seeds in Florida	5	SANITATION	49	No-Nib'l Rabbit Repellent	64
What and When to Sow Flower		SAFE COMBINATION OF PESTICIDES	51	Rat and Mouse Baits	64
Seeds in Florida	5	DON'TS (PESTICIDES NOT TO USE ON		Roach Killers	64
What Will Your Harvest Be?	34	CERTAIN CROPS)	51	WEED KILLERS	
SEED		SEED TREATMENT	52	LIVE STOCK PREPARATIONS	
Alfalfa	41	PESTICIDE DILUTION TABLE	52	POULTRY PREPARATIONS	
Beans, Bush, Pole, Lima, Green	6-8	STANDARD HOUSEHOLD		POULTRY SUPPLIES	
Beans, Velvet	46	MEASUREMENTS	52	FERTILIZERS	
Beets	9	AMOUNT OF SPRAY AND DUST		SOLUBLE PLANT FOODS	
Beets, Spinach	9	PER ACRE	52	NUTRIENTS	
Beets, Sugar and Stock	9	INSECT AND DISEASE		Activo	68
Bene or Sesame	41	RECOMMENDATIONS	53-58	Adco	68
Broccoli	9	Insects	53-58	Aluminum Sulphate	68
Brussels Sprouts	9	Livestock Pests	58	Calcium Chloride Flakes	68
Cabbage	10-11	Termite Control	58	Es-Min-el	68
Cabbage, Chinese	11	INSECTICIDES		Hydrated Lime	68
Cantaloupe	12	Aldrin 2 Spray	59-60	Kilgore's Improved CMZ	68
Carrots	13	Aramite	59	Kilgore's Iron-Aid	68
Cauliflower	12	Arsenate of Lead	59	Manganese Sulphate	68
Celery	14-15	BENZENE HEXACHLORIDE		Magnesium Sulphate (Epsom Salts)	68
Chicken Corn	43	PREPARATIONS	59	Sulphur-Manganese Dust	68
Chufas	43	Black Leaf 40	59	Zinc Sulphate	68
Clover, Alyce	41	Calcium Arsenate Powder	59	HORMONES	
Clover, White Dutch	42	CHLORDANE PREPARATIONS	59	Fruitone	68
Clover, Hubam	42	Corn Earworm Oil	59	Hormodin Powder	68
Clover, Floranna	42	DDT PREPARATIONS	59	Rootone	68
Clover, Kenland Red	42	Dieldrin Emulsion	59	Transplantone	68
Clover, Ladino	42	Endrin	59	SEED INOCULANTS	
Collard	15	Heptachlor	59	KNAPSACK and HAND DUSTERS	
Corn, Field and Roasting Ear	38	Kelthane	59	RESPIRATORS	
Corn, Sweet	18	KRYOCIDE PREPARATIONS	59	FLAME SPRAYERS	
Cowpeas, Field and Table	42	Lime Sulphur Solution	59	POWER and HAND SPRAYERS	
Crotalaria	43	LINDANE PREPARATIONS (ISOTOX)	59	LAWN SPRINKLERS	
Cucumber	16-17	Malathion	60	FARM and GARDEN TOOLS	
Dill	20	Marlate	60	Cyclone Seeder	72
Eggplant	19	Oil Emulsion	60	Insco Soil Fumigant Applicator	72
Egyptian Wheat	43	Orthal Garden Spray	63	LAWN and GARDEN TOOLS	
Endive	20	Pyrocid Dust	60	MISCELLANEOUS FARM and	
Flower Seed	48	PARATHION PREPARATIONS	60	GARDEN SUPPLIES	
Field Seed	41-46	Paris Green	60	Bamboo Stakes	75
Garlic	20	POISON BAITS	60	Cotton Twine	75
Grass, Lawn	39-40	Red Arrow Insect Spray	60	Hot Kaps	75
Grass, Pasture	41	RHOTHANE PREPARATIONS	60	Jute Twine	75
Gladiolus Bulbs	47	Rotenone Dust	60	Label Paste	75
Hegari (Higear)	45	Scalecide	60	Raffia	75
Hairy Indigo	43	Sixtox	60	Soil Test Kits	75
Kale	20	TOXAPHENE PREPARATIONS	60	WEATHER THERMOMETERS	
Kohl Rabi	20	Vapotone XX Spray	60	CANNING DEPARTMENT	
Leek	20	Volck Oil Spray	60	Can Sealers	76
Lettuce	21	FUNGICIDES		Cans	77
Lupine	43-44	Arsan	61-62	Fruit Jars	77
Millet, Pearl or Cattail	44	Antibiotics Sprays	61	Honey Jars	77
Mustard	22	Ceresan	61	Jelly Glasses	77
Oats	44	COPPER COMPOUNDS	61	Pressure Cookers	76
Okra	23	COPPER A COMPOUNDS PREPARATIONS		BEE-KEEPERS SUPPLIES	
Onion	24	KILCOP "53" PREPARATIONS	61	78-79	
Onion Sets	24	DITHANE PREPARATIONS	61		
Parsley	25	FERMATE PREPARATIONS	61		
Parsnip	25	Karathane	61		
Peanuts	44-55	Manzate	61		
Peas, English or Garden	25	Orthocide	61		
Peas, Southern (Cowpeas)	42	Parzate	61		
Pepper	26-27	Phygon XL	62		
Potatoes	28	Semesan	62		
Pumpkin	28	Semesan Bel	62		
Radish	29	Spergon	62		
Rape, Dwarf Essex	45	Sulfocide	62		
Rhubarb	28	Sulphur	62		
Rice	45	Tersan 75	62		
Rose Bushes	47	Zerlate	62		
Roselle	28				
Rutabaga	35				
Rye	45				
Sage	29				



THE KILGORE SEED COMPANY

PLANT CITY, FLORIDA

Name

P. O. Box, Street,
or R.F.D. No.

Post Office

County State

Ship by ☐ Parcel Post ☐ Express ☐ Truck ☐ Freight

Exp. or Frt. Office _____ Date _____ 19____

AMOUNT ENCLOSED

Check

Money Order..		
---------------	--	--

Cash.....		
-----------	--	--

Total.....		
------------	--	--

IMPORTANT! Be sure address and shipping instructions are correct. Please write plainly.

SPECIAL NOTICE: Remittances should be made by postal money orders, checks, or cash. **DO NOT SEND STAMPS.** On items not marked postpaid be sure your remittance includes the necessary amount for postage and also the "State Sales Tax" on non-exempt merchandise.

Although we have an ample supply of all items listed in this catalog as we go to press, owing to the extra heavy demand for seeds, some items may be short before the end of the season. Therefore we would ask you wherever possible in the case of seeds to allow us to substitute with items of equal value and of the nearest variety. Just mark with a cross in the space provided at the bottom of this order sheet.

We warrant that the seeds, bulbs and plants we sell are, at the time of delivery, as described on the container within recognized tolerances; but we limit our liability on this warranty to the amount of the purchase price of such seeds, bulbs and plants. We give no other or further warranty express or implied.

Notice of any defect or variation resulting from the failure of any seed to be as represented must be given within a reasonable time and not later than 15 days from the time such defect or variation becomes apparent and is observed by the purchaser. THE KILGORE SEED COMPANY. Section 578.26, Florida Statutes.

[illegible]

SUBSTITUTES:

☐ -You may substitute the next best or most similar of any variety you cannot supply.

☐ —Please do not substitute without first notifying me.

(Mark "X" in above desired space)

We would like to send our annual catalog to your friends. Space on the reverse side of this sheet is provided for this purpose.

If you are interested in flower culture and home beautification and wish a copy of **Kilgore's Flower Guide for Florida** be sure to mark X in this space ☐

PLEASE DO NOT FORGET TO GIVE YOUR NAME AND ADDRESS


[SEE OTHER SIDE]

Tear Off Along This Line

Tear Off Along This Line

The names of neighbors or friends who might be interested in receiving a copy of our annual catalog.

NAMES	POST OFFICE	P.O. Box, Street, or R.F.D.	STATE
(SEE OTHER SIDE)			



A PARTIAL VIEW OF KILGORE'S MODERN INSECTICIDE MANUFACTURING PLANT, PLANT CITY, FLA.




PLANT

KILGORE'S
Bred-Rite
SEEDS


TRADE MARK

FOR BETTER CROPS


FOR HIGHER RETURNS




Wilt Resistant Grothen
Globe Tomato



Ft. Myers Market Eggplant



Florida Giant Pepper



Early
Yellow Summer Crookneck
(Baby Crookneck)
Squash



Smith's Perfect Cantaloupe



Marketer
(Early Green Market)
Cucumber

THE KILGORE SEED COMPANY

General Offices and Mail Order Department • • • PLANT CITY, FLORIDA